

Acta Medica Okayama

Volume 58, Issue 4

2004

Article 2

AUGUST 2004

2-methoxyestradiol enhances p53 protein transduction therapy-associated inhibition of the proliferation of oral cancer cells through the suppression of NFkappaB activity.

Hidehiko Takata*

Kazuhiro Tomizawa†

Masayuki Matsushita‡

Hideki Matsui**

*Okayama University,

†Okayama University,

‡Okayama University,

**Okayama University,

2-methoxyestradiol enhances p53 protein transduction therapy-associated inhibition of the proliferation of oral cancer cells through the suppression of NFkappaB activity.*

Hidehiko Takata, Kazuhito Tomizawa, Masayuki Matsushita, and Hideki Matsui

Abstract

Protein transduction therapy using poly-arginine peptide can deliver the biologically active proteins. A previous study showed that 11 poly-arginine fused p53 protein (11R-p53) effectively penetrated across the plasma membrane and inhibited the proliferation of oral cancer cells. However, the intracellular half-life of the delivered protein was less than 36 h. Previous studies also showed that 2-methoxyestradiol (2-ME), an endogenous non-toxic estrogenic metabolite, induces the stabilization of the wild-type p53 protein in human cancer cells posttranscriptionally. In the present study, we examined whether 2-ME induced the stabilization of 11R-p53 and had an inhibitory effect on the proliferation of oral cancer cells. The application of 2-ME significantly enhanced the inhibitory effect of 11R-p53 on the proliferation of oral cancer cells. However, 2-ME had no effect on the intracellular half-life of 11R-p53 in oral cancer cells. Of interest is the finding that 2-ME suppressed the transcriptional activity of NFkappaB, which has an important role in tumorigenesis, but did not affect p53 transcriptional activity. These results suggest that 2-ME synergistically enhances the 11R-p53-induced inhibition of the proliferation of oral cancer cells through the suppression of NFkB transcription.

KEYWORDS: tumor, TAT, poly arginine, gene therapy, protein therapy

*PMID: 15551755 [PubMed - indexed for MEDLINE]

Original Article

2-Methoxyestradiol Enhances p53 Protein Transduction Therapy-Associated Inhibition of the Proliferation of Oral Cancer Cells through the Suppression of NF κ B Activity

Hidehiko Takata, Kazuhito Tomizawa*, Masayuki Matsushita, and Hideki Matsui

*Department of Physiology, Okayama University Graduate School of
Medicine and Dentistry, Okayama 700-8558, Japan*

Protein transduction therapy using poly-arginine peptide can deliver the biologically active proteins. A previous study showed that 11 poly-arginine fused p53 protein (11R-p53) effectively penetrated across the plasma membrane and inhibited the proliferation of oral cancer cells. However, the intracellular half-life of the delivered protein was less than 36 h. Previous studies also showed that 2-methoxyestradiol (2-ME), an endogenous non-toxic estrogenic metabolite, induces the stabilization of the wild-type p53 protein in human cancer cells posttranscriptionally. In the present study, we examined whether 2-ME induced the stabilization of 11R-p53 and had an inhibitory effect on the proliferation of oral cancer cells. The application of 2-ME significantly enhanced the inhibitory effect of 11R-p53 on the proliferation of oral cancer cells. However, 2-ME had no effect on the intracellular half-life of 11R-p53 in oral cancer cells. Of interest is the finding that 2-ME suppressed the transcriptional activity of NF κ B, which has an important role in tumorigenesis, but did not affect p53 transcriptional activity. These results suggest that 2-ME synergistically enhances the 11R-p53-induced inhibition of the proliferation of oral cancer cells through the suppression of NF κ B transcription.

Key words: tumor, TAT, poly arginine, gene therapy, protein therapy

The human immunodeficiency virus (HIV) TAT protein transduction system has been shown to transduce biologically active proteins into cells [1-3]. However, this method has critical disadvantages in terms of its low efficiency of transduction and its influence on neuronal plasticity [4, 5]. To overcome these problems, a novel transduction system utilizing eleven poly-arginine (11R) has been developed [4]. 11R is highly efficient in delivering proteins into cells in multiple organs and is non-toxic with respect to neuronal function [4].

Recent advances in molecular oncology have led to the identification of proteins that are quantitatively, qualitatively or genetically altered in cancer cells. p53 protein is one of the altered proteins in cancer cells, and is mutated or deleted in more than half of human tumors [6]. Therefore, tumor suppressor protein therapy with wild-type p53 protein has been proposed as a possible therapy for malignancies lacking active p53. A previous study showed that 11R-fused p53 protein effectively penetrates across the plasma membrane of oral cancer cells (> 95% cells) and translocates into the nucleus [7]. The protein induces the activity of the p21/WAF1 promoter and inhibits the proliferation of human oral cancer cells, and its effect is equivalent to that of adenovirus-mediated p53

Received December 10, 2003; accepted February 9, 2004.

*Corresponding author. Phone: +81-86-235-7107; Fax: +81-86-235-7111
E-mail: tomikt@md.okayama-u.ac.jp (K. Tomizawa)

gene therapy [7]. However, the half-life of 11R-p53 is less than 36 h [7]. Therefore, repeated transduction of 11R-p53 is needed for the inhibition of oral cancer cell proliferation [7]. Although the immunogenicity of the transduced protein has not been determined, repeated administration of protein therapy may initiate an immune response. The development of protein therapy to transduce stable proteins or of methods to stabilize the delivered proteins is important in order to initiate clinical trials of protein therapy.

2-Methoxyestradiol (2-ME) is a metabolic byproduct of estrogen and an anti-mitotic agent that is present in human urine and blood [8]. 2-ME has been shown to inhibit the growth of various tumor cells through the induction of apoptosis [9, 10]. 2-ME can potently inhibit superoxide dismutase, resulting in enhanced formation of reactive oxygen species and thus toxicity to tumor cells [11]. Also, 2-ME interferes with NF κ B transcriptional activity, which plays an important role in tumorigenesis and which has been shown to be aberrant in many types of cancer cells and primary tumors [12–14]. Moreover, previous studies have shown that 2-ME induces stabilization of the wild-type p53 protein in human cancer cells posttranscriptionally [15, 16], and that adenoviral-vector-mediated p53 gene transfer followed by 2-ME treatment causes significant growth inhibition in human lung cancer cell lines [15]. These results suggest that 2-ME might induce the stabilization of 11R-p53 in oral cancer cells and enhance the inhibitory effect of 11R-p53 on the proliferation of oral cancer cells.

In the present study, we examined whether 2-ME reduced the degradation of 11R-p53 and enhanced the anti-tumor effect of the protein.

Materials and Methods

Cell culture. The oral cell line NOS-1, which was established from a patient with oral squamous cell carcinoma and in which the *p53* gene was point-mutated at site 248, was a gift from Drs. N. Oku and T. Komori (Kobe University, Kobe, Japan). The oral cancer cell line SAS, which contains the wild-type *p53* gene, and the HSC-3 and HSC-4 cell lines were provided by the Cell Resource Center for Biomedical Research (Tohoku University, Sendai, Japan). All cell lines were maintained in GIT medium (Wako Pure Chemical Industries, Osaka, Japan) supplemented with 100 units/ml penicillin and 100 units/ml streptomycin. The cells were

cultured in a 37 °C incubator with 5% CO₂.

Production of 11R-p53 proteins. Production of 11R-p53 was carried out as described previously [7]. Briefly, full-length human wild-type p53 cDNA was subcloned into p11R-HA vector. The plasmids were then transformed into BL21-DE3 *Escherichia coli* cells and 11R-p53 proteins were expressed in the cells by induction with 0.2 mM isopropyl-1-thio- β -D-galactopyranoside. The 11R-p53 proteins were purified using a column of Ni-NTA agarose (Invitrogen, San Diego, CA, USA).

Cell proliferation assay. Cell proliferation was assessed as described previously [7]. Briefly, cells were plated in 96-well plates at a density of 4×10^3 cells/well in quintuplicate. After 24 h, the cells were incubated with 1 μ M 11R-p53 with/without 10 μ M 2-ME (Sigma-Aldrich Chemical, St. Louis, MO, USA). The cell medium supplemented with 10 μ M 2-ME was exchanged every 24 h. Control cells received only the vehicle (0.1% DMSO). Cell growth was assessed every 24 h by measuring the conversion of the triazolium salt WST-1 (Roche Applied Science, Mannheim, Germany) to formazan according to the manufacturer's instructions.

Western blotting analysis. Western blotting analysis for p53 and actin was carried out as described previously [17]. Briefly, the harvested cells were lysed by boiling and sonication containing 1% SDS. Samples containing 50 μ g of total protein were electrophoresed on 10% SDS-PAGE gels and then transferred to nitrocellulose membranes (Hybond ECL; Amersham Biosciences, Piscataway, NJ, USA). The blots were probed with primary antibodies against human p53 (1:1000; Pab 1801; Santa Cruz Biotechnology, Santa Cruz, CA, USA) and actin (1:5000; AC-40; Sigma-Aldrich Chemical, St. Louis, MO, USA). The bands were visualized using a commercial ECL detection kit (Amersham Biosciences).

Reporter assay for p53- and NF κ B-driven transactivation. The reporter assays were performed as described previously [7]. For the p53 reporter assay, the luciferase reporter vector pGL2-basic (Promega, Madison, WI, USA) containing an inserted 2.4-kbp fragment of the human p21/WAF1 promoter was a gift from Drs. T. Akiyama (Tokyo University, Tokyo, Japan) and K. Yoshikawa (Osaka University, Osaka, Japan). pNF κ B-TA-Luc plasmid (BD Biosciences Clontech, Palo Alto, CA, USA) was used for the NF κ B reporter assay. The luciferase reporter vectors were transfected into ~70% confluent NOS-1 cells in

35-mm dishes using Lipofectamine plus (Invitrogen). The cells were then incubated with $1 \mu\text{M}$ 11R-p53 and $10 \mu\text{M}$ 2-ME for 24 h. The cells were harvested, and the luciferase activities were measured with a luminometer using a reagent kit (Tokyo Ink, Tokyo, Japan). The background luciferase activity was subtracted in all experiments.

Statistical analysis. Data were analyzed using either the Student's *t*-test to compare 2 conditions or ANOVA followed by planned comparisons of multiple conditions, and $P < 0.05$ was considered to be significant.

Results

Effect of co-treatment with 11R-p53 and 2-ME on the proliferation of oral cancer cells.

A previous study showed that $1 \mu\text{M}$ 11R-p53 inhibits the proliferation of both SAS and NOS-1 cells, in which wild-type and point-mutated p53 genes are expressed, respectively [7]. In the present study, the oral cancer cells were treated with $1 \mu\text{M}$ 11R-p53 on day 0. After 4 h, the cells were washed and treated with fresh medium in the presence or absence of $10 \mu\text{M}$ 2-ME, which is an optimal concentration for the inhibition of the proliferation of cancer cells [9-11, 14]. Treatment with 11R-p53 significantly suppressed the proliferation of SAS cells (Figs. 1A and C, day 4, control, 1.56 ± 0.1 ; 11R-p53, 1.24 ± 0.15 , $n = 5$ each, $*P < 0.05$) and NOS-1 cells (Figs. 1B and C, day 4, control, 1.68 ± 0.12 ; 11R-p53, 1.22 ± 0.13 , $n = 5$, $***P < 0.01$) as described previously [7]. Co-application of 11R-p53 and 2-ME inhibited the proliferation of the oral cancer cells more markedly than treatment with 11R-p53 alone (Fig. 1A-C, SAS, day 4, 11R-p53 + 2-ME, 0.56 ± 0.08 , $n = 5$, $**P < 0.001$ vs. 11R-p53; NOS-1, day 4, 11R-p53 + 2-ME, 0.55 ± 0.13 , $n = 5$, $**P < 0.001$ vs. 11R-p53). Moreover, the inhibitory effect of treatment with 2-ME on the proliferation was the same as that of 11R-p53 treatment (Fig. 1C, SAS, day 4, 2-ME, 0.91 ± 0.11 , $n = 5$, $P > 0.05$ vs. 11R-p53; NOS-1, day 4, 2-ME, 1.18 ± 0.15 , $n = 5$, $P > 0.05$ vs. 11R-p53). These results suggest that co-treatment with 11R-p53 and 2-ME causes enhancement of the individual effects of these agents on the proliferation of oral cancer cells.

Effect of 2-ME on the stabilization and level of transcription of 11R-p53 in oral cancer cells. To investigate the effect of 2-ME on the stabilization of 11R-p53 in NOS-1 cells, the effect of 2-ME on the time-dependent changes in the level of

11R-p53 were observed. A high level of 11R-p53 was detected 6 h after exogenous addition of the protein to the cells in the absence of 2-ME (Fig. 2A). The level was decreased after 12 h and the low level was maintained until 72 h after 11R-p53 addition. Co-application of $10 \mu\text{M}$




Fig. 1 2-ME induced the inhibitory effect of 11R-p53 on the proliferation of the oral cancer cell lines, SAS (A) and NOS-1 (B). $1 \mu\text{M}$ 11R-p53 with $10 \mu\text{M}$ 2-ME (11R-p53 + 2-ME) or without 2-ME (11R-p53) was added to the cells at day 0. The medium was changed to fresh medium every 24 h, and for 11R-p53 + 2-ME, $10 \mu\text{M}$ 2-ME was added in the medium. Cell growth was assessed every 24 h by WST assay as described in "Materials and Methods". C, Comparison of the inhibitory effect of each treatment on cell growth at day 4. Data are representative of 5 independent experiments. Bars, S.D. $*P < 0.05$, $**P < 0.001$ and $***P < 0.01$.

2-ME with 11R-p53 had no effect on the level of 11R-p53 (Fig. 2A). The level of 11R-p53 was high 6 h after it was added together with 2-ME, but the high level was not maintained 12 h after the addition. The level of 11R-p53 72 h after the addition was not different than the level observed in the absence of 2-ME. The effect of high-dose 2-ME (50 μ M) was also examined, but 50 μ M 2-ME did not affect the stabilization of 11R-p53 (data not shown).

The effect of 2-ME on the expression of endogenous p53 was next examined in various oral cancer cell lines. HSC-3 cells harbor an insertion mutation between codons 305 and 306. The p53 protein is prematurely terminated and the predicted molecular weight is 42 kDa. In HSC-4 cells, a heterozygous missense mutation corresponding to a CAG (Gln) \rightarrow CGG (Arg) transition in codon 248 was seen in the p53 gene. Each cell line was incubated with




Fig. 2 A, Time course of the degradation of the 11R-p53 proteins and the effect of 2-ME on the degradation in NOS-1 cells. After incubation with 11R-p53 for 4 h, the cells were washed with PBS twice and were then incubated with fresh medium in the presence or absence of 2-ME (10 μ M). The medium was replaced with fresh medium with/without 2-ME (10 μ M) every 24 h. The cells were collected at the indicated time-points, and Western blotting analysis was performed using anti-p53 and anti-actin antibodies. B, Effect of 2-ME on endogenous p53 expression in oral cancer cell lines. Each cell line was incubated with 10 μ M 2-ME for 48 h. The cells were then collected and Western blotting analysis was performed using anti-p53 and anti-Actin antibodies. The predicted molecular weight of the p53 protein in HSC-3 cells is 42 kDa. C, Effect of 2-ME (10 μ M) on p53 reporter activity. Data are representative of 5 independent experiments. Data represent means \pm S.D. * P < 0.0001.

11R-p53 for 48 h and was then collected (Fig. 2B). 2-ME did not induce the expression of endogenous p53 in any of the cell lines. These results suggest that 2-ME had no effect on the expression or stabilization of 11R-p53 and p53 in oral cancer cells.

We also examined the effect of 2-ME on the level of transcriptional activity of 11R-p53 in NOS-1 cells (Fig. 2C). Transduction of 11R-p53 induced the p53 transcriptional activity (23.5 ± 1.7 fold compared with the control, $n=6$ each, $*P < 0.0001$). The p53 transcriptional activity in the cells incubated with both 11R-p53 and 2-ME was the same as that in 11R-p53-transduced cells (11R-p53 + 2-ME, 21 ± 3.7 fold, $n=6$, $P > 0.05$ vs. 11R-p53), indicating that 2-ME had no effect on the 11R-p53 transcriptional activity. Moreover, the application of 2-ME did not induce endogenous p53 transcriptional activity without 11R-p53 (2-ME, 1.7 ± 0.9 , $n=6$, $P > 0.05$ vs. control).

Inhibitory effect of 2-ME on the NF κ B transcriptional activity. Previous studies have shown that 2-ME interferes with NF κ B transcriptional activity, resulting in the inhibition of the proliferation of cancer cells and the induction of apoptosis of the cells [12-14]. The effects of 11R-p53 and 2-ME on the level of NF κ B transcriptional activity in NOS-1 cells were examined (Fig. 3). A high level of NF κ B activity was observed in the intact cells, and 11R-p53 protein transduction had no effect on the level of transcriptional activity

(1.1 ± 0.1 , $n=5$, $P > 0.05$ vs. Cont.). In contrast, co-application of 11R-p53 and 2-ME significantly reduced the level of NF κ B transcriptional activity (0.23 ± 0.12 , $n=5$, $*P < 0.001$ vs. Cont.). Application of 2-ME alone also inhibited the level of NF κ B activity to the same extent as co-application of 11R-p53 and 2-ME (Fig. 3, 0.21 ± 0.05 , $n=5$, $*P < 0.001$ vs. Cont.). These results suggest that 2-ME inhibits the NF κ B transcriptional activity in the oral cancer cells.

Discussion

The results of the present study provided the three following important findings: 1) Co-application of 11R-p53 and 2-ME significantly inhibited the proliferation of oral cancer cells compared with the application of 2-ME alone; 2) 2-ME had no effect on the stabilization or expression of either 11R-p53 or endogenous p53 protein in oral cancer cells; 3) 2-ME reduced the NF κ B transcriptional activity but had no effect on the p53 transcriptional activity.

Protein transduction therapy using agents such as poly-arginine and TAT peptides is useful for the delivery of biologically active proteins. However, protein transduction therapy has a disadvantage in terms of the short-term expression of the delivered proteins. The present and previous studies show that the intracellular half-life of the delivered 11R-p53 is less than 24 h [7]. Previous studies have shown that 2-ME induces posttranscriptional stabilization of the wild-type p53 protein in human cancer cells [15, 16], and that adenoviral-vector-mediated p53 gene transfer followed by 2-ME treatment causes significant growth inhibition in human lung cancer cell lines [15]. In the present study, we investigated whether 2-ME enhanced the stabilization and expression of 11R-p53 in oral cancer cells. However, 2-ME had no effect on the stabilization or expression of either 11R-p53 or endogenous p53 protein in oral cancer cells. It is hard to explain the discrepancy between the previous and present studies because the mechanism of 2-ME-dependent induction of the stabilization of p53 protein is unclear. One possibility is that the origin of the cancer cell lines was different. Previous studies have shown the effect of 2-ME on the induction of p53 protein in human lung cancer cell lines and colorectal carcinoma cell lines [15, 16]. A method to stabilize the transduced 11R-p53 in oral cancer cells needs to be developed. A previous study showed that a p53 mutant in which lysine residues




Fig. 3 Effect of 11R-p53 and 2-ME on NF κ B transcriptional activity in NOS-1 cells. The cells were transfected with pNF B-TA-Luc plasmid. After 24 h, the cells were incubated with $1 \mu\text{M}$ 11R-p53 and/or $10 \mu\text{M}$ 2-ME for 24 h and were then harvested, and the luciferase activities were measured. Data represent means \pm S.D. $*P < 0.001$

370, 372, 373, 381, 382 and 386 were replaced by arginine residues was resistant to ubiquitin-proteasome-mediated degradation, and that the transcriptional activity was higher than that of wild-type p53 [18]. This result suggests that stable expression of the fusion protein of the p53 mutant with 11R is maintained in oral cancer cells. Another possible method of stabilization is the transduction of a phosphorylated form of 11R-p53. Phosphorylation of p53 at Ser15 and/or Ser20 induces stabilization of p53 by inhibiting of the binding with MDM2, which is critical for ubiquitin-proteasome-mediated degradation [19]. Direct addition of phospho-11R-p53, which was previously phosphorylated *in vitro*, may solve the problem of the stabilization of 11R-p53.

In the present study, the growth of SAS cells with wild-type p53 gene was suppressed by wild-type p53 protein transduction. The expression of p53 is regulated by the action of Mdm2, an oncogenic E3 ligase, to induce p53 ubiquitination and degradation [20, 21]. The human Mdm2 is overexpressed in a wide range of human cancers in which wild-type p53 is expressed, and the overexpression of Mdm2 serves to inactivate p53 function in these cells [22, 23]. These results may suggest that Mdm2 overexpresses in SAS cells and induces the rapid degradation of p53. Therefore, 11R-p53 transduction may effectively inhibit the proliferation of the cells.

Interestingly, the present results showed that 2-ME inhibited the NF κ B transcriptional activity. Numerous studies have examined the tumor-inhibiting potential of 2-ME [8-11, 14, 24]. Moreover, co-application of adenoviral-vector-mediated p53 gene transfer and 2-ME causes significant growth inhibition in human lung cancer cell lines [15]. However, the exact mechanism by which 2-ME inhibits cancer cell proliferation is unknown. A very recent report showed that 2-ME preferentially inhibits the growth of brain tumor cells by blocking cell cycle progression in the G₂/M phase and induces apoptosis of the tumor cells [14]. In addition, 2-ME reduces the transcriptional activity and DNA binding activity of NF κ B but does not influence the expression of p53 [14]. The present results agree with those results, and show that 2-ME did not alter p53 expression or a p53 transcriptional activity. NF κ B has been suggested to have a wide-ranging role in tumor progression, and its activity is associated with angiogenesis and metastasis [12, 13]. Also it is known to be required for the transforming properties of many oncogenes, such as Bcr-Abl, Ras and the human T cell lymphotropic virus

1 (HTLV 1) protein Tax [25-28]. These results suggest that NF κ B activation may be crucial for cancer cell growth and that 2-ME may inhibit cell growth through the inhibition of NF κ B activation, not through a p53-mediated pathway in oral cancer cells.

In contrast, we previously showed that 11R-p53 inhibited the proliferation of oral cancer cells without any significant induction of apoptosis [7]. Moreover, 11R-p53 induces the activity of the promoter of p21/WAF, which mediates G1 arrest via its inhibitory effects on the cyclin-dependent kinases (CDKs) required for S-phase entry [29, 30]. These results may suggest that 11R-p53 inhibits the growth of oral cancer cells by means of the induction of cell-cycle arrest, and that the mechanism of the growth inhibition induced by 11R-p53 is different from that induced by 2-ME. Therefore, co-application of 11R-p53 and 2-ME might cause a significant additive inhibitory effect on the proliferation of oral cancer cells compared with the independent application of each agent alone in the present study. 2-ME is a natural metabolic by-product of estrogen and causes no adverse reactions such as hair loss, intestinal damage or infection [31]. These results suggest that co-application of 11R-p53 and 2-ME has great potential as a chemotherapeutic treatment for oral cancer.

In conclusion, 2-ME had no effect on the stabilization or expression of 11R-p53, but significantly inhibited oral cancer proliferation. Our current findings may be useful for the development of a novel oral cancer therapy.

Acknowledgments. This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas "Cancer" from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Schwarze SR, Ho A, Vocero-Akbani A and Dowdy SF: *In vivo* protein transduction: Delivery of a biologically active protein into the mouse. *Science* (1999) 285: 1569-1572.
- Schwarze SR and Dowdy SF: *In vivo* protein transduction: Intracellular delivery of biologically active proteins, compounds and DNA. *Trends Pharmacol Sci* (2000) 21: 45-48.
- Matsui H, Tomizawa K, Lu YF and Matsushita M: Protein Therapy: *In vivo* protein transduction by polyarginine (11R) PTD and subcellular targeting delivery. *Curr Protein Pept Sci* (2003) 4: 151-157.
- Matsushita M, Tomizawa K, Moriwaki A, Li ST, Terada H and Matsui H: A high-efficiency protein transduction system demonstrating the role of PKA in long-lasting long-term potentiation. *J Neurosci* (2001) 21: 6000-6007.
- Li ST, Matsushita M, Moriwaki A, Lu YF, Tomizawa K and Matsui H: Human immunodeficiency virus type 1 Tat potentiates extrasynaptic N-methyl-D-Aspartate receptor-mediated calcium influx and attenuates

- spatial learning. *Annals Neurol* (2004): in press.
6. Greenblatt MS, Bennett WP, Hollstein M and Harris CC: Mutations in the p53 tumor suppressor gene: Clues to cancer etiology and molecular pathogenesis. *Cancer Res* (1994) 54: 4855-4878.
 7. Takenobu T, Tomizawa K, Matsushita M, Li ST, Moriwaki A, Lu YF and Matsui H: Development of p53 protein transduction therapy using membrane-permeable peptides and the application to oral cancer cells. *Mol Cancer Ther* (2002) 1: 1043-1049.
 8. Zhu BT and Conney AH: Is 2-methoxyestradiol an endogenous estrogen metabolite that inhibits mammary carcinogenesis? *Cancer Res* (1998) 58: 2269-2277.
 9. Schumacher G, Kataoka M, Roth JA and Mukhopadhyay T: Potent antitumor activity of 2-methoxyestradiol in human pancreatic cancer cell lines. *Clin Cancer Res* (1999) 5: 493-499.
 10. Ghosh R, Ott AM, Seetharam D, Slaga TJ and Kumar AP: Cell cycle block and apoptosis induction in a human melanoma cell line following treatment with 2-methoxyestradiol: Therapeutic implications? *Melanoma Res* (2003) 13: 119-127.
 11. Wood L, Leese MR, Leblond B, Woo LW, Ganeshpillai D, Purohit A, Reed MJ, Potter BV and Packham G: Inhibition of superoxide dismutase by 2-methoxyestradiol analogues and oestrogen derivatives: Structure-activity relationships. *Anticancer Drug Des* (2001) 16: 209-215.
 12. Baldwin AS: Control of oncogenesis and cancer therapy resistance by the transcription factor NF-kappaB. *J Clin Invest* (2001) 107: 241-246.
 13. Garg A and Aggarwal BB: Nuclear transcription factor-kappaB as a target for cancer drug development. *Leukemia* (2002) 16: 1053-1068.
 14. Kumar AP, Garcia GE, Orsborn J, Levin VA and Slaga TJ: 2-Methoxyestradiol interferes with NF kappa B transcriptional activity in primitive neuroectodermal brain tumors: Implications for management. *Carcinogenesis* (2003) 24: 209-216.
 15. Mukhopadhyay T and Roth JA: Superinduction of wild-type p53 protein after 2-methoxyestradiol treatment of Ad5p53-transduced cells induces tumor cell apoptosis. *Oncogene* (1998) 17: 241-246.
 16. Carothers AM, Hughes SA, Ortega D and Bertagnolli MM: 2-Methoxyestradiol induces p53-associated apoptosis of colorectal cancer cells. *Cancer Lett* (2002) 187: 77-86.
 17. Tomizawa K, Iga N, Lu YF, Moriwaki A, Matsushita M, Li ST, Miyamoto O, Itano T and Matsui H: Oxytocin improves long-lasting spatial memory during motherhood through MAP kinase cascade. *Nat Neurosci* (2003) 6: 384-390.
 18. Rodriguez MS, Desterro JM, Lain S, Lane DP and Hay RT: Multiple C-terminal lysine residues target p53 for ubiquitin-proteasome-mediated degradation. *Mol Cell Biol* (2000) 20: 8458-8467.
 19. Ashcroft M, Taya Y and Vousden KH: Stress signals utilize multiple pathways to stabilize p53. *Mol Cell Biol* (2000) 20: 3224-3233.
 20. Haupt Y, Maya R, Kazanietz A and Oren M: Mdm2 promotes the rapid degradation of p53. *Nature* (1997) 387: 296-299.
 21. Kubbutat MH, Jones SN and Vousden KH: Regulation of p53 stability by Mdm2. *Nature* (1997) 387: 299-303.
 22. Reifenberger G, Liu L, Ichimura K, Schmidt EE and Collins VP: Amplification and overexpression of the MDM2 gene in a subset of human malignant gliomas without p53 mutations. *Cancer Res* (1993) 53: 2736-2739.
 23. Bueso-Ramos CE, Manshoury T, Haidar MA, Yang Y, McCown P, Ordonez N, Glassman A, Sneige N and Albitar M: Abnormal expression of MDM-2 in breast carcinomas. *Breast Cancer Res Treat* (1996) 37: 179-188.
 24. Carothers AM, Hughes SA, Ortega D and Bertagnolli MM: 2-Methoxyestradiol induces p53-associated apoptosis of colorectal cancer cells. *Cancer Lett* (2002) 187: 77-86.
 25. Mayo MW, Wang CY, Cogswell PC, Rogers-Graham KS, Lowe SW, Der CJ and Baldwin AS Jr: Requirement of NF-kappaB activation to suppress p53-independent apoptosis induced by oncogenic Ras. *Science* (1997) 278: 1812-1815.
 26. Mosialos G: The role of Rel/NF-kappa B proteins in viral oncogenesis and the regulation of viral transcription. *Semin Cancer Biol* (1997) 8: 121-129.
 27. Reuther JY, Reuther GW, Cortez D, Pendergast AM and Baldwin AS Jr: A requirement for NF-kappaB activation in Bcr-Abl-mediated transformation. *Genes Dev* (1998) 12: 968-981.
 28. Jo H, Zhang R, Zhang H, McKinsey TA, Shao J, Beauchamp RD, Ballard DW and Liang P: NF-kappa B is required for H-ras oncogene induced abnormal cell proliferation and tumorigenesis. *Oncogene* (2000) 19: 841-849.
 29. Brugarolas J, Chandrasekaran C, Gordon JI, Beach D, Jacks T and Hannon GJ: Radiation-induced cell cycle arrest compromised by p21 deficiency. *Nature* (1995) 377: 552-557.
 30. Waldman T, Kinzler KW and Vogelstein B: p21 is necessary for the p53-mediated G1 arrest in human cancer cells. *Cancer Res* (1995) 55: 5187-5190.
 31. Klauber N, Parangi S, Flynn E, Hamel E and D'Amato RJ: Inhibition of angiogenesis and breast cancer in mice by the microtubule inhibitors 2-methoxyestradiol and taxol. *Cancer Res* (1997) 57: 81-86.