

*Physics*  
*Light & Optics fields*

---

Okayama University

*Year* 2006

---

Cyclic phase in f=2 spinor  
condensate:long-range order, kinks, and  
roughening transition

W V. Pogosov  
Okayama University

K Machida  
Okayama University

This paper is posted at eScholarship@OUDIR : Okayama University Digital Information Repository.

[http://escholarship.lib.okayama-u.ac.jp/light\\_and\\_optics/7](http://escholarship.lib.okayama-u.ac.jp/light_and_optics/7)

# Cyclic phase in $F=2$ spinor condensate: Long-range order, kinks, and roughening transition

W. V. Pogosov and K. Machida

Department of Physics, Okayama University, Okayama 700-8530, Japan

(Received 23 April 2006; published 18 August 2006)

We study the effect of thermal fluctuations on homogeneous infinite Bose-Einstein condensate with spin  $F=2$  in the cyclic state, when atoms occupy three hyperfine states with  $m_F=0, \pm 2$ . We use both the approach of small-amplitude oscillations and mapping of our model on the sine-Gordon model. We show that thermal fluctuations lead to the existence of the rough phase in one- and two-dimensional systems, when the presence of kinks is favorable. The structure and energy of a single kink are found. We also discuss the effect of thermal fluctuations on spin degrees of freedom in  $F=1$  condensate.

DOI: [10.1103/PhysRevA.74.023611](https://doi.org/10.1103/PhysRevA.74.023611)

PACS number(s): 03.75.Kk

## I. INTRODUCTION

Recent progress in the experimental physics of cold atoms allows for the creating of quasi-two-dimensional atomic gas either using one-dimensional optical lattices or applying tight axial trapping [1–4]. Many properties of low-dimensional systems differ from that for the three-dimensional ones. For instance, it is well known that Bose-Einstein condensation is impossible in the two-dimensional (2D) situation at any non-zero temperature, since long-wavelength thermal fluctuations destroy phase coherence. However, at low temperatures there is still a quasi-long-range order, which disappears at the temperature of Berezinskii-Kosterlitz-Thouless (BKT) transition. The low-temperature phase is often called quasicondensate, and this concept was introduced in the context of physics of alkali-metal atom gases in Refs. [5,6]; see also Ref. [7]. It was shown in those works that, in *finite* low-dimensional systems, there is still a long-range order at low temperatures, and one has a true condensate, but at higher temperatures coherence is again lost, and we have a quasicondensate.

The aim of the present paper is to study thermal fluctuations in homogeneous *spinor* condensates. We are mainly interested in the cyclic phase of the  $F=2$  condensate, since it has unusual properties such as phase locking and kinks, which are absent in other phases of the  $F=2$  condensate. In the recent experiments [8–12],  $F=2$  spinor Bose-Einstein condensates were created and studied. The order parameter in the  $F=2$  system has five components, and, in the cyclic phase, all the particles populate three hyperfine states with  $m_F=0, \pm 2$ . The characteristic feature of the cyclic phase is the fact that the ground-state energy depends on the relative angle  $\chi$  among the phases  $S_j$  of different components of the order parameter,  $\chi=2S_0-S_2-S_{-2}$ , through the spin-mixing term. This leads to the peculiar phase-locking phenomena, since the energy has a term proportional to  $\cos \chi$ . In this paper, we concentrate on thermal fluctuations of phases  $S_j$  of different components of the order parameter as well as of  $\chi$ . First, we solve the Bogoliubov–de Gennes equations and calculate the mean-square fluctuations of  $S_j$ . Each individual phase  $S_j$  in this case behaves similarly to the case of the scalar system. Namely, there is no long-range order for  $S_j$  in one and two dimensions, whereas this order is kept for the 3D situation. At the same time, the long-range order is still preserved for  $\chi$ . After this, we note that the Bogoliubov–de

Gennes equations can be insufficient for the analysis of thermal fluctuations in the cyclic state, since this approach uses an expansion of the energy in the vicinity of one of the infinite number of equivalent minima at  $\chi=2\pi l$ ,  $l$  being an integer number, and does not take into account a global periodic structure of the energy in functional space. In other words, the  $\cos \chi$  contribution to the energy is changed by the quadratic potential well with *infinite* height. One of the possible solutions of the Gross-Pitaevskii equations for the cyclic phase is a *kink*, which separates two spatial domains with  $\chi$  different by  $2\pi$  from each other. At zero temperature, kink is energetically unfavorable. However, at finite temperature the formation of kinks can become favorable due to entropic reasons. We determine analytically the structure and energy of a single kink. Also we note that our model can be mapped on the well-known sine-Gordon model. For this model, it was established before that, in low dimensions, a so called roughening transition can occur, when the system becomes unlocked from one of the minima of  $\chi$ . In the 1D situation, a long-range order for  $\chi$  is absent for any nonzero temperature due to thermally excited kinks, and the system is rough. If the system has a finite length, a nonzero temperature appears, below which there is still a long-range order for  $\chi$ . We estimate this temperature from simple entropic arguments. In two dimensions, a roughening transition occurs at some finite nonzero temperature. From known results for the sine-Gordon model and some simple qualitative considerations, we show that the temperature of the roughening transition is close to the BKT critical temperature and possibly coincides with it. We also discuss briefly the case of the ferromagnetic  $F=1$  condensate. Based on the solutions of the Bogoliubov–de Gennes equations, we find that there is no long-range order in the one- and two-dimensional systems for the direction of spin. In finite systems, the order still exists at low temperatures, similarly to the phase coherence in scalar condensates.

The paper is organized as follows. In Sec. II, we solve the Bogoliubov–de Gennes equations for the  $F=2$  cyclic phase and find correlation functions. In Sec. III, we determine the structure and energy of a single kink. In Sec. IV, we discuss the order-disorder transition due to the proliferation of kinks. In Sec. V, we study the case of the ferromagnetic  $F=1$  condensate. We conclude in Sec. VI.

## II. CYCLIC PHASE: SMALL-AMPLITUDE OSCILLATIONS

We consider the infinite homogeneous  $F=2$  condensate with a given density of particles  $n$  in zero magnetic field. The energy of the system depends on three interaction parameters  $\alpha$ ,  $\beta$ , and  $\gamma$ , which can be defined as [13,14]

$$\alpha = \frac{1}{7}(4g_2 + 3g_4), \quad (1)$$

$$\beta = -\frac{1}{7}(g_2 - g_4), \quad (2)$$

$$\gamma = \frac{1}{5}(g_0 - g_4) - \frac{2}{7}(g_2 - g_4), \quad (3)$$

where  $(q=0, 2, 4)$

$$g_q = \frac{4\pi\hbar^2}{m}a_q \quad (4)$$

and  $a_q$  is the scattering lengths characterizing collisions between atoms with the total spin 0, 2, and 4. In real atomic condensates,  $\alpha \gg \beta, \gamma$ .

The order parameter in the  $F=2$  case has five components  $\Psi_i (i=-2, -1, 0, 1, 2)$ . The energy of the system is given by [15,16]

$$\begin{aligned} F = \int d\mathbf{r} \left[ -\frac{\hbar^2}{2m} \Psi_j^* \Delta \Psi_j + \frac{\alpha}{2} \Psi_j^* \Psi_k^* \Psi_j \Psi_k \right. \\ \left. + \frac{\beta}{2} \Psi_j^* \Psi_l^* (F_a)_{jk} (F_a)_{lm} \Psi_k \Psi_m \right. \\ \left. + \frac{\gamma}{2} \Psi_j^* \Psi_k^* \Psi_{-j} \Psi_{-k} (-1)^j (-1)^k \right], \quad (5) \end{aligned}$$

where integration is performed over the system volume, repeated indices are summed, and  $F_a$  ( $a=x, y, z$ ) is the angular momentum operator, which can be expressed in a usual matrix form.

In the absence of magnetic field and rotation, the condensate can be in three different states [13], as seen from Eq. (5). These states are called ferromagnetic, cyclic, and polar [13]. In the cyclic phase,  $\Psi_{\pm 1}=0$  and  $\Psi_{-2}=\frac{\sqrt{n}}{2}e^{i\theta}$ ,  $\Psi_0=\frac{\sqrt{n}}{2}$ , and  $\Psi_2=-\frac{\sqrt{n}}{2}e^{-i\theta}$ , where  $\theta$  is an arbitrary phase (the energy of the system is degenerate with respect to  $\theta$ ). The ground-state value of  $\chi$  in these notations is zero. Depending on the values of scattering lengths  $a_q$ , the ferromagnetic, cyclic, or polar phase has the lowest energy [13]. Extended Gross-Pitaevskii equations can be obtained as usual from the condition of the minimum of the free energy of the system Eq. (5). In the cyclic state, the last term on the right-hand side of Eq. (5) contains a spin-mixing contribution,

$$F_{\text{sm}} = \gamma |\Psi_0|^2 |\Psi_2| |\Psi_{-2}| \cos \chi. \quad (6)$$

Now we analyze small-amplitude oscillations of the order parameter in the cyclic phase. The fluctuations of phase of the given component of the order parameter can be expressed through the fluctuations of the order parameter itself as  $\delta S_j = \text{Im}(\delta \Psi_j) / |\Psi_j|$ . For  $\delta \chi$ , we have  $\delta \chi = 2\delta S_0 - \delta S_2 - \delta S_{-2}$ .

The deviations of the five components of the order parameter from their equilibrium values  $\delta \Psi_j$  can be represented as

$$\delta \Psi_j = \sum_{\mathbf{k}} \{c_{\mathbf{k}}^{(j)} \exp(i\mathbf{k} \cdot \mathbf{r} + i\omega_{\mathbf{k}} t) + d_{\mathbf{k}}^{(j)*} \exp(-i\mathbf{k} \cdot \mathbf{r} - i\omega_{\mathbf{k}} t)\}, \quad (7)$$

where  $c_{\mathbf{k}}^{(j)}$  and  $d_{\mathbf{k}}^{(j)}$  are constants to be found. Fluctuations of  $\Psi_{\pm 1}$  are decoupled from that for  $\Psi_0, \Psi_{\pm 2}$  as follows for the Bogoliubov–de Gennes equations for the cyclic phase. The spectrum for oscillations of  $\Psi_0, \Psi_{\pm 2}$  has three branches. It was obtained for the first time in Ref. [17], and our results coincide with those results,

$$\hbar \omega_{\mathbf{k}}^{(1)} = \frac{\hbar^2 \mathbf{k}^2}{2m} + 2\gamma n, \quad (8)$$

$$\hbar \omega_{\mathbf{k}}^{(2)} = \sqrt{\left(\frac{\hbar^2 \mathbf{k}^2}{2m}\right)^2 + 2\alpha n \frac{\hbar^2 \mathbf{k}^2}{2m}}, \quad (9)$$

$$\hbar \omega_{\mathbf{k}}^{(3)} = \sqrt{\left(\frac{\hbar^2 \mathbf{k}^2}{2m}\right)^2 + 4\beta n \frac{\hbar^2 \mathbf{k}^2}{2m}}. \quad (10)$$

The thermal distribution of quasiparticles is given by ( $j=1, 2, 3$ )

$$N_{\mathbf{k}}^{(j)} = \frac{1}{\exp(\hbar \omega_{\mathbf{k}}^{(j)} / k_B T) - 1}. \quad (11)$$

One can also find eigenvectors corresponding to the branches (8)–(10). Eigenvectors for Eqs. (9) and (10) give no contribution to  $\delta \chi$ , as can be seen from the solutions of the Bogoliubov–de Gennes equations. This is the first branch (8), which is responsible for the fluctuations of  $\chi$ . Physically, this is due to the fact that only branch (8) depends on the value of  $\gamma$ , which yields the potential well for  $\chi$  via the spin-mixing term (6). Finally, we obtain a simple expression for  $\delta \chi$ ,

$$\delta \chi = \frac{1}{2i\sqrt{V}\sqrt{n}} \sum_{\mathbf{k}} \{\exp(i\mathbf{k} \cdot \mathbf{r} + i\omega_{\mathbf{k}} t) - \exp(-i\mathbf{k} \cdot \mathbf{r} - i\omega_{\mathbf{k}} t)\}. \quad (12)$$

Next we can determine the behavior of the mean-square fluctuations of  $\chi$  at large distances. We found that this quantity in all dimensions tends to a constant, when the distance tends to infinity. This means that there is a long-range order for  $\chi$ . The reason is the presence of the gap in the excitation energy (8). As an example, we present below the derivation and results for the 2D situation. From Eq. (12), we have

$$[\delta \chi(r) - \delta \chi(0)]^2 = -\frac{1}{4Vn} \sum_{\mathbf{k}_1, \mathbf{k}_2} A_{\mathbf{k}_1} A_{\mathbf{k}_2}, \quad (13)$$

where

$$\begin{aligned} A_{\mathbf{k}} &= \exp(i\omega_{\mathbf{k}}^{(1)} t) \\ &\times [\exp(i\mathbf{k} \cdot \mathbf{r}) - 1] - \exp(-i\omega_{\mathbf{k}}^{(1)} t) [\exp(-i\mathbf{k} \cdot \mathbf{r}) - 1]. \end{aligned}$$

After averaging over the time, only terms with  $k_1=k_2$  survive in the expansion (13) and for the correlator we have

$$\langle [\delta\chi(\mathbf{r}) - \delta\chi(0)]^2 \rangle_T = -\frac{1}{8\pi V n} \sum_{\mathbf{k}, \varphi_1, \varphi_2} N_{\mathbf{k}}^{(j)} \times (e^{ikr \cos \varphi_1} - 1)(e^{ikr \cos \varphi_2} - 1), \quad (14)$$

where  $\varphi_1$  and  $\varphi_2$  are angles between  $\mathbf{r}$  and  $\mathbf{k}_1$  and  $\mathbf{k}_2$ , respectively. Now we can switch from the integration to the summation, and after the integration over  $\varphi_1$  and  $\varphi_2$  we get

$$\langle [\delta\chi(\mathbf{r}) - \delta\chi(0)]^2 \rangle_T = \frac{1}{2\pi n} \int k dk N_{\mathbf{k}}^{(1)} [1 - J_0(kr)]^2, \quad (15)$$

where  $J_0(r)$  is the Bessel function. This integral can be evaluated analytically at large  $r$  and in the limits of low and high temperatures. At low temperatures,  $k_B T \ll \gamma n$ ,  $N_{\mathbf{k}}^{(1)} = \exp(-\hbar \omega_{\mathbf{k}}^{(j)} / k_B T)$ . At high temperatures,  $k_B T \gg \gamma n$ ,  $N_{\mathbf{k}}^{(1)} = k_B T / \hbar \omega_{\mathbf{k}}^{(j)}$ . Finally, we have

$$\langle [\delta\chi(\mathbf{r}) - \delta\chi(0)]^2 \rangle_T = \begin{cases} T/T_d \exp(-2\gamma n / k_B T), & k_B T \ll \gamma n, \\ T/T_d \ln(k_B T / 2\gamma n), & k_B T \gg \gamma n, \end{cases} \quad (16)$$

where  $T_d = 2\pi \hbar^2 n / k_B m$  is the temperature of quantum degeneracy.

The second and third branches of the spectrum, given by Eqs. (9) and (10), are responsible for the fluctuations of phases of individual components of the order parameter,  $S_0$ ,  $S_{\pm 2}$ . Our calculations of the mean-square fluctuations of these phases revealed the behavior that is similar to the scalar condensate. The long-range order is absent in one- and two-dimensional situations. It is interesting to note that, at the same time, the long-range order exists for the linear combination of these phases,  $\chi$ . Fluctuations of populations of different hyperfine states are small at  $T \ll T_d$ .

### III. KINK: STRUCTURE AND ENERGY

In the previous section we studied small-amplitude oscillations in the vicinity of a homogeneous solution to the extended Gross-Pitaevskii equations. However, these equations also have a nonhomogeneous solution corresponding to the kink or domain wall between two spatial regions with values of  $\chi$  different by  $2\pi l$  from each other, since the energy of the system is degenerate with respect to  $l$ . Kinks are known to play an important role in the physics of low-dimensional systems, see, e.g., the textbook [18]. Now we find the structure and energy of the one-dimensional kink in the cyclic phase for the most important case, when  $l=1$ . We assume that all the quantities depend only on one coordinate  $x$ . For  $\chi$  we have a boundary conditions  $\chi(\infty) = 2\pi$ ,  $\chi(-\infty) = 0$ . The phase-dependent part of the energy consists of two contributions: one is the kinetic energy, and another one is the spin-mixing term (6),




FIG. 1. The structure of a single kink in the cyclic state. Solid curve denotes  $\chi$ , dashed one corresponds to  $S_0$ , and dotted curve shows  $S_2 = S_{-2}$ .

$$F_{ph} = \int_{-\infty}^{+\infty} dx \left[ \frac{\hbar^2}{2m} \sum_{j=-1}^1 |\Psi_{2j}|^2 (\nabla S_{2j})^2 + \gamma |\Psi_0|^2 |\Psi_2| \times |\Psi_{-2}| \cos \chi \right]. \quad (17)$$

We can rewrite this expression in the diagonal form as

$$F_{ph} = \int_{-\infty}^{+\infty} dx \left[ \frac{\hbar^2 n}{8m} \frac{1}{4} (\nabla \chi)^2 + \frac{\gamma n^2}{4} \cos \chi + \frac{\hbar^2 n}{8m} \left( \frac{1}{2} (\nabla S_2 - \nabla S_{-2})^2 + \frac{1}{4} (\nabla S_2 + \nabla S_{-2} + 2 \nabla S_0)^2 \right) \right]. \quad (18)$$

One can easily see from Eq. (18) that the minimum energy solution for the kink corresponds to the conditions  $\nabla S_2 - \nabla S_{-2} = 0$  and  $\nabla S_2 + \nabla S_{-2} + 2 \nabla S_0 = 0$ . In this case, the energy (18) can be mapped on the sine-Gordon energy functional. The function  $\chi(x)$  is found from the condition of minimum of  $F_{ph}$ :  $\delta F_{ph} / \delta \chi = 0$  and boundary conditions. Finally, we have

$$\chi(x) = 2\pi - 4 \tan^{-1} \exp(-x / \xi_\chi), \quad (19)$$

$$S_2(x) = S_{-2}(x) = -S_0(x) = -\frac{1}{4} \chi(x), \quad (20)$$

where

$$\xi_\chi = \sqrt{\frac{\hbar^2}{2m} \frac{1}{4\gamma n}} \quad (21)$$

is the healing length for  $\chi$ . It also gives a characteristic length of the kink. The spatial structure of a single kink is presented in Fig. 1, where we have plotted  $x$  dependences of  $\chi$ ,  $S_{\pm 2}$ , and  $S_0$ . A single kink moving in the space is a *soliton*.

As seen from Eq. (18), the energy of the kink  $F_{\text{kink}}$  scales as  $\gamma\xi_\chi \sim \sqrt{\gamma}$ . More accurate calculation based on Eq. (19) yields

$$F_{\text{kink}} = n^{3/2} \sqrt{\frac{\hbar^2 \gamma}{m}}. \quad (22)$$

Note that in real atomic condensates,  $\gamma$  has to be much smaller than  $\alpha$  and, therefore,  $\xi_\chi$  should far exceed the coherence length, which has a meaning of a length scale for the density modulations.

The same structure of the kink can be obtained from the extended Gross-Pitaevskii equations supplemented by the boundary conditions for  $\chi$ . These equations are rather cumbersome and we do not present them here (see, e.g., our previous work [19]). It is easy to see by the direct substitution that Eqs. (19)–(21) with the *constant* populations of each magnetic sublevel yield the *exact* solution to these equations.

#### IV. ROUGHENING

At zero temperature, the sine-Gordon system in any dimension is locked in one of equivalent minima corresponding to  $\chi = 2\pi l$ . At nonzero temperature, the presence of kinks can be favorable due to their entropic contribution to the free energy [18]. Kinks destroy the long-range order and lead to the rough phase, in which the height of the fluctuations diverges while tending the system size to infinity. In a one-dimensional system of length  $L$  ( $L \gg \xi_\chi$ ), the entropy corresponding to the single kink can be estimated as  $\ln L/\xi_\chi$ , where  $L/\xi_\chi$  is just a number of places to put a kink. Therefore, a roughening temperature is

$$T_R^{\text{1D}} \approx \frac{F_{\text{kink}}}{k_B \ln L/\xi_\chi}. \quad (23)$$

In the case of an infinite 1D system ( $L \rightarrow \infty$ ),  $T_R^{\text{1D}} = 0$ , and this result is very different from the result of the small-amplitude oscillations approach, which shows the presence of long-range order for  $\chi$ . Relation (23) is well known for the sine-Gordon model and can be derived using more rigorous analysis, see, e.g., Ref. [20] and references therein. Note that in a 1D finite system,  $T_R^{\text{1D}} \rightarrow 0$ , as  $\gamma \rightarrow 0$ .

In two dimensions, a roughening temperature  $T_R^{\text{2D}}$  can also be estimated in a simple manner. In this case,  $F_{\text{kink}}$  given by Eq. (22) is the energy of a unit length of a kink. The “critical nucleus” for the formation of the rough phase is represented by the step for  $\chi$  of height  $2\pi$  in a functional space with a perimeter  $\approx 2\pi\xi_\chi$ . The total energy of such a nucleus is of the order of  $2\pi F_{\text{kink}}\xi_\chi = \pi\hbar^2 n_s / 2m$ , where we have changed density of atoms  $n$  by superfluid density  $n_s$  just below the roughening transition. It is interesting to note that this energy is independent of  $\gamma$ . A roughening temperature  $T_R^{\text{2D}}$  can be estimated by equating the nucleus energy to the thermal one,  $T_R^{\text{2D}} \approx \pi\hbar^2 n_s / 2k_B m$ . A more rigorous calculation for the 2D sine-Gordon model, based on renormalization-group analysis [18], yields the same result,

$$T_R^{\text{2D}} = \frac{\pi\hbar^2 n_s}{2k_B m}, \quad (24)$$

which is independent of  $\gamma$ . The expression (24) for the roughening temperature coincides with the well-known result for the temperature of the BKT transition, see, e.g., [7]. To describe properly the behavior of the system in this strongly fluctuative region at high temperatures, one needs more careful and detailed analysis. However, from the considerations presented here, we can conclude that a long-range order for  $\chi$  in 2D systems survives up to quite high temperatures (of the order of the BKT critical temperature). Possibly, the roughening transition occurs simultaneously with the BKT transition, and both types of topological defects, namely kinks and vortices, proliferate together.

If the size of a 1D or 2D finite system is less than  $\xi_\chi$  (which can be much larger than the coherence length), the formation of a kink is impossible, and, therefore, we expect that  $\chi$  is nearly constant inside the cloud. At the same time, a value of  $\chi$  can be different from  $2\pi l$  due to thermal fluctuations. We have studied a similar situation before in Ref. [21], where the case of a harmonically trapped quasi-two-dimensional  $F=1$  condensate was treated. An infinite homogeneous  $F=1$  condensate in zero magnetic field can be either in polar or ferromagnetic states, where atoms populate only one or two hyperfine states and the spin-mixing terms in the energy are zero in the equilibrium. However, in the case of a trapped rotated condensate containing vortices, there are some regions on the phase diagram where all three hyperfine states are populated, and the energy depends on the relative phase among them [22–27]. This vortex phase can be both locally and globally stable, and some of them, like the Mermin-Ho vortex, have an axial symmetry, which makes them rather simple. Another method to create such states was recently used experimentally in Ref. [28], where a microwave energy was injected into the system leading to the redistribution of particles from the spin  $-1$  state to spin 0 and 1 states. We see that the cyclic phase in the  $F=2$  condensate is one more example of the systems, where spin mixing is important. It is remarkable that, in this case, spin mixing is happening in the ground state and even in the absence of rotation, applied magnetic fields, and other external perturbations, as in the  $F=1$  case. For the spin-mixing dynamics in the  $F=1$  condensate, see Ref. [29]. Note that recently, a new experimental method for nondestructive study of internal degrees of freedom of spinor condensates was proposed in Ref. [30].

#### V. FERROMAGNETIC STATE IN THE $F=1$ SPINOR CONDENSATE

In this section, we discuss thermal fluctuations in a spinor  $F=1$  condensate. We concentrate on the ferromagnetic state. The energy of the system is independent of the direction of the spin but it depends on the gradients of spin. A similar situation exists for the phase of the order parameter in a scalar condensate (the energy is independent of its value, but depends on the gradient), and therefore we can expect similar behavior, namely, the absence of long-range order



in low dimensions. This result was obtained before for a large class of discrete and continuous spin models [18]. We assume that in the ground state all the atoms occupy only one hyperfine state,  $\Psi_{-1}=\Psi_0=0$ ,  $\Psi_1=\sqrt{n}$ , and study small-amplitude oscillations induced by the temperature. In the ground state, spin is oriented along the  $z$  direction,  $S_z=1$ ,  $S_x=S_y=0$ . Perturbations of  $\Psi_0$  lead to that of  $S_x$  and  $S_y$ ,

$$\delta S_x = \frac{1}{\sqrt{2n}}(\delta\Psi_0 + \delta\Psi_0^*), \quad (25)$$

$$\delta S_y = \frac{i}{\sqrt{2n}}(\delta\Psi_0 - \delta\Psi_0^*). \quad (26)$$

The spectrum for the  $F=1$  condensate was obtained in Refs. [15,16]. The frequency of the mode, responsible for fluctuations of  $\Psi_0$ , is given by

$$\hbar\omega = \frac{\hbar^2 \mathbf{k}^2}{2m}. \quad (27)$$

Using Eqs. (25)–(27), we have calculated mean-square fluctuations for  $S_x$  and  $S_y$  at large distances and found an absence of long-range order in 1D and 2D cases. As an example, we show here the correlator in the 2D situation,

$$\langle [S_x(\mathbf{r}) - S_x(0)]^2 \rangle_T = \frac{T}{T_d} \ln \frac{r}{\lambda_T}, \quad (28)$$

where

$$\lambda_T = \frac{\hbar}{\sqrt{2mk_B T}}. \quad (29)$$

The derivation is similar to the case of the cyclic phase in the  $F=2$  condensate, presented in Sec. IV. The result for  $S_y$  is the same. For the 1D case, the correlators behave as  $\sim r$ . This implies that in low-dimensional systems there is no long-range order in the direction of spin. However, the 2D system can be divided into blocks of a characteristic size  $L$ ,  $\xi_s \ll L \ll \lambda_T \exp(T_d/T)$ , with nearly the same direction of spin in each block ( $\xi_s \sim \sqrt{\frac{\hbar^2}{2m\beta n}}$  is the spin healing length). Different blocks are uncorrelated with each other. At temperatures of the order of  $2T_d/\ln(k_B T_d/\beta n)$ ,  $\xi_s$  becomes comparable to  $\lambda_T \exp(T_d/T)$ . In this case, the size of each block becomes of the order of  $\xi_s$ .

In the quasi-1D and -2D trapped condensates, which have finite sizes exceeding  $\xi_s$ , at low temperatures, one can expect the same orientation of spins throughout the system, and at higher temperatures, the cloud should consist of uncorrelated blocks. This is similar to the problem of a fluctuating phase in quasi-1D and -2D scalar condensates [5–7]. Note that the case of the ferromagnetic  $F=2$  condensate is similar to that of the spin-1 system.

## VI. CONCLUSIONS

We have studied the effect of thermal fluctuations on a homogeneous infinite Bose-Einstein condensate with spin  $F=2$ . We were interested in the cyclic state of this system, in which all the particles occupy three hyperfine states with  $m_F=0, \pm 2$ , and the energy depends on the relative phase  $\chi=2S_0-S_2-S_{-2}$  through the spin-mixing term. Using Bogoliubov–de Gennes equations, we have calculated mean-square fluctuations of  $S_0, S_{\pm 2}$  and found an absence of long-range order in one- and two-dimensional situations, but the presence of this order for  $\chi$ . Then, we went beyond the small-amplitude oscillations approach and mapped our problem on the sine-Gordon model. A structure and energy of a single kink separating two topological sectors with different values of  $\chi$  were found. In 1D and 2D situations, thermal proliferation of kinks can lead to the roughening transition destroying a long-range order for  $\chi$ . The one-dimensional infinite system is always in a rough phase, and the finite 1D system, as well as the infinite 2D system, experience a roughening transition at finite temperatures, whose values we have estimated. At the end, we have also discussed thermal fluctuations in the ferromagnetic  $F=1$  system and found an absence of long-range order in the direction of spin in low-dimensional situations, as can be expected from the Mermin-Wagner-Berezinskii theorem. We defined the typical size of each block, within which still there is an order in spin degrees of freedom. We expect that finite systems should be in the ordered phase at low temperatures.

## ACKNOWLEDGMENTS

We acknowledge very useful discussions with H. Adachi and T. K. Ghosh. W.V.P. is supported by the Japan Society for the Promotion of Science.

- 
- [1] A. Gorlitz, J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle, Phys. Rev. Lett. **87**, 130402 (2001).
  - [2] V. Schweikhard, I. Coddington, P. Engels, V. P. Mogendorff, and E. A. Cornell, Phys. Rev. Lett. **92**, 040404 (2004).
  - [3] D. Rychtarik, B. Engeser, H.-C. Nagerl, and R. Grimm, Phys. Rev. Lett. **92**, 173003 (2004).
  - [4] N. L. Smith, W. H. Heathcote, G. Hechenblaikner, E. Nugent, and C. J. Foot, J. Phys. B **38**, 223 (2005).
  - [5] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000).
  - [6] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **87**, 050404 (2001).
  - [7] D. S. Petrov, D. M. Gangardt, and G. V. Shlyapnikov, J. Phys. IV **116**, 5(2005).
  - [8] H. Schmaljohann, M. Erhard, J. Kronjäger, M. Kottke, S. van Staa, L. Cacciapuoti, J. J. Arlt, K. Bongs, and K. Sengstock, Phys. Rev. Lett. **92**, 040402 (2004).
  - [9] M.-S. Chang, C. D. Hamley, M. D. Barrett, J. A. Sauer, K. M.

- Fortier, W. Zhang, L. You, and M. S. Chapman, Phys. Rev. Lett. **92**, 140403 (2004).
- [10] A. Görlitz, T. L. Gustavson, A. E. Leanhardt, R. Löw, A. P. Chikkatur, S. Gupta, S. Inouye, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **90**, 090401 (2003).
- [11] T. Kuwamoto, K. Araki, T. Eno, and T. Hirano, Phys. Rev. A **69**, 063604 (2004).
- [12] A. Widera, F. Gerbier, S. Fölling, T. Gericke, O. Mandel, and I. Bloch, e-print cond-mat/0604038.
- [13] C. V. Ciobanu, S.-K. Yip, and Tin-Lun Ho, Phys. Rev. A **61**, 033607 (2000).
- [14] M. Koashi and M. Ueda, Phys. Rev. Lett. **84**, 1066 (2000).
- [15] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. **67**, 1822 (1998).
- [16] T.-L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
- [17] J. P. Martikainen and K. A. Suominen, J. Phys. B **34**, 4091 (2001).
- [18] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge Univ. Press, Cambridge, 1995).
- [19] W. V. Pogosov, R. Kawate, T. Mizushima, and K. Machida, Phys. Rev. A **72**, 063605 (2005).
- [20] S. Ares, J. A. Cuesta, A. Sanchez, and R. Toral, Phys. Rev. E **67**, 046108 (2003).
- [21] W. V. Pogosov and K. Machida, e-print cond-mat/0602119.
- [22] T. Mizushima, K. Machida, and T. Kita, Phys. Rev. Lett. **89**, 030401 (2002).
- [23] T. Isoshima and K. Machida, Phys. Rev. A **66**, 023602 (2002).
- [24] T. Mizushima, K. Machida, and T. Kita, Phys. Rev. A **66**, 053610 (2002).
- [25] T. Mizushima, N. Kobayashi, and K. Machida, Phys. Rev. A **70**, 043613 (2004).
- [26] T. Kita, T. Mizushima, and K. Machida, Phys. Rev. A **66**, 061601(R) (2002).
- [27] T. Isoshima, K. Machida, and T. Ohmi, J. Phys. Soc. Jpn. **70**, 1604 (2001).
- [28] M. S. Chang, Q. Qin, W. Zhang, L. You, and M. S. Chapman, Nat. Phys. **1**, 111 (2005).
- [29] H. Pu, C. K. Law, S. Raghavan, J. H. Eberly, and N. P. Bigelow, Phys. Rev. A **60**, 1463 (1999).
- [30] J. M. Higbie, L. E. Sadler, S. Inouye, A. P. Chikkatur, S. R. Leslie, K. L. Moore, V. Savalli, and D. M. Stamper-Kurn, Phys. Rev. Lett. **95**, 050401 (2005).