
OKAYAMA UNIVERSITY

Earth Science Reports

Vol. 19, No. 1, 2012

CONTENTS

—◆—

Holocene environmental and sea-level changes of the Okayama Plain – evidence from peat bed – Shigeyuki SUZUKI	1
Folded structure of the Carboniferous Ko-yama Limestone Group, Akiyoshi Belt, SW JAPAN Noriyuki INADA, Shigeyuki SUZUKI, Keisuke ISHIDA	5
The temperature and pressure conditions of synthetic Sibirskite and Parasibirskite Masahiro KANO, Junji YAMAKAWA	13
Estimation of SPM concentration distribution over the central Seto Inland Sea Takahiro EBI, Hirofumi MATSUMOTO, Junji YAMAKAWA	19
The oldest fossil forest in Japan discovered from the Upper Triassic Nariwa Group, Okayama Prefecture, SW Japan – Implication for reconstruction of depositional environment and paleovegetation – Hirokazu YUKAWA, Kazuo TERADA, Ge SUN, Shigeyuki SUZUKI	25
A case study on persistent appearance of the ensemble of precipitation bands in the warm sector of the Baiu front in the western Japan around 19 June 200 Yasuhiro GODA, Kuranoshin KATO, Osamu TSUKAMOTO	39



Department of Earth Sciences
Faculty of Science, OKAYAMA UNIVERSITY
1-1, Tsushima-Naka 3chome, Kita-ku OKAYAMA 700-8530, JAPAN
