



# Chloroplast Genome Sequences of Seven Strains of the Bloom-Forming Raphidophyte *Heterosigma akashiwo*

Sergio Seoane,<sup>a</sup>  Kiwamu Hyodo,<sup>b</sup>  Shoko Ueki<sup>b</sup>

Department of Plant Biology and Ecology, University of the Basque Country (UPV/EHU), Lejona, Spain<sup>a</sup>;  
Institute of Plant Science and Resources, Okayama University, Okayama, Japan<sup>b</sup>

**ABSTRACT** We report here the complete chloroplast genome sequences of seven strains of the bloom-forming raphidophyte *Heterosigma akashiwo*. These ~160-kb sequences contain 124 protein-, 6 rRNA-, and 34 tRNA-coding sequences. Notable sequence variations were observed among these seven sequenced and two previously characterized strains.

*Heterosigma akashiwo* is a eukaryotic, unicellular, bloom-forming alga that belongs to the family *Raphidophyceae*. It has been identified in the Pacific and Atlantic oceans and in both the northern and southern hemispheres, thus spanning a wide climate range from arctic to tropic (1–12). Because its blooms may exert a negative impact on local ecosystems, its population dynamics in the environment are of great interest and importance. Establishing strain markers will allow the monitoring of geographical and temporal dynamics of the species. We previously sequenced whole-mitochondrial genomes of four strains of this species and identified a hypervariable protein-coding sequence associated with the geographic origins of the strains (13–15).

Here, as a part of our continuous effort to establish strain-specific markers for *H. akashiwo*, we sequenced full-length chloroplast DNA (ptDNA) of seven *H. akashiwo* strains with different geographic origins: CCAP934/8 (Puget Sound, WA, USA), EHUSP1 (Bay of Biscay, Spain), CCAP934/4 (Tampa Bay, FL, USA), CCMP2274 (San Francisco Bay, CA, USA), CCMP3374 (East Greenwich, RI, USA), HaFk01 (Fukuoka, Japan), and CCMP1596 (Narragansett, RI, USA). Total DNA was extracted from *H. akashiwo* cells using the DNeasy blood and tissue kit (Qiagen). The libraries were sequenced with the Illumina MiSeq platform, and the obtained paired-end reads were mapped to the previously published sequences, *H. akashiwo* ptDNA of NIES293 (EU168190) and CCMP452 (EU168191) (16), using Burrows-Wheeler alignment software. The mapped reads were extracted using Samtools software and assembled using the Platanus software (17). The obtained contigs were aligned with the reference sequences, and the corresponding segments in the references were replaced with the contigs. The process was repeated up to three times to obtain the final assemblages, and the accuracy of the assembly was validated by mapping the reads on the obtained sequences (18). Protein-coding sequence (open reading frame) prediction and gene annotation were performed with ORFfinder and database searches with BLASTx; tRNAs were predicted by tRNA-scan server, and rRNA coding sequences were identified by a database search with BLASTn.

The lengths of the ptDNAs of strains CCAP934/8, EHUSP01, CCAP934/4, CCMP2274, CCMP3374, HaFk01, and CCMP1596 were 159,918 bp, 160,150 bp, 160,099 bp, 159,321 bp, 160,152 bp, 159,492 bp, and 159,691 bp, respectively, and G+C contents were 30.5% for all the strains. Twenty-four genes related to photosynthesis (two *cfxQs*, *psaA*, *psaB*, *psaC*, *psaD*, *psaF*, *psaL*, *psb28*, two *psbAs*, *psbB*, two *psbCs*, two *psbDs*, *psbE*, *psbH*, *psbV*, two *rbclS*, *ycf3*, *ycf34*, and *ycf35*) were identified in each strain, as reported

**Received** 19 August 2017 **Accepted** 23 August 2017 **Published** 12 October 2017

**Citation** Seoane S, Hyodo K, Ueki S. 2017. Chloroplast genome sequences of seven strains of the bloom-forming raphidophyte *Heterosigma akashiwo*. *Genome Announc* 5:e01030-17. <https://doi.org/10.1128/genomeA.01030-17>.

**Copyright** © 2017 Seoane et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Shoko Ueki, [shokoueki@okayama-u.ac.jp](mailto:shokoueki@okayama-u.ac.jp).

previously for strains CCMP452 and NIES293. Similarly, 34 previously known tRNA genes, including 2 pseudogenes, were identified in all strains. As previously reported, all strains possessed ~22-kb invert repeats, and the strains CCAP934/8, CCMP2274, and HaFk01 contained an ~8.0-kb inversions, flanked between the *tyrC* and *psb28* genes, compared to those of strains EHUSP01, CCAP934/4, CCMP3374, and CCMP1596. The parts of the *H. akashiwo* ptDNA sequences, between 17.5 and 18 kb and 66 and 66.5 kb, showed notable polymorphisms, which may be utilized as strain-specific markers.

**Accession number(s).** The sequences reported here were deposited in DDBJ/EMBL/GenBank under the accession numbers [LC269918](#) (CCAP934/8), [LC269919](#) (EHUSP01), [LC269920](#) (CCAP934/4), [LC269921](#) (CCMP2274), [LC269922](#) (CCMP3374), [LC269923](#) (HaFk01), and [LC269924](#) (CCMP1596).

## ACKNOWLEDGMENTS

This work was supported by KAKENHI (15H01263 and 16H06449a8).

We thank Tomoyuki Shikata (National Research and Development Agency, Japan Fisheries Research and Education Agency) for providing us with strain HaFk01. Strain CCAP934/8 was provided by the Culture Collection of Algae and Protozoa of the Scottish Marine Institute, and strains CCMP2274, CCMP3374, and CCMP1596 were provided by Bigelow Laboratory for Ocean Sciences.

## REFERENCES

- Honjo T. 1993. Overview on bloom dynamics and physiological ecology of *Heterosigma akashiwo*, p 33–41. In Smayda TJ, Shimizu Y (ed), Toxic phytoplankton blooms in the sea. Elsevier, Amsterdam, Netherlands.
- Chang FH, Anderson C, Boustead NC. 1990. First record of a *Heterosigma* (Raphidophyceae) bloom with associated mortality of cage-reared salmon in Big Glory Bay, New Zealand. N Z J Mar Freshwat Res 24: 461–469. <https://doi.org/10.1080/00288330.1990.9516437>.
- Mackenzie L. 1991. Toxic and noxious phytoplankton in Big Glory Bay, Stewart Island, New Zealand. J Appl Phycol 3:19–34. <https://doi.org/10.1007/BF00003916>.
- Black EA, Whyth JNC, Bagshaw JW, Ginther NG. 1991. The effects of *Heterosigma-akashiwo* on juvenile *Oncorhynchus-tshawytscha* and its implications for fish culture. J Appl Ichthyol 7:168–175. <https://doi.org/10.1111/j.1439-0426.1991.tb00523.x>.
- Rensel J, Homer RA, Postel LE. 1989. Effects of phytoplankton blooms on salmon aquaculture in Puget Sound, Washington: initial research. Northwest Environ J 5:53–59.
- Taylor FJR. 1993. Current problems with harmful phytoplankton blooms in British Columbia waters, p 699–703. In Smayda TJ, Shimizu Y (ed), Toxic phytoplankton blooms in the sea. Elsevier, Amsterdam, Netherlands.
- O'Halloran C, Silver MW, Holman TR, Scholin CA. 2006. *Heterosigma akashiwo* in central California waters. Harmful Algae 5:124–132. <https://doi.org/10.1016/j.hal.2005.06.009>.
- Lackey JB, Lackey EW. 1963. Microscopic algae and protozoa in the waters near Plymouth in August 1962. J Mar Biol Assoc 43:797–805. <https://doi.org/10.1017/S0025315400025698>.
- Thronsen J. 1969. Flagellates of Norwegian coastal waters. Nytt Magazin for Botanikk 16:161–216.
- Rojas de Mendiola B. 1979. Red tide along the Peruvian coast, p 183–190. In Taylor DL, Seliger HH (ed), Toxic dinoflagellate blooms. Elsevier, Amsterdam, Netherlands.
- Tseng CK, Zhou MJ, Zou JZ. 1993. Toxic phytoplankton studies in China, p 347–352. In Smayda TJ, Shimizu Y (ed), Toxic phytoplankton blooms in the sea. Elsevier, Amsterdam, Netherlands.
- Engesmo A, Eikrem W, Seoane S, Smith K, Edvardsen B, Hofgaard A, Tomas CR. 2016. New insights into the morphology and phylogeny of *Heterosigma akashiwo* (Raphidophyceae), with the description of *Heterosigma minor* sp. nov. Phycologia 55:279–294. <https://doi.org/10.2216/15-115.1>.
- Higashi A, Nagai S, Seone S, Ueki S. 2017. A hypervariable mitochondrial protein coding sequence associated with geographical origin in a cosmopolitan bloom-forming alga, *Heterosigma akashiwo*. Biol Lett 13. <https://doi.org/10.1098/rsbl.2016.0976>.
- Ogura Y, Nakayama N, Hayashi T, Ueki S. 2016. Mitochondrial genome sequences of four strains of the bloom-forming raphidophyte *Heterosigma akashiwo*. Genome Announc 4(6):e01288-01216. <https://doi.org/10.1128/genomeA.01288-16>.
- Higashi A, Nagai S, Salomon PS, Ueki S. 2017. A unique, highly variable mitochondrial gene with coding capacity of *Heterosigma akashiwo*, class Raphidophyceae. J Appl Phycol. <https://doi.org/10.1007/s10811-017-1142-2>.
- Cattolico RA, Jacobs MA, Zhou Y, Chang J, Duplessis M, Lybrand T, McKay J, Ong HC, Sims E, Rocap G. 2008. Chloroplast genome sequencing analysis of *Heterosigma akashiwo* CCMP452 (West Atlantic) and NIES293 (West Pacific) strains. BMC Genomics 9:211. <https://doi.org/10.1186/1471-2164-9-211>.
- Kajitani R, Toshimoto K, Noguchi H, Toyoda A, Ogura Y, Okuno M, Yabana M, Harada M, Nagayasu E, Maruyama H, Kohara Y, Fujiyama A, Hayashi T, Itoh T. 2014. Efficient de novo assembly of highly heterozygous genomes from whole-genome shotgun short reads. Genome Res 24:1384–1395. <https://doi.org/10.1101/gr.170720.113>.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.