

Thrombocytopenia in Preterm Infants with Intrauterine Growth Restriction

Hidehiko Maruyama*

Masako Shinozuka[†]

Yo-ichi Kondoh[‡]

Yo-ichiro Akahori**

Miwa Matsuda^{††}

Seiji Inoue^{‡‡}

Yumi Sumida[§]

Tsuneo Morishima[¶]

*Department of Pediatrics, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, hidemaru@md.okayama-u.ac.jp

[†]Department of Pediatrics, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

[‡]Department of Pediatrics, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

**Department of Obstetrics and Gynecology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

^{††}Department of Obstetrics and Gynecology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

^{‡‡}Department of Obstetrics and Gynecology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

[§]Department of Obstetrics and Gynecology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

[¶]Department of Pediatrics, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences,

Thrombocytopenia in Preterm Infants with Intrauterine Growth Restriction*

Hidehiko Maruyama, Masako Shinozuka, Yo-ichi Kondoh, Yo-ichiro Akahori,
Miwa Matsuda, Seiji Inoue, Yumi Sumida, and Tsuneo Morishima

Abstract

Sick preterm infants often have thrombocytopenia at birth, and this is often associated with intrauterine growth restriction (IUGR), or birth weights less than the 10th percentile. The pathogenesis of the thrombocytopenia and its importance in IUGR are still unclear. We studied the characteristics of preterm IUGR infants with thrombocytopenia. Twenty-seven singleton Japanese preterm IUGR infants were born between January 2002 and June 2007 at Okayama University Hospital. Infants with malformation, chromosomal abnormalities, alloimmune thrombocytopenia, sepsis, and maternal aspirin ingestion were excluded. The infants were divided into group A (n=8), which had thrombocytopenia within 72h after birth, and group B (n=19), which did not. There were significant differences in birth weight, head circumference, umbilical artery (UA)-pulsatility index (PI), middle cerebral artery-PI, UA-pH, UA-pO₂, and UA-pCO₂. The infants in group A were smaller, had abnormal blood flow patterns, and were hypoxic at birth. We speculate that the infants with thrombocytopenia were more severely growth-restricted by chronic hypoxia. Thrombocytopenia is an important parameter for chronic hypoxia in the uterine.

KEYWORDS: thrombocytopenia, intrauterine growth restriction, chronic hypoxia

Original Article

Thrombocytopenia in Preterm Infants with Intrauterine Growth Restriction

Hidehiko Maruyama^{a*}, Masako Shinozuka^a, Yo-ichi Kondoh^a,
Yo-ichiro Akahori^b, Miwa Matsuda^b, Seiji Inoue^b,
Yumi Sumida^b, and Tsuneo Morishima^a

*Departments of ^aPediatrics, and ^bObstetrics and Gynecology, Okayama University Graduate School of Medicine,
Dentistry and Pharmaceutical Sciences, Okayama 700-8558 Japan*

Sick preterm infants often have thrombocytopenia at birth, and this is often associated with intrauterine growth restriction (IUGR), or birth weights less than the 10th percentile. The pathogenesis of the thrombocytopenia and its importance in IUGR are still unclear. We studied the characteristics of preterm IUGR infants with thrombocytopenia. Twenty-seven singleton Japanese preterm IUGR infants were born between January 2002 and June 2007 at Okayama University Hospital. Infants with malformation, chromosomal abnormalities, alloimmune thrombocytopenia, sepsis, and maternal aspirin ingestion were excluded. The infants were divided into group A (n=8), which had thrombocytopenia within 72h after birth, and group B (n=19), which did not. There were significant differences in birth weight, head circumference, umbilical artery (UA)-pulsatility index (PI), middle cerebral artery-PI, UA-pH, UA-pO₂, and UA-pCO₂. The infants in group A were smaller, had abnormal blood flow patterns, and were hypoxic at birth. We speculate that the infants with thrombocytopenia were more severely growth-restricted by chronic hypoxia. Thrombocytopenia is an important parameter for chronic hypoxia in the uterine.

Key words: thrombocytopenia, intrauterine growth restriction, chronic hypoxia

Sick preterm infants are often thrombocytopenic, and thrombocytopenia occurs in 18-40% of all preterm infants admitted to neonatal intensive care units [1, 2]. Thrombocytopenia is often associated with intrauterine growth restriction (IUGR) [3, 4]. The thrombocytopenia in IUGR infants usually appears within 72h of birth and resolves within 1 week, and the platelet count rarely falls below $5 \times 10^4/\mu\text{l}$ [5, 6]. In contrast, thrombocytopenia in

IUGR is related to intraventricular hemorrhage, preterm delivery, and so on [7]. In IUGR infants, impaired delivery of oxygen and other essential nutrients is thought to limit organ growth and musculoskeletal maturation. The cause of thrombocytopenia is also thought to be this hypoxia and low level of nutrients, but not all IUGR infants are thrombocytopenic. The pathogenesis of thrombocytopenia and its importance in IUGR are still unclear. Therefore, we studied the characteristics of preterm IUGR infants with thrombocytopenia.

Received April 22, 2008; accepted June 9, 2008.

*Corresponding author. Phone: +81-86-235-7247; Fax: +81-86-221-4745
E-mail: hidemaru@md.okayama-u.ac.jp (H. Maruyama)

Materials and Methods

Between January 2002 and June 2007, 952 infants, including 246 preterm infants, were admitted to the neonatal intensive care unit at Okayama University Hospital. These included 27 singleton preterm infants with IUGR, or birth weights less than the 10th percentile for Japanese, and who had no malformations, chromosomal abnormalities, alloimmune thrombocytopenia, sepsis, or maternal aspirin ingestion within 4 weeks of delivery.

Generally, thrombocytopenia in full term infants is defined as a platelet count of less than $15 \times 10^4/\mu\text{l}$. According to the reports on the fetal platelet count of cord blood, the platelet count reaches $15 \times 10^4/\mu\text{l}$ by the end of first trimester [8] and 17.5×10^4 to $25 \times 10^4/\mu\text{l}$ by the end of the second trimester [9]. Therefore, thrombocytopenia in preterm infants is also defined as a platelet count of less than $15 \times 10^4/\mu\text{l}$ at any gestational age. The platelet count of venous blood from the baby was analyzed at day 0 and 1. If it was below $15 \times 10^4/\mu\text{l}$ or if it seemed to drop below $15 \times 10^4/\mu\text{l}$, we continue to check on it until the platelet count was over $15 \times 10^4/\mu\text{l}$. The infants were divided into 2 groups: group A ($n=8$) had thrombocytopenia within 72h after birth, and the remainder constituted group B ($n=19$).

Gestational age was calculated as the best obstetrical estimate according to the last menstrual period combined with a first-trimester ultrasound. Ultrasound scans were performed using 3.5- or 5-MHz probes (SSD 2200; Aloka, Tokyo, Japan). The data on the umbilical artery (UA)-pulsatility index (PI) and middle cerebral artery (MCA)-PI at the last measurement within 1 week before delivery were used. The definition of pregnancy-induced hypertension published by Japan Society of Obstetrics and Gynecology in 2005 was used. The mode of delivery vaginal delivery or cesarean section (C/S) was decided by monitoring indicative conditions such as abnormalities and fetal growth restriction. An umbilical artery gas analysis was performed immediately after birth. In this study, hypoglycemia was defined as a blood glucose level of less than 40 mg/dl. The classification of intraventricular hemorrhage proposed by Papile *et al.* was used [10].

The Japanese standards for birth weight and head circumference (HC) published in 1998 and those for

UA-PI and MCA-PI on fetal ultrasound published in 2003 were used. The PI was defined as (peak velocity–end-diastolic velocity)/mean velocity. As we could not find a good standard for placental weight for Japanese, we used data from Norway [11]. When analyzing the birth weight data, HC, UA-PI, MCA-PI, and placental weight, we considered the gestational age. We also analyzed the standard deviation (SD), presuming that each set of values was normally distributed.

The results are expressed as the mean $\pm$ SD or mean (range). The Mann-Whitney *U* test and Fisher's exact test were used to analyze the relationship between the 2 groups. Values of $p < 0.05$ were considered statistically significant. The data were processed using StatView statistical software (SAS Institute, Cary, NC, USA). The study was approved by the ethics board at the Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences. All parents provided written informed consent upon entry into this study.

Results

The characteristics of the infants are shown in Table 1. The mean gestational age was 32.6 (28.1–35.7) weeks in group A and 33.9 (28.4–36.6) weeks in group B ($p=0.24$). There were no extremely premature babies who were born before 28 weeks. There was no difference in maternal age, primiparity ratio, gender, incidence of pregnancy-induced hypertension, C/S ratio, or Apgar score between the 2 groups.

The birth weight in group A, 1,167 (749–1,784) g, was significantly lower than that in group B, 1,572 (769–2,248) g ($p=0.020$), and the SD in group A (-2.62 ± 0.80) was lower than that in group B (-1.95 ± 0.67) ($p=0.034$) (Fig. 1). The HC in group A, 27.2 (25.0–30.8) cm, was significantly smaller than that in group B, 29.2 (24.3–32.5) cm ($p=0.011$), as was the SD (-1.14 ± 0.58 vs. 0.40 ± 0.67 , group A vs. group B; $p=0.012$) (Fig. 2). The placental weight in group A, 271 (180–368) g, tended to be lower than that in group B, 359 (180–550) g ($p=0.053$), although the SDs were not significantly different (-1.85 ± 0.52 vs. -1.33 ± 0.80 , group A vs. group B; $p=0.094$).

The UA-PI in group A, 1.555 (0.878–2.121), was not significantly larger than that in group B, 1.132 (0.641–1.696) ($p=0.063$), although the SDs were

Table 1 Characteristics of the study infants

	Group A (n = 8)	Group B (n = 19)	p-value
Gestational age (weeks)	32.6 (28.1–35.7)	33.9 (28.4–36.6)	0.24
Maternal age	32.0 (19–41)	32.6 (23–39)	0.94
Primiparity	3 (37.5%)	10 (52.6%)	0.68*
Male	3 (37.5%)	8 (42.1%)	> 0.99*
Pregnancy-induced hypertension	7 (87.5%)	9 (47.4%)	0.090*
C/S	8 (100%)	17 (89.5%)	> 0.99*
Emergency C/S	4 (50%)	9 (47%)	> 0.99*
Apgar score 1 minute	6.3 ± 1.8	7.5 ± 1.6	0.059
Apgar score 5 minutes	8.5 ± 0.9	8.8 ± 0.8	0.26
BW (g)	1167 (749–1784)	1572 (769–2248)	0.020
BW (SD)	–2.62 ± 0.80	–1.95 ± 0.67	0.034
HC (cm)	27.2 (25.0–30.8)	29.2 (24.3–32.5)	0.011
HC (SD)	–1.14 ± 0.58	0.40 ± 0.67	0.012
PW (g)	271 (180–368)	359 (180–550)	0.053
PW (SD)	–1.85 ± 0.52	–1.33 ± 0.80	0.094
BW/PW	4.35 ± 0.69	4.63 ± 1.52	0.98
UA-PI	1.555 (0.878–2.121)	1.132 (0.641–1.696)	0.063
UA-PI (SD)	4.27 ± 3.28	1.30 ± 1.60	0.028
MCA-PI	1.311 (1.016–1.637)	1.800 (1.229–2.434)	0.0010
MCA-PI (SD)	–1.90 ± 0.92	–0.24 ± 1.03	0.0011
UA-pH	7.246 ± 0.109	7.346 ± 0.061	0.026
UA-pO ₂ (mmHg)	11.7 ± 3.8	16.3 ± 4.7	0.038
UA-pCO ₂ (mmHg)	65.1 ± 16.6	48.3 ± 8.5	0.013
UA-BE (mmol/L)	0.65 ± 6.2	0.62 ± 4.6	0.85
UA-HCO ₃ (mmol/L)	28.2 ± 5.2	26.4 ± 4.4	0.44
UA-lactate (mmol/L)	5.3 ± 2.2	3.6 ± 1.7	0.067
Platelets ($\times 10^4/\mu\text{l}$) (day 0)	14.3 ± 2.8	28.2 ± 7.4	0.0001
WBC ($/\mu\text{l}$) (day 0)	8169 ± 6512	9351 ± 4393	0.31
Hemoglobin (g/dl) (day 0)	17.6 ± 1.9	17.6 ± 2.1	0.54
Hypoglycemia	3 (37.5%)	4 (21%)	0.63*
IVH	0 (0%)	0 (0%)	> 0.99*
NEC	0 (0%)	0 (0%)	> 0.99*

There were significant differences in BW, HC, UA-PI, MCA-PI, UA-pH, UA-pO₂, UA-pCO₂, and platelet count, indicating that the infants in group A were more growth restricted and more hypoxic *in utero*.

*Fisher's exact test; the others, Mann-Whitney *U* test.

C/S, Cesarean section; BW, birth weight; HC, head circumference; PW, placental weight; UA, umbilical artery; PI, pulsatility index; MCA, middle cerebral artery; BE, base excess; WBC, white blood cells; IVH, intraventricular hemorrhage; NEC, necrotizing enterocolitis.

4.27 ± 3.28 and 1.30 ± 1.60, respectively, for the 2 groups ($p=0.028$) (Fig. 3). No infant had absent or reversed end-diastolic (ARED) flow in the umbilical artery. The MCA-PI in group A, 1.311 (1.016–1.637), was significantly smaller than that in group B, 1.800 (1.229–2.434) ($p=0.0010$), as was the SD (–1.90 ± 0.92 vs. –0.24 ± 1.03, group A vs. group B; $p=0.0011$) (Fig. 4).

The respective values of the umbilical artery gas analysis in groups A and B were as follows: UA-pH, 7.246 ± 0.109 vs. 7.346 ± 0.061; UA-pO₂, 11.7 ± 3.8 vs. 16.3 ± 4.7 mmHg; and UA-pCO₂, 65.1 ± 16.6 vs. 48.3 ± 8.5 mmHg ($p=0.026$, 0.038, and 0.013, respectively). UA-pH and UA-pO₂ were lower and

UA-pCO₂ was higher in group A compared with group B.

The platelet count at day 0 was significantly lower in group A than in group B ($14.3 \times 10^4 \pm 2.8 \times 10^4/\mu\text{l}$ vs. $28.2 \times 10^4 \pm 7.4 \times 10^4/\mu\text{l}$; $p=0.00010$). The platelet count in group A never dropped below $10 \times 10^4/\mu\text{l}$ within 72h after birth and had increased to more than $15 \times 10^4/\mu\text{l}$ by day 9. There was no difference between the 2 groups in the white blood cell count or hemoglobin at day 0, and also no intergroup difference in the incidence of hypoglycemia. No infant had intraventricular hemorrhage or necrotizing enterocolitis.




Fig. 1 Plot of individual birth weights versus gestational age. The infants with thrombocytopenia (group A, open circles) were smaller than the infants without thrombocytopenia (group B, filled circles) ($p=0.034$). The standard curves for males and multiparity are shown as references.




Fig. 2 Plot of individual head circumferences versus gestational age. The head circumferences of infants with thrombocytopenia (group A, open circles) were smaller than those of the infants without thrombocytopenia (group B, filled circles) ($p=0.012$). The standard curves for males are shown as references.




Fig. 3 Plot of the individual umbilical artery-pulsatility index values versus gestational age. The UA-PI of the infants with thrombocytopenia (group A, open circles) was higher than that of the infants without thrombocytopenia (group B, filled circles) ($p=0.028$). The standard curves are shown as references.




Fig. 4 Plot of the individual middle cerebral artery-pulsatility index values versus gestational age. The MCA-PI of the infants with thrombocytopenia (group A, open circles) was lower than that of the infants without thrombocytopenia (group B, filled circles) ($p=0.0011$). The standard curves are shown as references.

Discussion

The birth weight and HC of the infants with thrombocytopenia (group A) were significantly smaller than those of infants without thrombocytopenia (group B). In this study, all of the infants had IUGR, or weights less than the 10th percentile at birth. Therefore, this result indicated a strong relationship between thrombocytopenia and birth weight.

There are 2 types of IUGR: symmetrical and asymmetrical. Generally, symmetrical IUGR infants have a fetal factor such as a chromosomal anomaly, and birth weight and HC are reduced proportionately for the gestational age. By contrast, asymmetrical

IUGR infants have maternal or placental factors such as pregnancy-induced hypertension or placental insufficiency. Fetal weight is reduced and brain growth is usually spared. The infants in the present study were considered to be asymmetrical, because we excluded cases with malformations and chromosomal abnormalities. Nevertheless, the smaller HC in group A indicates that the infants in group A were severely growth-restricted.

Blood flow analysis of UA and MCA is performed routinely and is used to evaluate fetal well-being [12]. Worsening of these indicators means circulatory failure, *i.e.*, oxygen supply breakdown [13–15]. Placental insufficiency causes inadequate gas exchange

[16]. With circulatory failure, UA-PI generally increases owing to the increased impedance of the placenta, while MCA-PI decreases due to the increased cerebral blood flow. In the present study, UA-PI was significantly higher and MCA-PI was lower in group A compared with group B, indicating that the infants in group A were chronically hypoxic. When ARED flow is recognized in the umbilical artery, the fetal cardiac output is severely redistributed secondary to hypoxia. Fetuses with ARED flow in the UA have a poor prognosis [17, 18]; however, there was no case with ARED in this study.

Platelets are initially produced mainly in the fetal liver, and the production site is transferred to the bone marrow in the third trimester [5]. The liver is the first organ to suffer the effects of growth restriction, because in hypoxia there is a decrease in blood flow to the liver, rather than an increase in blood flow to the ductus venosus [19]. It is not clear how chronic hypoxia affects progenitor cells and megakaryocytes in platelet production.

Umbilical artery gas analysis showed that the UA-pH and UA-pO₂ were lower and the UA-pCO₂ was higher in group A compared with group B. The UA gas analysis is influenced by the delivery method, but there were no differences in C/S, the emergency C/S ratio, or Apgar scores between the 2 groups. Nevertheless, the differences in the UA gas analysis showed that the infants in group A were hypoxic during delivery.

In summary, in the present study there were significant differences in birth weight, HC, UA-PI, MCA-PI, UA-pH, UA-pCO₂, and UA-pO₂ between the groups of infants with and without thrombocytopenia. Infants with thrombocytopenia were more severely growth-restricted by chronic hypoxia. Thrombocytopenia is an important parameter of chronic hypoxia in the uterine. Although there were no differences in short-term complications, the long-term complications in preterm IUGR infants with thrombocytopenia need to be studied.

References

- Mehta P, Vasa R, Neumann L and Karparkin M: Thrombocytopenia in the high-risk infant. *J Pediatr* (1980) 97: 791–794.
- Oren H, Irken G, Oren B, Olgun N and Ozkan H: Assessment of clinical impact and predisposing factors for neonatal thrombocytopenia. *Indian J Pediatr* (1994) 61: 551–558.
- Chakravorty S, Murray N and Roberts I: Neonatal thrombocytopenia. *Early Hum Dev* (2005) 81: 35–41.
- Roberts IA and Murray NA: Thrombocytopenia in the newborn. *Curr Opin Pediatr* (2003) 15: 17–23.
- Murray NA and Roberts IA: Circulating megakaryocytes and their progenitors in early thrombocytopenia in preterm neonates. *Pediatr Res* (1996) 40: 112–119.
- Murray NA, Watts TL and Roberts IA: Endogenous thrombopoietin levels and effect of recombinant human thrombopoietin on megakaryocyte precursors in term and preterm babies. *Pediatr Res* (1998) 43: 148–151.
- Beiner ME, Simchen MJ, Sivan E, Chetrit A, Kuint J and Schiff E: Risk factors for neonatal thrombocytopenia in preterm infants. *Am J Perinatol* (2003) 20: 49–54.
- Pahal GS, Jauniaux E, Kinnon C, Thrasher AJ and Rodeck CH: Normal development of human fetal hematopoiesis between eight and seventeen weeks' gestation. *Am J Obstet Gynecol* (2000) 183: 1029–1034.
- Forestier F, Tissot JD, Vial Y, Daffos F and Hohlfield P: Haematological parameters of parvovirus B19 infection in 13 fetuses with hydrops foetalis. *Br J Haematol* (1999) 104: 925–927.
- Papile LA, Burstein J, Burstein R and Koffler H: Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr* (1978) 92: 529–534.
- Thompson JM, Irgens LM, Skjaerven R and Rasmussen S: Placenta weight percentile curves for singleton deliveries. *Bjog* (2007) 114: 715–720.
- Toal M, Chan C, Fallah S, Alkazaleh F, Chaddha V, Windrim RC and Kingdom JC: Usefulness of a placental profile in high-risk pregnancies. *Am J Obstet Gynecol* (2007) 196: 363 e1–7.
- Figuerola-Diesel H, Hernandez-Andrade E, Acosta-Rojas R, Cabero L and Gratacos E: Doppler changes in the main fetal brain arteries at different stages of hemodynamic adaptation in severe intrauterine growth restriction. *Ultrasound Obstet Gynecol* (2007) 30: 297–302.
- Jackson MR, Walsh AJ, Morrow RJ, Mullen JB, Lye SJ and Ritchie JW: Reduced placental villous tree elaboration in small-for-gestational-age pregnancies: relationship with umbilical artery Doppler waveforms. *Am J Obstet Gynecol* (1995) 172: 518–525.
- Mari G, Hanif F, Kruger M, Cosmi E, Santolaya-Forgas J and Treadwell MC: Middle cerebral artery peak systolic velocity: a new Doppler parameter in the assessment of growth-restricted fetuses. *Ultrasound Obstet Gynecol* (2007) 29: 310–316.
- Lackman F, Capewell V, Gagnon R and Richardson B: Fetal umbilical cord oxygen values and birth to placental weight ratio in relation to size at birth. *Am J Obstet Gynecol* (2001) 185: 674–682.
- Hartung J, Kalache KD, Heyna C, Heling KS, Kuhlrig M, Wauer R, Bollmann R and Chaoui R: Outcome of 60 neonates who had ARED flow prenatally compared with a matched control group of appropriate-for-gestational age preterm neonates. *Ultrasound Obstet Gynecol* (2005) 25: 566–572.
- Kush ML, Gortner L, Harman CR and Baschat AA: Sustained hematological consequences in the first week of neonatal life secondary to placental dysfunction. *Early Hum Dev* (2006) 82: 67–72.
- Pennati G, Corno C, Costantino ML and Bellotti M: Umbilical flow distribution to the liver and the ductus venosus in human fetuses during gestation: an anatomy-based mathematical modeling. *Med Eng Phys* (2003) 25: 229–238.

