

Acta Medica Okayama

Volume 41, Issue 1

1987

Article 7

FEBRUARY 1987

Metaplastic bone formation in a hyperplastic polyp of the stomach: a case report.

Yuji Ohtsuki*

Yoshifumi Danbara[†]

Isao Takeda[‡]

Kiyoshi Takahashi**

Kazuhiko Hayashi^{††}

Hiroshi Sonobe^{‡‡}

Tadashi Yoshino[§]

Tadaatsu Akagi[¶]

*Kochi Medical School,

[†]Tonan Hospital,

[‡]Okayama University,

**Kochi Medical School,

^{††}Okayama University,

^{‡‡}Okayama University,

[§]Okayama University,

[¶]Okayama University,

Metaplastic bone formation in a hyperplastic polyp of the stomach: a case report.*

Yuji Ohtsuki, Yoshifumi Danbara, Isao Takeda, Kiyoshi Takahashi, Kazuhiko Hayashi, Hiroshi Sonobe, Tadashi Yoshino, and Tadaatsu Akagi

Abstract

Metaplastic bony tissue along with hyperplastic mucosal epithelium showing no atypism was detected in biopsy materials from a Yamada type I gastric polyp. The tissue was metaplastic woven bone associated with calcification. Histogenesis of the bone formation is as yet unknown. This is the first reported case of the presence of metaplastic bone accompanied by hyperplastic gastric mucosa so far.

KEYWORDS: stomach, hyperplastic polyp, metaplastic bone, histopathology

*PMID: 3105253 [PubMed - indexed for MEDLINE]

Metaplastic Bone Formation in a Hyperplastic Polyp of the Stomach : A Case Report

Yuji Ohtsuki, Yoshifumi Danbara*, Isao Takeda**, Kiyoshi Takahashi, Kazuhiko Hayashi, Hiroshi Sonobe, Tadashi Yoshino*** and Tadaatsu Akagi***

*Department of Pathology, Kochi Medical School, Kochi 781-51, Japan, *Division of Internal Medicine, Tonan Hospital, Kochi 780, Japan and **First Department of Surgery and ***Second Department of Pathology, Okayama University Medical School, Okayama 700, Japan*

Metaplastic bony tissue along with hyperplastic mucosal epithelium showing no atypism was detected in biopsy materials from a Yamada type I gastric polyp. The tissue was metaplastic woven bone associated with calcification. Histogenesis of the bone formation is as yet unknown. This is the first reported case of the presence of metaplastic bone accompanied by hyperplastic gastric mucosa so far.

Key words : stomach, hyperplastic polyp, metaplastic bone, histopathology

Bone formation in the human stomach has been found, though rarely, among cases of cancer (1-4). However, there has been no report of the presence of bony tissue associated with gastric epithelium showing no atypism so far. We report a case in which bone formation with calcification occurred along with mucosal epithelium in a hyperplastic polyp of the stomach.

Case report

The patient was a 71 year-old male, who complained of poor appetite and fatigue for about one month. Gastrofiberscopy revealed Yamada type I and type IV polyps at the cardiac region of the stomach (Fig. 1). Biopsy materials from the Yamada type IV polyp were necrotic granulation tissue, and mucosal tissue from the Yamada type I polyp showed metaplastic bone formation with calcification along with hyperplastic glandular

tissue revealing no atypism (Fig. 2). This bony tissue was metaplastic woven bone, and




Fig. 1 Endoscopic view of the stomach revealing polyps of Yamada type IV (arrowhead) and type I (double arrowheads).




Fig. 2 Metaplastic bony tissue (arrowheads) associated with calcification along with hyperplastic epithelium obtained from the Yamada type I polyp in Fig. 1. H.E. stain, $\times 60$.

Fig. 3 Higher magnification of metaplastic woven bone. H.E. stain, $\times 350$.

Fig. 4 An osteocyte-like cell (arrowhead) embedded in the osteoid tissue. H.E. stain, $\times 140$.

Fig. 5 Fragmented short collagen bundles in the metaplastic bone revealed by polarized microscopy. H.E. stain, polarized, $\times 140$.

Fig. 6 Strongly positive reaction for calcium deposition in the metaplastic bone. von Kossa's reaction, $\times 140$.

osteoblastic cells were not detected around the bony tissue (Fig. 3). Osteocyte-like flat cells embedded in the osteoid tissue were detected (Fig. 4). The bony tissue was composed of fragmented bundles of collagen fibers, as revealed by polarized microscopy (Fig. 5), and calcium deposition in the central area was demonstrated by von Kossa's reaction (Fig. 6).

The main laboratory data are as follows; *Serum*: total protein 7.2 g/dl (albumin 55.2 %, α_1 -globulin 3.1 %, α_2 -globulin 9.7 %, β -globulin 7.9 %, γ -globulin 24.1 %), total cholesterol 160 mg/dl, gamma-GTP 146 mU, LAP 204 mU, total bilirubin 1.3 mg/dl, (direct 0.6 mg/dl), ALP 239 IU/l, GOT 59 IU/l, GPT 27 IU/l, LDH 59 IU/l, ZTT 10.5, BUN 10.1 mg/dl, uric acid 3.9 mg/dl, creatinine 0.7 mg/dl, Na 140 mEq/l, K 3.9 mEq/l, Cl 100 mEq/l, Ca 8.2 mg/dl (normal range, 8.2-10.8 mg/dl), P 3.1 mg/dl; *Blood cells*: RBC $438 \times 10^4/\text{mm}^3$, WBC $8000/\text{mm}^3$ (St. 5 %, Seg. 32 %, Eo. 2 %, Ba 0 %, Mo. 4 %, Lym 57 %), Hb 16.8 g/dl, Ht 43 %, platelets $22.1 \times 10^4/\text{mm}^3$; *Urine*: n.p. Laboratory data disclosed only a mild degree of liver dysfunction. Serum electrolytes (Na, K, Cl, Ca and P) were within the normal range. The clinical course of the patient has been favorable.

Discussion

Four cases of ossification in stomach can-

cer, including both primary and metastatic foci, have been reported in the literature (1-4). Papillary and tubular adenocarcinomas were the predominant histological types (1-4). On the other hand, calcification alone has frequently been observed in mucinous type carcinoma (4). Although ossification was experimentally observed in precancerous lesions induced by chemical carcinogens in the remaining stomach following subtotal gastrectomy in rats (5,6), or in the transplanted gastric mucosa of rats (7), no report has been observed on the presence of bony tissue associated with atypical or hyperplastic epithelium in the human stomach so far in the literature. In the present case, bony tissue with calcification was observed along with non-malignant, but hyperplastic epithelium with calcification.

As to the histogenesis of ossifications, the tissue induction theory, in which fibroblastic cells are purported to be activated and differentiate to form metaplastic bone in the stroma under the influence of the cancerous epithelium, is widely accepted (4,6,7). As for the bone formation beneath mucosal epithelium with no atypism, the same woven bone as in the present case was reported in a case of juvenile polyp in the rectum (8). Considering the hamartomatous origin of this kind of polyp, the primitive mesenchyme may play an important role in stromal osteogenesis (8).

It is likely that the resection of the fundic glands, resulting in increased alkalinity, could induce bony tissue as reported previously (5,9), but such an acid deficiency was not likely in the present case. Calcification preceding ossification as seen in the present case has been emphasized in cases of colon cancer (10). In addition, it is of interest that bone formation was exclusively observed just beneath the regenerating epithelium, but not in the originally transplanted gastric mucosa in the abdominal wall in rats (7).

Therefore, not only malignant epithelium, but also regenerating or even hyperplastic epithelium as in the present case could induce stromal osteogenesis under certain special conditions as yet undetermined.

However, as reported previously (3), the total number of stomach cancers associated with calcification is very small, when compared with the frequent occurrence of the disease. This fact is not fully explained yet.

Acknowledgement. The authors are grateful to Mr. T. Yamaguchi and Miss M. Ohkuchi for their technical assistance.

References

1. Gruber GB: Knochenbildung in einem Magenkarzinom. Zieglers Beitr Pathol Anat (1913) **55**, 368-370.
2. Hasegawa T: Zur Kenntnis der Stromaverknöcherung in Karzinomen des Digestionstraktes. Wien Klin Wschr (1923) **36**, 653-656.
3. Yasuma T, Hashimoto K, Miyazawa R and Hiyama Y.: Bone formation and calcification in gastric cancer — Case report and review of literature —. Acta Pathol Jpn (1973) **23**, 155-172.
4. Rhone DP and Horowitz RN: Heterotopic ossification in the pulmonary metastases of gastric adenocarcinoma — Report of a case and review of the literature —. Cancer (1976) **38**, 1773-1780.
5. Ackerman NB: Bone formation in the gastric mucosa following subtotal gastrectomy in rats. Exp Mol Pathol (1968) **9**, 125-130.
6. Werner B, Dahm K and Breucker H: Ossification in cancer of the stomach. An experimental study. Z Krebsforsch (1976) **86**, 147-154.
7. Miyairi B: Experimental studies on bone formation induced by the epithelium of gastric mucosa. Tr Soc Pathol Jpn (1956) **45**, 645-654. (in Japanese).
8. Ohtsuki Y, Takeda I, Takahashi K, Akagi T, Ohmori M, Watanabe R and Sasaki S: A case of juvenile polyp with stromal ossification in the rectum. Geka (1984) **46**, 540-542. (in Japanese).
9. Gemell NI: Calcification within a gastric carcinoma. A case report. Am J Roentgenol (1964) **91**, 779-783.
10. van Patter HT and Whittick JW: Heterotopic ossification in intestinal neoplasms. Am J Clin Pathol (1955) **31**, 73-84.

Received: August 27, 1986

Accepted: November 13, 1986

Correspondence to:

Yuji Ohtsuki
Department of Pathology
Kochi Medical School
Kochi 780, Japan