

Acta Medica Okayama

Volume 37, Issue 6

1983

Article 11

DECEMBER 1983

Type C virus particles produced in human T-cell lines derived from acute lymphoblastic leukemia and a leukemic T-lymphoid malignancy.

Takuzo Oda*

Sekiko Watanabe[†]

Takashi Nakamura[‡]

*Okayama University,

[†]Okayama University,

[‡]Okayama University,

Type C virus particles produced in human T-cell lines derived from acute lymphoblastic leukemia and a leukemic T-lymphoid malignancy.*

Takuzo Oda, Sekiko Watanabe, and Takashi Nakamura

Abstract

Electron microscopy of four human T-cell lines revealed the production of type C virus particles in two T-cell lines: one derived from acute lymphoblastic leukemia and the other from a leukemic T-lymphoid malignancy. Virus particles isolated from these cells had reverse transcriptase activity and the major internal structural protein of 30,000 daltons (p30). The indirect immunofluorescence test of these virus-producing cells with sera of patients with adult T-cell leukemia (ATL) was negative. The data indicate that these retroviruses are different from adult T-cell leukemia virus (ATLV).

KEYWORDS: type C virus particles, human T-cell lines, electron microscopy, virion proteins, immunofluorescence test

*PMID: 6198871 [PubMed - indexed for MEDLINE]

Copyright (C) OKAYAMA UNIVERSITY MEDICAL SCHOOL

Acta Med. Okayama 37, (6), 529-533 (1983)

— BRIEF NOTE —

**TYPE C VIRUS PARTICLES PRODUCED IN HUMAN T-CELL
LINES DERIVED FROM ACUTE LYMPHOBLASTIC
LEUKEMIA AND A LEUKEMIC T-
LYMPHOID MALIGNANCY**

Takuzo ODA, Sekiko WATANABE and Takashi NAKAMURA

Department of Biochemistry, Cancer Institute, Okayama University Medical School, Okayama 700, Japan

Received October 19, 1983

Abstract. Electron microscopy of four human T-cell lines revealed the production of type C virus particles in two T-cell lines : one derived from acute lymphoblastic leukemia and the other from a leukemic T-lymphoid malignancy. Virus particles isolated from these cells had reverse transcriptase activity and the major internal structural protein of 30,000 daltons (p30). The indirect immunofluorescence test of these virus-producing cells with sera of patients with adult T-cell leukemia (ATL) was negative. The data indicate that these retroviruses are different from adult T-cell leukemia virus (ATLV).

Key words : type C virus particles, human T-cell lines, electron microscopy, virion proteins, immunofluorescence test.

Type C retroviruses (HTLV-I, II and ATLTV) were isolated from T-cells and T-cell lines derived from patients with cutaneous T-cell lymphoma and leukemia (1-2), adult T-cell leukemia (3-5), and hairy cell leukemia (6-7). HTLV-I and ATLTV have been shown to be identical or closely related (8). We recently isolated a type C retrovirus (TALV) from a human T-cell line (TALL-1) derived from acute lymphoblastic leukemia (9). TALV has the major internal structural protein of 34,000 daltons (p34) and is different from HTLV-I, II and ATLTV (9-12). In the present paper, we report the production of type C retroviruses from two other human T-cell lines ; Molt-4 derived from acute lymphoblastic leukemia and HPB-MLT from a leukemic T-lymphoid malignancy. These viruses were found to be different from ATLTV and TALV.

Materials and Methods. The human T-cell lines examined for the presence of virus particles were Molt-4 (13) and CCRF-CEM (14) derived from acute lymphoblastic leukemia, Jurkat from T-cell lymphoma, and HPB-MLT from a leukemic T-lymphoid malignancy (15). Molt-4 was provided by Dr. H. Hidaka of the Hayashibara Research Institute for Biological Chemistry, and other cell lines were provided by Dr. T. Tanaka of Kagawa Medical College. MT-2, an ATLTV-producing T-cell line, was a generous gift of Dr. I. Miyoshi of Kochi


Medical College. These cell lines were cultured in RPMI-1640 medium supplemented with heat-inactivated 20 % fetal bovine serum at 37 °C in an atmosphere of 5 % CO₂ in air under conditions precluding laboratory contamination. The fetal bovine serum was free of viral particles. Cell suspensions were centrifuged at $1,000 \times g$ for 10 min, the supernatant was removed, and the pellet was fixed with 2.5 % glutaraldehyde and 1 % osmium tetroxide, dehydrated in an ethanol series, and embedded in Epon 812. Thin-sectioned specimens were stained with 2 % uranyl acetate and lead citrate, and examined under an electron microscope. Viral particles were isolated as described (10-12). Virion proteins were analyzed by 13 % polyacrylamide-sodium dodecyl sulfate (SDS) slab gel electrophoresis according to the method of Laemmli (16). The indirect immunofluorescence test of virus-producing cells was performed as described elsewhere (3).

Results. Electron microscopy of thin sections of these human T-cells revealed the production of type C viruses in two cell lines, Molt-4 and HPB-MLT (Figs. 1 and 2). To distinguish these viruses from HTLV/ATLV and TALV, the type C virus detected in Molt-4 was tentatively designated as MOLV and that in HPB-MLT as MLTV. Many virus particles were found extracellularly, and some in the cisternal spaces. Typical type C virus particles budding from cell membranes were also observed (Figs. 1 and 2 b-e). Most of the extracellular MOLV were typical mature type C particles measuring $118.4 \text{ nm} \pm 12.9 \text{ nm}$ in diameter with a centrally located and homogeneously condensed nucleoid ($78.6 \text{ nm} \pm 11.1 \text{ nm}$). Intracisternal MOLV particles having budded or budding from the endoplasmic reticulum membranes were immature type A particles with a ring- or arc-form nucleoid (Fig. 1f). Most of the extracellular MLTV were mature type C particles measuring $119.4 \text{ nm} \pm 18.3 \text{ nm}$ in diameter with a centrally located and unhomogeneously condensed nucleoid ($79.3 \text{ nm} \pm 12.0 \text{ nm}$). Intracisternal immature type A particles of MLTV having budded or budding from the endoplasmic reticulum membranes were also observed (Fig. 2f).

The indirect immunofluorescence test of MOLV-producing Molt-4 cells and MLTV-producing HPB-MLT cells using sera from patients with adult T-cell leukemia was negative, while that of ATLAV-producing MT-2 cells using the same sera was strongly positive. Molt-4 cells and HPB-MLT cells were not stained with antisera to TALV, bovine leukemia virus (BLV), feline leukemia/sarcoma

Fig. 1. Electron micrographs of MOLV-producing Molt-4 cells (a), type C virus particles budding from cell membranes (b, c, d and e), intracisternal budding and immature particles (f), and protein bands of partially purified MOLV electrophoresed in the SDS-polyacrylamide gel and stained with Coomassie blue (g). Mr, molecular markers in kilodaltons (KD); AT, ATLAV proteins; MO, MOLV proteins. a-f, $\times 46,600$.

Fig. 2. Electron micrographs of MLTV-producing HPB-MLT cells (a), budding viruses in various stages (b, c, d and e), intracisternal budding and immature particles (f), and protein bands of partially purified MLTV electrophoresed in the SDS-polyacrylamide gel and stained with Coomassie blue (g) ML, MLTV proteins. a-f, $\times 46,600$.



viruses (FeLV/FeSV), simian sarcoma virus (SSV-1/SSAV-1), avian leukemia virus (ALV), avian myeloblastosis virus (AMV), and vesicular stomatitis virus (VSV), but were faintly stained with antisera to murine sarcoma/leukemia viruses (MSV/MLV).

Both MOLV and MLTV had the major internal structural protein of 30,000 daltons (p30) (Figs. 1 g and 2 g) and showed reverse transcriptase activity preferring Mn^{2+} in a reaction with polyriboadenylate·oligodeoxyribothymidylate [poly (A)·oligo (dT)₁₂₋₁₈] as a synthetic template-primer.

Discussion. The data presented in this report strongly suggest that MOLV and MLTV are different from ATL (HTLV) and TALV. The major internal structural protein of MOLV and MLTV is p30, while that of ATL (HTLV) and TALV is p24 and p34, respectively. Although the ultrastructures of MOLV and MLTV appear to be slightly different, electrophoretic mobilities of their virion proteins were similar. Further characterization of proteins, RNA, and immunological properties of purified particles of these retroviruses is now under way to determine the origin of these viruses. Comparative studies of the mode of integration and expression of ATL (HTLV), TALV, MOLV and MLTV genomes in human T-cells and T-cell lines should be quite significant in obtaining further information on the viral origin of human T-cell lymphoma and leukemia.

Acknowledgments. We thank Dr. T. Tanaka for generously providing CCRF-CEM, Jurkat, and HPB-MLT cell lines, Dr. H. Hidaka and Dr. T. Kunitomi for Molt-4, Dr. I. Miyoshi and Dr. S. Hiraki for MT-2, Dr. K. Kitajima and Dr. K. Takahashi for antisera from patients with ATL, Dr. N. Ida for antisera to MSV/MLV, Dr. H. Nakashima for antisera to BLV/BSV, and Dr. I. Tomoda and Dr. A. Hasegawa for antisera to FeLV/FeSV, and Ortho Diagnostic Systems Inc. for OK series of monoclonal antibodies.

REFERENCES

1. Poiesz, B.J., Ruscetti, F.W., Gazdar, A.F., Bunn, P.A., Minna, J.D. and Gallo, R.C.: Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proc. Natl. Acad. Sci. U.S.A.* **77**, 7415-7419, 1980.
2. Poiesz, B.J., Ruscetti, F.W., Reitz, M.S., Kalyanaraman, V.S. and Gallo, R.C.: Isolation of a new type C retrovirus (HTLV) in primary uncultured cells of a patient with Sézary T-cell leukemia. *Nature* **294**, 268-271, 1981.
3. Hinuma, Y., Nagata, K., Hanaoka, M., Nakai, M., Matsumoto, T., Kinoshita, K.I., Shirakawa, S. and Miyoshi, I.: Adult T-cell leukemia: Antigen in an ATL cell line and detection of antibodies to the antigen in human sera. *Proc. Natl. Acad. Sci. U.S.A.* **78**, 6476-6480, 1981.
4. Miyoshi, I., Kubonishi, I., Yoshimoto, S., Akagi, T., Ohtsuki, Y., Shiraishi, Y., Nagata, K. and Himura, Y.: Type C virus particles in a cord T-cell line derived by co-cultivating normal human cord leukocytes and human leukemic T-cells. *Nature* **294**, 770-771, 1981.
5. Yoshida, M., Miyoshi, I. and Hinuma, Y.: Isolation and characterization of retrovirus from cell lines of human adult T-cell leukemia and its implication in the disease. *Proc. Natl. Acad. Sci. U.S.A.* **79**, 2031-2035, 1982.
6. Kalyanaraman, V.S., Sarngadharan, M.G., Robert-Guroff, M., Miyoshi, I., Blayney, D., Golde,

- D. and Gallo, R.C.: A new subtype of human T-cell leukemia virus (HTLV-II) associated with a T-cell variant of hairy cell leukemia. *Science* **218**, 571-573, 1982.
7. Gallo, R.C. and Reitz, Jr., M.S.: Human retroviruses and adult T-cell leukemia-lymphoma. *J. Natl. Cancer Inst.* **69**, 1209-1214, 1982.
 8. Popovic, M., Reitz, Jr., M.S., Sarngadharan, M.G., Robert-Guroff, M., Kalyanaraman, V.S., Nakao, Y., Miyoshi, I., Minowada, J., Yoshida, M., Ito, Y. and Gallo, R.C.: The virus of Japanese adult T-cell leukemia is a member of the human T-cell leukemia virus group. *Nature* **300**, 63-66, 1982.
 9. Oda, T., Watanabe, S., Shigehara, T. and Nakamura, T.: Budding type C virus particles in a human T-cell line derived from acute lymphoblastic leukemia. *J. Electron Microsc.* **32**, 172-175, 1983.
 10. Oda, T., Ikeda, S., Tsutsui, T. and Watanabe, S.: Isolation and properties of a retrovirus from a T-cell line derived from acute lymphoblastic leukemia. *Seikagaku* **55**, 769, 1983.
 11. Oda, T., Watanabe, S., Ikeda, S., Tsutsui, K. and Maki, Y.: Comparative studies of a type C retrovirus in a human T-cell line from acute lymphoblastic leukemia and adult T-cell leukemia virus. In *Proc. Jpn. Cancer Assoc.*, 42nd Ann. Meet., Nagoya, p. 371, 1983.
 12. Ikeda, S., Tsutsui, K., Watanabe, S. and Oda, T.: Isolation of a retrovirus from a T-cell line derived from human acute lymphoblastic leukemia and analysis of its constitutional proteins. In *Proc. Jpn. Cancer Assoc.*, 42nd Ann. Meet., Nagoya, P.374, 1983.
 13. Minowada, J., Ohnuma, T. and Moore, G.E.: Rosette-forming human lymphoid cell lines. I. Establishment and evidence for origin of thymus-derived lymphocytes. *J. Natl. Cancer Inst.* **49**, 891-894, 1972.
 14. Foley, G.E., Lazarus, H., Farber, S., Uzman, B.G., Boon, B.A. and McCarthy, R.E.: Continuous culture of human lymphoblasts from peripheral blood of a child with acute leukemia. *Cancer* **18**, 522-529, 1965.
 15. Morikawa, S., Tatsumi, E., Baba, M., Harada, T. and Yasuhira, K.: Two E-rosette-forming lymphoid cell lines. *Int. J. Cancer* **21**, 166-170, 1978.
 16. Laemmli, U.K.: Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685, 1970.