

# *Mathematical Journal of Okayama University*

---

*Volume 32, Issue 1*

1990

*Article 13*

JANUARY 1990

---

## Some Characterizations of $\pi$ -Regular Rings of Bounded Index

Yasuyuki Hirano\*

\*Okayama University

Copyright ©1990 by the authors. *Mathematical Journal of Okayama University* is produced by  
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

## SOME CHARACTERIZATIONS OF $\pi$ -REGULAR RINGS OF BOUNDED INDEX

Dedicated to Professor Manabu Harada on his 60th birthday

YASUYUKI HIRANO

In this paper, all rings contain an identity element. A ring  $R$  is said to be of *bounded index (of nilpotence)* if there is a positive integer  $n$  such that  $a^n = 0$  for all nilpotent elements  $a$  of  $R$ . The least such integer is called the *index* of  $R$ , and we denote it by  $i(R)$ . Recall that  $R$  is said to be  $\pi$ -*regular* if for each element  $a$  of  $R$ , there exists a positive integer  $n$  and an element  $x$  of  $R$  such that  $a^n = a^n x a^n$ . The purpose of this paper is to give some characterizations of  $\pi$ -regular rings of bounded index. We show that a ring  $R$  is a  $\pi$ -regular ring of index at most  $n$  if and only if the endomorphism ring  $\text{End}_R(M)$  of any cyclic module  $M$  is of index at most  $n$ . From this result, we obtain that every finite extension of a  $\pi$ -regular ring of bounded index is also a  $\pi$ -regular ring of bounded index. We also show that a ring  $R$  of bounded index is  $\pi$ -regular if and only if every prime factor ring of  $R$  is Artinian. Using this result, we prove that if  $S$  is a  $\pi$ -regular ring of bounded index and if  $S$  is a finite normalizing extension of a ring  $R$ , then  $R$  is also a  $\pi$ -regular ring of bounded index.

Let  $R$  be a ring of index  $n$ . Then it is proved in [2] that the following statements are equivalent :

- 1)  $R$  is a  $\pi$ -regular ring.
- 2) It holds that  $a^n R = a^{n+1} R$  for all  $a \in R$ .
- 3) It holds that  $R a^n = R a^{n+1}$  for all  $a \in R$ .

Thus, if  $R$  is a  $\pi$ -regular ring of index  $n$ , then every factor ring of  $R$  is of index at most  $n$ . However the converse is not true. For example, for any positive integer  $k$ , the  $k$ -th Weyl algebra  $A_k(Q)$  over the field  $Q$  of rational numbers is a simple domain, but it is not  $\pi$ -regular. Nevertheless, we can show that the center of such a ring is  $\pi$ -regular.

**Proposition 1.** *If every factor ring of a ring  $R$  is of index at most  $n$ , then the center  $Z(R)$  of  $R$  is a  $\pi$ -regular ring of index at most  $n$ .*

*Proof.* Let  $a$  be an element of  $Z(R)$ . Consider the ideal  $I = a^{n+1}R$ . Then  $a+I$  is a nilpotent element of the ring  $R/I$ . Since  $R/I$  is of index at

most  $n$  by hypothesis,  $a^n \in I$ . Then there exists an element  $b \in R$  such that  $a^n = a^{2n}b$ . Hence  $a^n$  is strongly regular, and so by [2, Lemma 1] there exists  $z \in Z(R)$  such that  $a^n = a^{2n}z$ . This proves that  $Z(R)$  is a  $\pi$ -regular ring of bounded index.

**Corollary 1.** *Let  $R$  be a PI-ring. If every factor ring of  $R$  is of index at most  $n$ , then  $R$  is a  $\pi$ -regular ring of index at most  $n$ .*

*Proof.* By [5, Theorem 2.3] it suffices to prove that every prime factor ring of  $R$  is a simple Artinian ring. So, without loss of generality, we may assume that  $R$  is a prime ring. By Proposition 1 the center  $Z(R)$  of  $R$  is  $\pi$ -regular, and so  $Z(R)$  is a field. Hence  $R$  is a simple Artinian ring by [10, Corollary to Theorem 2].

To characterize a  $\pi$ -regular ring of bounded index in this direction, we need consider the endomorphism rings of cyclic modules.

**Theorem 1.** *The following statements are equivalent :*

- 1)  $R$  is a  $\pi$ -regular ring of index at most  $n$ .
- 2) For any cyclic module  $M$ ,  $i(\text{End}_R(M)) \leq n$ .

*In this case, for any module  $N$  generated by  $m$  elements, it holds that  $i(\text{End}_R(N)) \leq i(M_m(R))$ .*

*Proof.* Suppose that 1) holds. Let  $M$  be a cyclic right  $R$ -module. Then  $M$  is isomorphic to  $R/K$  for some right ideal  $K$  of  $R$ . Let  $I_R(K)$  denotes the idealizer of  $K$  in  $R$ . Then  $\text{End}_R(M)$  is isomorphic to the ring  $I_R(K)/K$ . We claim that  $I_R(K)/K$  is of index at most  $n$ . Let  $a$  be an element of  $I_R(K)$  with  $a^p \in K$  for some positive integer  $p$ . Since  $R$  is a  $\pi$ -regular ring of index at most  $n$ , there exists  $x \in R$  such that  $a^n = a^{n+1}x$ . If  $p > n$ , then  $a^{p-1} = a^p x \in K$ , because  $K$  is a right ideal of  $R$ . Continuing this process, we obtain that  $a^n \in K$ .

Let  $N$  be a right  $R$ -module generated by  $a_1, a_2, \dots, a_m$  and let  $K = \{z \in M_m(R) \mid (a_1, \dots, a_m)z = (0, \dots, 0)\}$ . Given  $g \in \text{End}_R(N)$ , we can write  $g(a_1, \dots, a_m) = (a_1, \dots, a_m)(r_{ij})$  for some  $(r_{ij}) \in I_{M_m(R)}(K)$ . Then the map  $\phi: \text{End}_R(N) \rightarrow I_{M_m(R)}(K)/K$  defined by  $\phi(g) = (r_{ij}) + K$  is a ring isomorphism. By [11, Theorem 5]  $M_m(R)$  is also a  $\pi$ -regular ring of bounded index. Hence, by the same way as above, we can prove that the index of  $I_{M_m(R)}(K)/K$  is less than or equal to  $i(M_m(R))$ .

2)  $\Rightarrow$  1). Let  $a$  be an element of  $R$ . Consider the cyclic module

$M = R/a^{n+1}R$ . Then  $\text{End}_R(M)$  is isomorphic to  $I_R(a^{n+1}R)/a^{n+1}R$ . Since  $a \in I_R(a^{n+1}R)$  and  $a^{n+1} \in a^{n+1}R$ , we conclude that  $a^n \in a^{n+1}R$ . This implies that  $R$  is a  $\pi$ -regular ring of index at most  $n$ .

A ring  $R$  is called a *strongly regular ring* if  $R$  is a von Neumann regular ring and  $R$  has no non-zero nilpotent element. As an immediate corollary of Theorem 1, we have

**Corollary 2.** *The following statements are equivalent :*

- 1)  $R$  is a strongly regular ring.
- 2) For any cyclic module  $M$ ,  $\text{End}_R(M)$  has no non-zero nilpotent elements.

In this case, for any module  $N$  generated by  $m$  elements, we have  $i(\text{End}_R(N)) \leq m$ .

Let  $R$  be a subring of a ring  $S$ . If  $S$  is finitely generated as a right  $R$ -module,  $S$  is called a *finite extension* of  $R$ .

**Corollary 3.** *Let  $S$  be a finite extension of a ring  $R$ . If  $R$  is a  $\pi$ -regular ring of bounded index, then so is  $S$ .*

*Proof.* Let  $M$  be a cyclic right  $S$ -module. Let  $S$  be generated by  $m$  elements as a right  $R$ -module. Then  $M$  is generated by  $m$  elements as a right  $R$ -module. By Theorem 1,  $i(\text{End}_R(M)) \leq i(M_m(R))$ , and by [11, Theorem 5]  $n = i(M_m(R))$  is finite. Since  $\text{End}_S(M)$  is a subring of  $\text{End}_R(M)$ , we obtain that  $i(\text{End}_S(M)) \leq n$ . Again, by Theorem 1,  $S$  is a  $\pi$ -regular ring of index  $i(S) \leq n$ .

We shall sharpen [7, Theorem 2.3].

**Proposition 2.** *Let  $R$  be a  $\pi$ -regular ring of index  $n$  and let  $P$  be a prime ideal of  $R$ . Then  $R/P$  is isomorphic to a matrix ring  $M_k(D)$  for some division ring  $D$  and some  $k \leq n$ .*

*Proof.* Since  $S = R/P$  is  $\pi$ -regular, the Jacobson radical  $J$  of  $S$  is a nil ideal. Hence  $x^n = 0$  for all  $x \in J$ . Since  $S$  is a prime ring,  $J = 0$  by [6, Lemma 1.1]. Moreover, by [8, Lemma 2],  $S$  has no infinite set of orthogonal idempotents. By [7, Theorem 2.1]  $S$  is a simple Artinian ring (of index at most  $n$ ), and hence  $S$  is isomorphic to a matrix ring  $M_k(D)$  for some division ring  $D$  and some  $k \leq n$ .

Combining this proposition with [5, Theorem 2.1], we obtain

**Theorem 2.** *Let  $R$  be a ring of bounded index. Then the following statements are equivalent :*

- 1)  $R$  is  $\pi$ -regular.
- 2) All prime factor rings of  $R$  are Artinian.

According to [5, Theorem 2.3], a PI-ring  $R$  is  $\pi$ -regular if and only if all prime factor rings of  $R$  are Artinian. We shall show that a similar result of Corollary 3 holds for rings all of whose prime factor rings are Artinian. We say  $R$  is *strongly  $\pi$ -regular* if for each  $a \in R$  there exists a positive integer  $n$  such that  $a^n R = a^{n+1} R$ . By [4, Théorème 1] this definition is left-right symmetric, and so such a ring is  $\pi$ -regular. The following is an easy consequence of [1, Theorem 1.1].

**Proposition 3.** *The following statements are equivalent :*

- 1) For each  $n \geq 1$ ,  $M_n(R)$  is strongly  $\pi$ -regular.
- 2) Every finite extension of  $R$  is strongly  $\pi$ -regular.

Let  $R$  be a ring whose prime factor rings are Artinian. By virtue of [5, Theorem 2.1],  $R$  satisfies the condition 1) of Proposition 3. Hence we have

**Corollary 4.** *Let  $R$  be a ring whose prime factor rings are Artinian. Then every finite extension of  $R$  is strongly  $\pi$ -regular.*

Let  $S$  be a finite extension of a ring  $R$ . If all prime factor rings of  $S$  are Artinian, is  $R$  strongly  $\pi$ -regular? We shall show that this is true when  $S$  is a finite normalizing extension of  $R$ . Recall that  $S$  is called a *finite normalizing extension* of  $R$  if there exists finitely many elements  $x_1, x_2, \dots, x_n$  in  $S$  such that  $S = Rx_1 + Rx_2 + \dots + Rx_n$  and  $Rx_i = x_i R$  for all  $i = 1, 2, \dots, n$ .

**Proposition 4.** *Let  $S$  be a finite normalizing extension of a ring  $R$ . If  $S$  is a ring whose prime factor rings are Artinian, then so is  $R$ .*

*Proof.* Let  $P$  be a prime ideal of  $R$ . By [3, Theorem 2.3] there is a prime ideal  $Q$  of  $S$  such that  $P$  is one of the minimal primes over  $Q \cap R$ . By hypothesis,  $S/Q$  is a simple Artinian ring. Since  $S/Q$  is a finite normalizing extension of  $R/(Q \cap R)$ ,  $R/(Q \cap R)$  is right Artinian by [9, Proposition 5 (iii)]. Since  $Q \cap R$  is semiprime,  $R/(Q \cap R)$  is a semisimple

Artinian ring. Then  $R/P$  is isomorphic to one of the simple components of  $R/(Q \cap R)$ . This proves that  $R/P$  is a simple Artinian ring.

By virtue of Theorem 2, we have

**Corollary 5.** *Let  $S$  be a finite normalizing extension of a ring  $R$ . If  $S$  is a  $\pi$ -regular ring of bounded index, then so is  $R$ .*

**Acknowledgment.** During the summer of 1989 the author was a visitor at The Pusan National University, where part of this work was done. The author thanks the Department of Mathematics Education for their hospitality.

#### REFERENCES

- [1] E. P. ARMENDARIZ, J. W. FISHER and R. L. SNIDER : On injective and surjective endomorphisms of finitely generated modules, *Comm. Algebra* **6** (1978), 659–672.
- [2] G. AZUMAYA : Strongly  $\pi$ -regular rings, *J. Fac. Sci. Hokkaido Univ.* **13** (1954), 34–39.
- [3] J. BIT-DAVID and J. C. ROBSON : Normalizing extensions I, *Lecture Notes in Math.* **825** (1980), 1–5, Springer-Verlag.
- [4] F. DISCHINGER : Sur les anneaux fortement  $\pi$ -réguliers, *C. R. Acad. Sci. Paris* **283 A** (1976), 571–573.
- [5] J. W. FISHER and R. L. SNIDER : On the von Neumann regularity of rings with regular prime factor rings, *Pacific J. Math.* **54** (1974), 135–144.
- [6] I. N. HERSTEIN : *Topics in Ring Theory*, Univ. of Chicago Press, 1969.
- [7] I. KAPIANSKY : Topological representations of algebras, II, *Trans. Amer. Math. Soc.* **68** (1950), 62–75.
- [8] M. OHORI : On non-commutative generalized p. p. rings, *Math. J. Okayama Univ.* **26** (1984), 157–167.
- [9] R. RESCO : Radicals of finite normalizing extensions, *Comm. Algebra* **9** (1981), 713–725.
- [10] L. H. ROWEN : Some results on the center of a ring with polynomial identity, *Bull. Amer. Math. Soc.* **79** (1973), 219–223.
- [11] H. TOMINAGA : Some remarks on  $\pi$ -regular rings of bounded index, *Math. J. Okayama Univ.* **4** (1955), 135–141.

DEPARTMENT OF MATHEMATICS  
OKAYAMA UNIVERSITY  
OKAYAMA, 700 JAPAN

(Received November 2, 1989)