

Mathematical Journal of Okayama University

Volume 25, Issue 2

1983

Article 4

DECEMBER 1983

Some remarks on the set of idempotents and the set of elements square zero

Yasuyuki Hirano^{*}

Hisao Tominaga[†]

^{*}Okayama University

[†]Okayama University

Copyright ©1983 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

SOME REMARKS ON THE SET OF IDEMPOTENTS AND THE SET OF ELEMENTS SQUARE ZERO

YASUYUKI HIRANO and HISAO TOMINAGA

Throughout the present paper, R will represent an (associative) ring different from 0. Given a subset S of R , we denote by $\langle S \rangle$ the subring of R generated by S . Let E be the set of idempotents in R , and N the set of nilpotents in R . We set $N^* = \{x \in R \mid x^2 = 0\}$. We say R is *normal* if and only if E is contained in the center of R : $[E, R] = 0$. As is well known, if $[E, E] = 0$ or $[E, N^*] = 0$ then R is normal. But, $[N^*, N^*] = 0$ need not yield the normality of R . If every left (resp. right) ideal of R is idempotent, R is said to be *fully left* (resp. *right*) *idempotent*. Finally, R is called a *chain ring* if the lattice of ideals of R forms a chain.

In §1, we investigate the structure of π -regular rings with only finitely many idempotents (Theorem 1), which leads us to the main theorem of Putcha and Yaqub [7]. Also, we give the conditions for an algebra generated by idempotents over a commutative regular ring to be commutative (Theorem 2). In §2, our interest will be directed towards to rings whose elements square zero commute with each other. First, such a semiprime P.I. ring is shown to be reduced (Theorem 3). Next, by making use of a generalization of the corrected statement of [8, Proposition 4], we prove that such a fully left and right idempotent chain ring is a simple ring without nontrivial idempotents (Theorem 4).

1. We state first the following

Theorem 1. *Let R be a ring in which every non-nil right ideal contains a nonzero idempotent. If E is finite then either R is a nil ring or the residue class ring $\bar{R}$ of R modulo its (Jacobson) radical J is a finite direct sum of finite simple rings and/or division rings.*

Proof. If $R = J$ then R is obviously a nil ring. If $R \neq J$ then, in view of [4, Theorem 2.1], we see that $\bar{R}$ is Artinian (semiprimitive) and, needless to say, every simple constituent of $\bar{R}$ is either finite or a division ring.

Corollary 1 ([7, Theorem 1]). *If $R \setminus N$ is finite, then R is either finite or nil (and conversely).*

Proof. Obviously, R is strongly π -regular. Hence the radical J of R is a nil ideal, and $\bar{R}=R/J$ also has only a finite number of nonnilpotents. Suppose $\bar{R}$ is nonzero. Then it is easy to see that J is finite. On the other hand, by Theorem 1, $\bar{R}$ is a finite semiprimitive ring, and therefore R is finite.

Remark 1. On contrast with Corollary 1, in an unpublished paper, M.S. Putcha and A. Yaqub proved that if $R \setminus E$ is finite then R is either finite or Boolean (and conversely).

In advance of stating the second corollary to Theorem 1, we prove the next, which concerns [10, Problem 94].

Proposition 1. *The following statements are equivalent :*

- 1) R has a finite subset F such that every nonzero right ideal of R has a nonzero (nonempty) intersection with F .
- 2) The right socle S of R is essential in R_R and the lattice $\mathcal{L}(S_R)$ of its submodules is finite. When this is the case, $\bar{S}=(S+P)/P$ is a finite direct sum of finite simple rings and/or division rings, where P is the prime radical of R .

Proof. It suffices to show that 1) implies 2). We may assume that F does not contain 0. For any $f \in F$, we denote by M_f the least right ideal of R containing f . After suitable arrangement of F , we may further assume that $M_f \cap M_{f'} = 0$ provided $f \neq f'$. Then it is easy to see that M_f ($f \in F$) are the minimal right ideals of R .

Let $\bar{R}=R/P$. Since $\bar{S}_R$ is finitely generated, the ideal $\bar{S}$ is a direct summand of $\bar{R}_R$, and therefore $\bar{S}$ is a ringtheoretic direct summand of the semiprime ring $\bar{R}$. This proves that $\bar{S}$ is an Artinian semiprimitive ring and $\mathcal{L}(\bar{S}_R)$ is finite. Consider now the $n \times n$ full matrix ring $T = \sum_{i,j=1}^n D e_{ij}$ over a division ring D ($n > 1$). As is easily seen, the different idempotents $e_{11} + d e_{21}$ ($d \in D$) generate different right ideals of T . This enables us to see that $\bar{S}$ is a finite direct sum of finite simple rings and/or division rings.

Combining Proposition 1 with Theorem 1, we readily obtain

Corollary 2. *The following are equivalent :*

- 1) R is a semiprime ring and contains a finite subset F such that every nonzero right ideal of R has a nonzero intersection with F .
- 2) R is a semiprimitive π -regular ring and E is finite.
- 3) R is a finite direct sum of finite simple rings and/or division rings.

Next, we consider the commutativity conditions for certain algebras generated by its idempotents.

Proposition 2. *If $R = \langle E \rangle$ and is commutative, then R is a subdirect sum of R_α ($\alpha \in A$), where $R_\alpha = \mathbb{Z}$ or $\mathbb{Z}/p_\alpha^{k_\alpha}\mathbb{Z}$ with some prime p_α and $k_\alpha > 0$.*

Proof. We may, and shall, assume that R is subdirectly irreducible. Then, it is easy to see that R has 1 as the unique nonzero idempotent. If $N=0$, then we see that R is either $\mathbb{Z}$ or $\mathbb{Z}/p\mathbb{Z}$ with some prime p . On the other hand, if $N \neq 0$, then we can easily see that the additive order of 1 is p^k for some prime p and $k > 1$. Hence $R = \mathbb{Z}/p^k\mathbb{Z}$.

Theorem 2. *Let K be a commutative regular ring with 1. If a K -algebra R is generated by E over K , then the following are equivalent :*

- 1) R is commutative.
- 2) R is a subdirect sum of fields homomorphic to K .
- 3) R is a reduced ring.
- 4) R is a strongly regular ring.

Proof. Obviously, 4) implies 3) and 2) does 3). Furthermore, as every idempotent in a reduced ring is central, 3) implies 1).

1) implies 2) and 4). Since every ideal of R is necessarily an algebra ideal, a slight modification of the proof of Proposition 2 proves 2), and so R is a reduced ring. Now, let M be a proper prime ideal of R . Then $\hat{R} = R/M$ is a domain generated by $\hat{E}$ over K . Hence, $\hat{R}$ has the identity ϵ as the unique nonzero idempotent, and $\hat{R} = K \cdot \epsilon$ is a field. Thus, M is a maximal ideal, and therefore R is a regular ring (see, e.g. [1, Theorem]).

2. We begin this section with the following

Theorem 3. *Let R be a semiprime P.I. ring. If $[N^*, N^*] = 0$ then R is a reduced ring.*

Proof. By [5, Theorem 5], R has a maximal left and right quotient ring Q which is a regular ring. It suffices to show that Q is normal. Let e be an arbitrary idempotent in Q . Then there exists an essential ideal A of R such that $eA(1-e) \subseteq R$ and $(1-e)Ae \subseteq R$ (see [5, Theorem 4]). By hypothesis, $eA(1-e)Ae = (1-e)AeA(1-e) = 0$. Since R is semiprime, we then have $eA(1-e) = 0 = (1-e)Ae$. This means that $[e, A] = 0$. Now, let x be an arbitrary element of Q , and choose an essential ideal B of R such

that $xB \subseteq R$. We claim that $[e, x]BA = 0$. In fact, if $a \in A$ and $b \in B$ then $exba = xbae = xe ba$. Hence, $[e, x] = 0$, proving the normality of Q .

The next corrects and generalizes [8, Proposition 4].

Proposition 3. (1) Let R be a fully left idempotent chain ring. If $I \neq R$ is an ideal of R then $I = (I \cap N^*)R$.

(2) Let R be a fully left and right idempotent chain ring. If $I \neq R$ is an ideal of R then $I = \langle I \cap N^* \rangle$.

Proof. (1) Let x be an arbitrary element of I . Then there exists $e \in RxR$ such that $x = ex$. Furthermore, we can find $f \in ReR \subseteq RxR$ such that $fe = e$. Since $f \in RfR \subseteq R(1-f)R$ by hypothesis, we can write $e = \sum_i r_i(1-f)s_i$ with some $r_i, s_i \in R$. Hence, we get

$$x = e^2x = \sum_i er_i(1-f)s_ix \in (I \cap N^*)R.$$

(2) Let x be an arbitrary element of I . Then there exists $e \in RxR$ such that $x = ex = xe$ (see the proof of [6, Lemma 1 (a)]). Furthermore, we can find $f \in ReR \subseteq RxR$ such that $e = fe = ef$. Noting that R is a chain ring and $I \neq R$, we can easily see that $RfR \subseteq R(1-f)^2R$. Thus we can write $e = \sum_i r_i(1-f)^2s_i$ with some $r_i, s_i \in R$, and therefore we get

$$x = e^2x = \sum_i \{er_i(1-f)\}\{(1-f)s_ix\} \in \langle I \cap N^* \rangle.$$

Lemma 1. (1) If $[N^*, N^*] = 0$, then $\langle N^* \rangle$ is a commutative nil subring. If, furthermore, R is semiprime, then $N^{*2} = 0$.

(2) If $e \in E$, then $eR(1-e)R \subseteq \langle (eR \cup Re) \cap N^* \rangle$.

Proof. (1) It suffices to prove the latter assertion. If $x, y \in N^*$, then $xyRxy = yRxyx = yRyx^2 = 0$, and hence $xy = 0$.

(2) In fact, for any $r, s \in R$ we have

$$er(1-e)s = er(1-e) \cdot (1-e)se + er(1-e)s(1-e).$$

We are now in a position to state the second main theorem of this section.

Theorem 4. Let R be a fully left and right idempotent chain ring. If $[N^*, N^*] = 0$, then R is a simple ring without nontrivial idempotents.

Proof. By Lemma 1 (1) and Proposition 3 (2), we see that R is a simple ring. Now, let e be a nonzero idempotent in R . Then, by Lemma 1 (2), $eR(1-e)R$ is a nilpotent right ideal, and hence $R(1-e)R =$

$ReR(1-e)R=0$, whence $eR=R=Re$ follows.

Corollary 3. *If R is a regular chain ring with $[N^*, N^*]=0$, then R is a division ring (and conversely).*

We conclude this paper with the following remarks.

Remark 2. If a regular chain ring R is a P.I. ring, then R is an Artinian simple ring. In fact, it is clear that the center of R is a field, and hence R is an Artinian simple ring by [9, Corollary 1.6.28]. Moreover, it is known that every regular ring R with $[N^*, N^*]=0$ is strongly regular (see [3, Lemma 2 (2)]).

Remark 3. A chain ring is called an n -chain ring if it has exactly n nontrivial ideals. Every simple ring with 1 is a fully left and right idempotent 0-chain ring. Given an algebra S with 1 over a field K , we denote

by $S^{(1)}$ the set consisting of the matrices having the form

$$\begin{bmatrix} A & & 0 \\ & k & \\ & & k \\ & & & \ddots \\ 0 & & & & \ddots \end{bmatrix}$$

where A is arbitrary finite matrix with entries in S and k is any element in K . If S is an n -chain ring then it is easy to see that the K -algebra $S^{(1)}$ is an $(n+1)$ -chain ring. According to [2], there is a simple algebra R over $\mathbb{Q}$ which is a domain (with 1) but not a division ring (compare Theorem 4 with Corollary 3). Obviously, $R^{(n)}=(R^{(n-1)})^{(1)}$ is a fully left and right idempotent n -chain ring which is not a regular ring.

REFERENCES

- [1] K. CHIBA and H. TOMINAGA: On strongly regular rings, Proc. Japan Acad. **49** (1973), 435—437.
- [2] J.H. COZZENS: Homological properties of the ring of differential polynomials, Bull. Amer. Math. Soc. **76** (1970), 75—79.
- [3] Y. HIRANO, H. TOMINAGA and A. YAQUB: On rings satisfying the identity $(x+x^2+\cdots+x^n)^{(n)}=0$, Math. J. Okayama Univ. **25** (1983), 13—18.
- [4] I. KAPLANSKY: Topological representation of algebras. II, Trans. Amer. Math. Soc. **68** (1950), 62—75.
- [5] W.S. MARTINDALE, 3RD: On semiprime P.I. rings, Proc. Amer. Math. Soc. **40** (1973), 365—369.
- [6] I. MOGAMI and M. HONGAN: Note on commutativity of rings, Math. J. Okayama Univ. **20** (1978), 21—24.

- [7] M.S. PUTCHA and A. YAQUB : Rings with a finite set of non-nilpotents, *Internat. J. Math. & Math. Sci.* 2 (1979), 121—126.
- [8] R. RAPHAEL : A note on ideal structure and unit generation in some fully idempotent and regular rings, *Kyungpook Math. J.* 17 (1977), 21—22.
- [9] L.H. ROWEN : *Polynomial Identities in Ring Theory*, Academic Press, New York, 1980.
- [10] F. SZÁSZ : *Radicals of Rings*, John Wiley & Sons, Chichester-New York-Brisbane-Toronto, 1981.

DEPARTMENT OF MATHEMATICS
OKAYAMA UNIVERSITY

(Received January 20, 1983)