# Mathematical Journal of Okayama University

Volume 27, Issue 1

1985 January 1985

Article 19

## Hadamard matrices of bush type

Noboru Ito<sup>\*</sup> Judith Q. Longyear<sup>†</sup>

\*Konan University <sup>†</sup>Wayne State University

Copyright ©1985 by the authors. *Mathematical Journal of Okayama University* is produced by The Berkeley Electronic Press (bepress). http://escholarship.lib.okayama-u.ac.jp/mjou

Math. J. Okayama Univ. 27 (1985), 127-133

### HADAMARD MATRICES OF BUSH TYPE

NOBORU ITO and JUDITH Q. LONGYEAR

In [1] Bush suggested a method for constructing a Hadamard matrix of order n using a Hadamard matrix of order  $\frac{1}{2}n-2$  and a skew Hadamard matrix of  $\frac{1}{4}n+1$ , where  $n \equiv 12 \pmod{16}$ . A Hadamard matrix of order n constructed by the method of Bush will be called a Hadamard matrix of Bush type of order n.

The purpose of this note is to prove two propositions on Hadamard matrices of Bush type of order n.

For basic facts on Hadamard matrices see  $\lfloor 2 \rfloor$ .

1. Introduction. We want to construct a Hadamard matrix of order n = 16u+12 under certain "inductive" assumptions, where u is a non-negative integer. Obviously it suffices to construct a symmetric 2-(16u+11, 8u+5, 4u+2) design D = (P, B), where  $P = \{1, 2, ..., 16u+11\}$  and B denote the sets of points and blocks of D respectively.

We make the following "inductive" assumptions: (1) There exists a Hadamard matrix L of order 8u+4, and (2) there exists a skew Hadamard matrix R of order 4u+4. Put  $L = (\lambda(i)), 1 \le i \le 8u+4$ , where  $\lambda(i)$  denotes the *i*-th row vector of L and we may assume that  $\lambda(1)$  is the all one vector. Let  $L(\lambda(1)) = (P(\ell), B(\ell))$  be the Hadamard 3-design associated with L at  $\lambda(1)$ . We put  $P(\ell) = \{1, 2, ..., 8u+4\}$  so that the block  $\sigma(i)$  of  $L(\lambda(1))$  corresponding to  $\lambda(i)$  contains the point j if and only if the j-th component of  $\lambda(i)$  equals 1, where  $2 \le i \le 8u+4$  and  $\sigma(i)^* = P(\ell) - \sigma(i)$  is also a block of  $L(\lambda(1))$ , for  $2 \le i \le 8u+4$ . Clearly we have that  $\sigma(i) \cap \sigma(i)^* = \emptyset$  and  $|\sigma(i) \cap \sigma(j)| = |\sigma(i) \cap \sigma(j)^*| = 2u+1$  for  $i \ne j$ .

We pick up any 4u+3 distinct disjoint block pairs from the  $\{\sigma(i), \sigma(i)^*\}, 2 \le i \le 8u+4$ . For simplicity of notation we denote them by  $\{\sigma(i), \sigma(i)^*\}, 2 \le i \le 4u+4$ . This configuration 2 consists of 8u+6 blocks of size 4u+2.

Next we may assume that R is in a skew-normalized skew form :

$$R = \begin{bmatrix} -1 & 1 & \cdots & 1 \\ -1 & -1 & & & \\ \vdots & & \vdots & & \\ \vdots & & & \ddots & \\ -1 & & & -1 \end{bmatrix} = (\rho(i)), \text{ where } \rho(i) \text{ denotes}$$

127

N. ITO and J. Q. LONGYEAR

128

the *i*-th row vector of R,  $1 \le i \le 4u+4$ . we label the *j*-th column of R by 8u+2j+2, for  $2 \le j \le 4u+4$ , and notice that the first column is still labelled 1.

Let D(r) = (P(r), B(r)) be a symmetric 2-(4u+3, 2u+2, u+1)design which is the complement of the symmetric 2-(4u+3, 2u+1, u)design associated with R at  $\rho(1)$  punctured at 1. We put  $P(r) = \{8u+6, 8u+8, ..., 16u+10\}$  so that the block  $\tau(i)$  of D(r) corresponding to  $\rho(i)$ contains the point 8u+2j+2 if and only if the j-th component of  $\rho(i)$  equals -1 ( $2 \le i, j \le 4u+4$ ). Let us define a mapping T from B(r) to P(r) by  $\tau(i)T = 8u+2i+2$ , for  $2 \le i \le 4u+4$ . Then by the skew property of Rwe have that  $\tau(i)T \in \tau(i)$  and that  $\tau(i)T \in \tau(j)$  if and only if  $\tau(j)T \notin$  $\tau(i)$  for  $i \ne j$ .

Now we are going to double points and blocks of D(r) as follows. The block  $\tau(i)$  will be developed into two blocks  $\tau(i1)$  and  $\tau(i2)$ ,  $2 \le i \le 4u +$ 4. If  $8u+2j+2 \in \tau(i)$  and  $i \ne j$ , then both  $\tau(i1)$  and  $\tau(i2)$  contain both 8u+2j+2 and 8u+2j+3. If i = j, then  $\tau(i1)$  contains only 8u+2i+2and  $\tau(i2)$  contains only 8u+2i+3. Then clearly we have that  $|\tau(i1) \cap$  $\tau(i2)| = 4u+2$ , for  $2 \le i \le 4u+4$ . Moreover, since  $|\tau(i) \cap \tau(j)| =$ u+1 and since  $\tau(i)T \in \tau(j)$  if and only if  $\tau(j)T \notin \tau(i)$  for  $i \ne j$ , we have that  $|\tau(ik) \cap \tau(jl)| = 2u+1$  for  $i \ne j$  and  $1 \le k, l \le 2$ . In this way we get a configuration  $\Re$  consisting of 8u+6 blocks of size 4u+3 =1+2(2u+1).

Finally we match  $\mathfrak{A}$  and  $\mathfrak{N}$  together in any possible way under the condition that  $\{\sigma(i), \sigma(i)^*\}$  and  $\{\tau(j1), \tau(j2)\}$  should be matched if a member of  $\{\sigma(i), \sigma(i)^*\}$  is matched together with a member of  $\{\tau(j1), \tau(j2)\}$ . For simplicity of notation we assume that  $\sigma(i)$  and  $\tau(i1)$ , and hence  $\sigma(i)^*$  and  $\tau(i2)$ , are matched together,  $2 \le i \le 4u+4$ .

Put  $\alpha(1) = P(\ell) \cup \{8u+5\}$ ,  $\alpha(2i-2) = \sigma(i) \cup \tau(i1)$  and  $\alpha(2i-1) = \sigma(i)^* \cup \tau(i2)$ , for  $2 \le i \le 4u+4$ . Then it is easy to see that  $|\alpha(i)| = 8u+5$ ,  $1 \le i \le 8u+7$  and  $|\alpha(i) \cap \alpha(j)| = 4u+2$  for  $i \ne j$ .

So the configuration  $\mathfrak{P} = (P, \{a(i)\}, 1 \le i \le 8u+7)$  is possibly a portion of a symmetric 2-(16u+11, 8u+5, 4u+2) design.

Now we prove the following proposition.

**Proposition 1.** A necessary and sufficient condition for  $\mathfrak{P}$  to be completed to a symmetric 2-(16u+11, 8u+5, 4u+2) design can be stated as follows.

There exist 8u+4 subsets  $\mu(j)$  of size 4u+2,  $1 \le j \le 8u+4$ , of  $\alpha(1)$ ,

#### HADAMARD MATRICES OF BUSH TYPE

called blocks again, such that  $D(\ell) = (\alpha(1), \{\sigma(i), \sigma(i)^*, 1 \le i \le 4u+3, \mu(j), 1 \le j \le 8u+4\})$  forms a 2-(8u+5, 4u+2, 4u+1, 16u+10, 8u+4) design, where the five parameters correspond to the usual notation  $v, k, \lambda, b$  and r respectively, with the following three conditions:

(1) Put  $\bar{\sigma}(i) = \sigma(i)$  or  $\sigma(i)^*$ ,  $1 \le i \le 4u+3$ . Then with any fixed  $\bar{\sigma}(i)$  one half of the  $\mu(k)$  intersects in 2u+1 points and the other half of the  $\mu(k)$  intersects in 2upoints.

(2) With each of any fixed  $\bar{\sigma}(i)$  and  $\bar{\sigma}(j)$  for  $i \neq j$  one quarter of the  $\mu(k)$  intersects in 2u+1 points and another quarter of the  $\mu(k)$  intersects in 2u points.

(3) Let a be a point such that  $1 \le a \le 8u+4$ . If a belongs to  $\overline{\sigma}(i)$ , then exactly 2u of the  $\mu(k)$  which intersects with  $\overline{\sigma}(i)$  in 2u points contain a. If a does not belong to  $\overline{\sigma}(i)$ , then exactly 2u+1 of the  $\mu(k)$  which intersects with  $\overline{\sigma}(i)$  in 2u points contain a.

*Proof.* Necessity. Suppose that  $\mathfrak{P}$  is completed to a symmetric 2-(16u+11, 8u+5, 4u+2) design *D*. New blocks will be denoted by a(i), for  $8u+8 \le i \le 16u+11$ . Put  $\mu(i-8u-7) = \alpha(1) \cap \alpha(i)$  for  $8u+8 \le i \le 16u+11$ . Then  $D(\ell) = (\alpha(1), \ |\sigma(i), \ \sigma(i)^*, \ 1 \le i \le 4u+1, \ \mu(j), \ 1 \le j \le 8u+4\}$ ) is a 2-(8u+5, 4u+2, 4u+1, 16u+10, 8u+4) design. In fact, let *a* and *b* be any two distinct points of  $\alpha(1)$ . Then *a* belongs to 8u+5 blocks of *D* including  $\alpha(1)$  and  $\{a, b\}$  is contained in 4u+2 blocks of *D* including  $\alpha(1)$ . Hence *a* belongs to 8u+4 blocks of  $D(\ell)$  and  $\{a, b\}$  is contained in 4u+1 blocks of  $D(\ell)$ .

If a(8u+7+k),  $1 \le k \le 8u+4$ , contains both 8u+2i+2 and 8u+2i+3, where  $2 \le i \le 4u+4$ , or if it contains neither 8u+2i+2 nor 8u+2i+3, then  $a(8u+7+k) \cap \tau(i1) = a(8u+7+k) \cap \tau(i2)$ . Put  $|a(8u+7+k) \cap \tau(i1)| = x$ . Then  $4u+2 = |a(2i-2) \cap a(8u+7+k)| = |\sigma(i) \cap \mu(k)| + x$ . Every  $\mu(k)$  contains the point 8u+5. So  $|\sigma(i) \cap \mu(k)| + |\sigma(i)^* \cap \mu(k)| = 4u+1$ . Hence we have a contradiction that 4u+3 = 2x. Thus we have that  $||a(8u+7+k) \cap \tau(i1)| - |a(8u+7+k) \cap \tau(i2)|| = 1$ , and that  $|\sigma(i) \cap \mu(k)| = 2u$  or 2u+1. Let *E* and *F* be the numbers of the  $\mu(k)$  such that  $|\sigma(i) \cap \mu(k)| = 2u$  or 2u+1. Let *E* and *F* be the numbers of the  $\mu(k)$  such that  $|\sigma(i) \cap \mu(k)| = 2u$  and 2u+1 respectively. Since every point of  $\sigma(i)$  belongs to 4u+1 of the  $\mu(k)$ , we have that (4u+2)(4u+1) = 2u E+(2u+1)F. Then *E* is a multiple of 2u+1 and this fact implies that E = F = 4u+2 proving (1).

129

N. ITO and J. Q. LONGYEAR

130

We notice that  $|\alpha(8u+7+k) \cap \sigma(i)| = 2u+1$  if and only if  $8u+2i+2 \notin \alpha(8u+7+k)$ . Let  $2 \leq i \neq j \leq 4u+4$ . Then, since D(r) is a symmetric 2-(4u+3, 2u+2, u+1) design, there exist 2(u+1)-1 = 2u + 1 of the  $\tau(\ell 1)$  and  $\tau(\ell 2)$  containing the points 8u+2i+2 and 8u+2j+2. So 4u+2-(2u+1) = 2u+1 of the  $\alpha(8u+7+k)$  contain the points 8u+2i+2 and 8u+2j+2, proving (2).

Let  $a \in \sigma(i)$ , for  $1 \le a \le 8u+4$ . Now there exist exactly 2(2u+2) - 1 = 4u+3 of the  $\tau(jk)$  containing the point 8u+2i+2. So there exist exactly (2u+1)+1 = 2u+2 of the  $\alpha(\ell)$  with  $\ell \le 8u+7$  containing both a and 8u+2i+2. Hence there exist exactly 4u+2-(2u+2) = 2u of the  $\alpha(\ell)$  with  $\ell \ge 8u+8$  containing both a and 8u+2i+2. These are the blocks  $\alpha(\ell)$  with  $\ell \ge 8u+8$  intersecting with  $\sigma(i)$  in 2u points. The rest is similar. This proves (3).

Sufficiency. Suppose that we have a 2-(8u+5, 4u+2, 4u+1, 16u+10, 8u+4) design  $D(\ell)$  satisfying (1), (2) and (3).

Clearly  $\mu(k)$  contains the point 8u+5, for  $1 \le k \le 8u+4$ . Since  $\sigma(i) \cup \sigma(i)^* = \alpha(1) - |8u+5|$ , we have that  $|\sigma(i) \cap \mu(k)| = 2u+1$  or 2u according as  $|\sigma(i)^* \cap \mu(k)| = 2u$  or 2u+1 respectively, for  $2 \le i \le 4u+4$  and  $1 \le k \le 8u+4$ .

We form a configuration consisting of 8u+4 blocks  $\{\nu(1), ..., \nu(8u+4)\}$ of size 4u+3 based on the set of points  $\{8u+6, 8u+7, ..., 16u+11\}$ .  $\nu(k)$ contains the point 8u+2+2i or 8u+3+2i according as  $|\sigma(i) \cap \mu(k)| =$ 2u or  $|\sigma(i)^* \cap \mu(k)| = 2u$  respectively, for  $2 \le i \le 4u+4$  and  $1 \le k \le$ 8u+4. Since  $\nu(k)$  contains exactly one point of  $\{8u+2+2i, 8u+3+2i\}$ for each *i*, such that  $2 \le i \le 4u+4$ , the size of  $\nu(k)$  equals 4u+3.

We put  $\alpha(8u+7+j) = \mu(j) \cup \nu(j)$ , for  $1 \le j \le 8u+4$ , and let  $B = \{\alpha(1), \alpha(2), ..., \alpha(16u+11)\}$ . Then we show that D = (P, B) is a symmetric 2-(16u+11, 8u+5, 4u+2) design.

First we show that D is a 1-design. Let a be a point. If  $1 \le a \le 8u+5$ , then, since  $D(\ell)$  has replication number 8u+4 and since a belongs to  $\alpha(1)$ , a belongs to (8u+4)+1 = 8u+5 blocks of B. So let  $8u+6 \le a \le 16u+11$ . Now every point of D(r) belongs to 2u+2 blocks. One of these blocks say  $\tau(i)$ , contains a or a-1 as  $\tau(i)T$ . So there exists 2(2u+1)+1 = 4u+3 blocks  $\alpha(i)$  with  $i \le 8u+7$  containing a. Now by assumption (1) on  $D(\ell)$  there exist exactly 4u+2 of the  $\mu(k)$  such that  $|\sigma(i) \cap \mu(k)| = 2u$  or  $|\sigma(i)^* \cap \mu(k)| = 2u$ , according as a is even or odd respectively. So there exist 4u+2 blocks  $\alpha(i)$  with  $i \le 8u+8$  containing a.

Next we show that D is a 2-design. Let a and b be two distinct points.

#### HADAMARD MATRICES OF BUSH TYPE

If  $1 \le a, b \le 8u+5$ , then since  $D(\ell)$  is a 2-(8u+5, 4u+2, 4u+1, 16u)+10, 8u+4) design and since both a and b belong to  $\alpha(1)$ , a and b belong to (4u+1)+1 = 4u+2 blocks of B. Let  $8u+6 \le a, b \le 16u+11$ . If  $|a,b| = |8u+6, 8u+7|, |8u+8, 8u+9|, \dots, \text{ or } |16u+10, 16u+11|, \text{ then}$ we may assume that a is even. Only blocks  $\alpha(i)$  with  $2 \le i \le 8u+7$  may contain  $\{a, b\}$ . Since the replication number of D(r) is 2u+2, and since a appears in exactly one of the  $\tau(i)$  as  $\tau(i)T$ ,  $\{a, b\}$  is contained in 2(2u +(2-1) = 4u+2 blocks of B. If  $\{a, b\} \neq \{8u+6+2i, 8u+7+2i\}, 0 \le i$  $\leq 4u+2$ , then it suffices to consider the case where a and b are even. Then  $\{a, b\}$  is contained in exactly u+1 blocks of D(r). By the skew property of T exactly one of these blocks of D(r), say  $\tau(j)$ , contains a or b as  $\tau(j)T$ . So exactly 1+2(u+1-1) = 2u+1 blocks  $\alpha(i)$  with  $i \le 8u+7$ contain  $\{a, b\}$ . By assumption (2) on  $D(\ell)$  and by the definition of  $\nu(k)$ , exactly 2u+1 of the  $\nu(k)$  contain  $\{a, b\}$ . So exactly 2u+1 blocks  $\alpha(i)$  with  $i \ge 8u+8$  contain  $\{a, b\}$ . Finally let  $1 \le a \le 8u+5$  and 8u+5 and 8u+6 $\leq b \leq 16u+11$ . If a = 8u+5, then a belongs to all of the  $\mu(k)$ ,  $1 \leq k \leq 16u+11$ . 8u+4. By assumption (3) on  $D(\ell)$ , b belongs to exactly 4u+2 of the  $\mu(k)$ . So  $\{a, b\}$  is contained in exactly 4u+2 blocks of B. Thus we may assume that  $1 \le a \le 8u + 4$ . Again we may assume that b is even. Now b belongs to exactly 2u+2 blocks of D(r) and only one of these blocks, say  $\tau(k)$ , contain b as  $\tau(k)T$ . Therefore 2u+1 pairs of blocks  $\tau(ij)$  contain b, and  $\tau(k1)$ , not  $\tau(k2)$ , contains b. So if a belongs to  $\sigma(k)$ , then exactly 2u+1blocks  $\alpha(i)$  with  $i \leq 8u+7$  contain |a, b|. But if a belongs to  $\sigma(k)^*$ , then exactly 2u blocks  $\alpha(i)$  with  $i \leq 8u+7$  contain  $\{a, b\}$ . Then by assumption (3) on  $D(\ell)$  exactly 2u or 2u+1 blocks  $\alpha(i)$  with  $i \ge 8u+8$  contain  $\{a, b\}$ according as a belongs to  $\sigma(1)$  or  $\sigma(1)^*$ . This completes the proof.

**Definition.** We call a symmetric 2-(16u+11, 8u+5, 4u+2) design *D* thus constructed a Hadamard design of Bush type. Furthermore we call a Hadamard matrix of order 16u+12 associated with *D* a Hadamard matrix of Bush type.

**Remark 1.** The main point of proposition 1 is the fact that the construction of a Hadamard matrix of Bush type of order 16u+12 is reduced to the construction of a 2-(8u+5, 4u+2, 4u+1, 16u+10, 8u+4) design satisfying (1), (2) and (3) for which 8u+6 blocks are predetermined.

Remark 2. There exists some freedom to construct Hadamard matri-

#### 132 N. ITO and J. Q. LONGYEAR

ces of Bush type of order 16u+12: (i) The choice of a Hadamard matrix H of order 8u+4; (ii) The choice of 4u+4 rows from H; (iii) The choice of a skew Hadamard matrix of order 4u+4; (iv) The choice of the mapping T; (v) The choice of 2-(8u+5, 4u+2, 4u+1, 16u+10, 8u+4) design and (vi) The choice of the matching between  $\mathfrak{L}$  and  $\mathfrak{N}$ .

Remark 3. For u = 0 it is very easy to write down a design of Bush type:  $\alpha(1) = \{1, 2, 3, 4, 5\}, \ \alpha(2) = \{1, 2, 6, 10, 11\}, \ \alpha(3) = \{3, 4, 7, 10, 11\}, \ \alpha(4) = \{1, 3, 6, 7, 8\}, \ \alpha(5) = \{2, 4, 6, 7, 9\}, \ \alpha(6) = \{1, 4, 8, 9, 10\}, \ \alpha(7) = \{2, 3, 8, 9, 11\}, \ \alpha(8) = \{1, 5, 7, 9, 11\}, \ \alpha(9) = \{2, 5, 7, 8, 10\}, \ \alpha(10) = \{3, 5, 6, 9, 10\}, \ and \ \alpha(11) = \{4, 5, 6, 8, 11\}.$  For u = 1 there are more than ten inequivalent Hadamard matrices of Bush type.

2. The purpose of this section is to prove the following proposition.

**Proposition 2.** The transpose of a Hadamard matrix of Bush type is of Bush type. More precisely, the dual of a Hadamard design of Bush type is of Bush type.

*Proof.* We use the notation in the proof of Proposition 1, and consider the dual  $D^{d}$  of the Hadamard design of Bush type in § 1, D = (P, B). It will suffice to recognize in  $D^{d}$  a configuration similar to  $\mathfrak{P} = (P, \{\alpha(i)\}, 1 \le i \le 8u+7)$ .

Let  $\beta(i)$  be the set of blocks of *B* containing the point *i* of *P*,  $1 \le i \le 16u+11$ . Let  $P^d$  and  $B^d$  denote the sets of points and blocks of  $D^d$  respectively. Then  $P^d = \{a(i), 1 \le i \le 16u+11\}$  and  $B^d = \{\beta(i), 1 \le i \le 16u+11\}$ .

Now the point  $\alpha(1)$ , the set of points  $\alpha(i)$  with  $8u+8 \le i \le 16u+11$ and the block  $\beta(8u+5) = \{\alpha(1), \alpha(i) \text{ with } 8u+8 \le i \le 16u+11\}$  play the roles of the point 8u+5,  $P(\ell)$  and the block  $\alpha(1)$  in D, respectively.

Furthermore,  $\beta(8u+5) \cap \beta(8u+2i)$  and  $\beta(8u+5) \cap \beta(8u+2i+1)$ , where  $3 \le i \le 4u+5$ , correspond to  $\sigma(i) = \alpha(1) \cap \alpha(2i-2)$  and  $\sigma(i)^* = \alpha(1) \cap \alpha(2i-1)$ , where  $2 \le i \le 4u+4$ , respectively. Lastly  $(P(r))^d = \{\alpha(2j), 1 \le j \le 4u+3\}, (P(r))^d \cap \beta(8u+2i)$  and  $T^d$  defined by  $((P(r))^d \cap \beta(8u+2i))T^d = \alpha(2i-4)$ , where  $3 \le i \le 4u+5$ , correspond to P(r);  $\tau(i)$  and T respectively, where  $2 \le i \le 4u+4$ .

The rest may be checked without difficulty.

#### HADAMARD MATRICES OF BUSH TYPE

133

#### References

- K. A. BUSH: On Hadamard embeddability, Linear Algebra and Its Applications 29 (1980), 39-52.
- [2] M. HALL, JR.: Combinatorial Theory, Blaisdell, 1967.

DEPARTMENT OF APPLIED MATHEMATICS KONAN UNIVERSITY KOBE 658, JAPAN DEPARTMENT OF MATHEMATICS WAYNE STATE UNIVERSITY DETROIT, MICHIGAN 48202, U. S. A.

(Received May 25, 1985)