

# *Mathematical Journal of Okayama University*

---

*Volume 15, Issue 2*

1971

*Article 6*

OCTOBER 1972

---

## Another proof of the invariance of Ulm's functions in commutative modular group rings

Paul F. Dubois\*

Sudarshan K. Sehgal†

\*University of Alberta

†University of Alberta

Copyright ©1971 by the authors. *Mathematical Journal of Okayama University* is produced by The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

# ANOTHER PROOF OF THE INVARIANCE OF ULM'S FUNCTIONS IN COMMUTATIVE MODULAR GROUP RINGS

PAUL F. DUBOIS and SUDARSHAN K. SEHGAL

In this note we give a short and natural proof of the following theorem due to Berman and Mollov [1] and May [2].

**Theorem.** *Let  $Z_p G$  be the group ring of a  $p$ -primary group  $G$  over  $Z_p$ , the field of  $p$  elements. Suppose  $\theta: Z_p G \cong Z_p H$ . Then  $G$  and  $H$  have the same Ulm's functions.*

The proof is a direct consequence of a lemma of Jennings [3] which we give below. First, we need some notation. We write all groups multiplicatively and define  $G^p = \{g \in G \mid g = x^p \text{ for some } x \in G\}$ . Inductively, for ordinals  $\beta$  we have

$$G^{p^{\beta+1}} = \left(G^{p^\beta}\right)^p \text{ and } G^{p^\beta} = \bigcap_{\alpha < \beta} G^{p^\alpha}$$

for  $\beta$  a limit ordinal.

If  $K$  is a subgroup of  $G$ , by  $\Delta(G; K)$  we mean the ideal of  $Z_p G$  generated by elements of the form  $1 - k$ ,  $k \in K$ . Sometimes we write  $\Delta(K)$  if the context is clear. We denote  $\{x \in K \mid x^p = 1\}$  by  $K[p]$ .

**Lemma.** *Let  $G$  be a  $p$ -primary abelian group and  $N$  a subgroup. Then*

- (1)  $G/G^p \cong \Delta(G)/\Delta^2(G)$ , and
- (2)  $N/N^p \cong \Delta(G; N)/\Delta(G) \cdot \Delta(G; N)$ .

*Proof.* Define  $\lambda: G \rightarrow \Delta(G)/\Delta^2(G)$  by  $\lambda(g) = g - 1 + \Delta^2(G)$ . Since

$$(*) \quad g_1 g_2 - 1 = (g_1 - 1) + (g_2 - 1) + (g_1 - 1)(g_2 - 1),$$

$\lambda$  is an epimorphism with kernel  $= \{g \in G \mid g - 1 \in \Delta^2(G)\} = G^p$  by Jennings [3], proving (1). Actually, Jennings proved this equality for finite groups but since in an equation  $g - 1 = \delta \in \Delta^2(G)$ , only a finite number of elements of  $G$  occur, his result is applicable to our case.

For the second part, define

$$\mu: N \rightarrow \Delta(G; N)/\Delta(G) \cdot \Delta(G; N) \text{ by } \mu(n) = \overline{n - 1}.$$

It follows from (\*) and

$$g(n-1) = n-1 + (g-1)(n-1)$$

that  $\mu$  is an epimorphism with kernel  $= \{n \mid n-1 \in \Delta(G) \cdot \Delta(G; N)\}$ . It remains to prove:

$$(**) \quad n-1 \in \Delta(G) \cdot \Delta(G; N) \Rightarrow n \in N^p.$$

Choose a transversal  $\{g_i\}$  of  $N$  in  $G$  with  $g_1=1$ . Define for  $g, n \in G$ ,  $\sigma(g, n) = n$  and extend this linearly to  $\sigma: Z_p G \rightarrow Z_p N$ . Now,

$$n-1 = \sum_i \gamma_i (n_i-1), \quad \gamma_i \in \Delta(G), \quad n_i \in N.$$

Therefore

$$n-1 = (n-1)^\sigma = \sum_i \gamma_i^\sigma (n_i-1) \quad \text{and} \quad n-1 \in \Delta^2(N; N).$$

Hence,  $n \in N^p$ . This proves (\*\*) and therefore (2).

**Remark.** The above lemma is a special case of a similar result that holds for arbitrary (not necessarily abelian or finite) groups. Also, there is a corresponding result for integral group rings (see, Sehgal [4]). For the purpose of this paper the above will suffice.

*Proof of Theorem.* Now, suppose  $\theta: Z_p G \cong Z_p H$ . We may assume here that  $\theta$  is normalized; if  $\theta(g) = \sum_{h \in H} \alpha_h h$  then  $\sum_{h \in H} \alpha_h = 1$ . By noting that  $\ell(g^p) = (\sum_{h \in H} \alpha_h h)^p = \sum_{h \in H} \alpha_h^p h^p$ , we have that  $\theta$  maps  $Z_p G^p$  isomorphically onto  $Z_p H^p$ . By a simple induction

$$\theta: Z_p G^{p^\beta} \cong Z_p H^{p^\beta} \quad \text{for all ordinals } \beta.$$

We show first that the finite Ulm invariants are equal. The  $i$ th Ulm invariant,  $i < \omega$  (the first limit ordinal), is the dimension of  $(G^{p^i})[p] / (G^{p^{i+1}})[p]$ . For convenience let us denote  $(G^{p^i})[p]$  by  $L_i$ .

By the lemma we have an isomorphism

$$L_i \cong \Delta(G; L_i) / \Delta(G) \cdot \Delta(G; L_i).$$

Under  $\theta$ ,  $\Delta(L_i)$  is isomorphic to  $\Delta(M_i)$  where  $M_i = (H^{p^i})[p]$ .

Thus we obtain for each  $i$  the commutative diagram below:

$$\begin{array}{ccccccc} L_i & \cong & \Delta(L_i) / \Delta(G) \Delta(L_i) & \cong & \Delta(M_i) / \Delta(H) \Delta(M_i) & \cong & M_i \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ L_{i+1} & \cong & \Delta(L_{i+1}) / \Delta(G) \Delta(L_{i+1}) & \cong & \Delta(M_{i+1}) / \Delta(H) \Delta(M_{i+1}) & \cong & M_{i+1} \end{array}$$

and thus  $L_i / L_{i+1} \cong M_i / M_{i+1}$ .

The observation that  $Z_p G^{p^\beta} \cong Z_p H^{p^\beta}$  allows us to conclude that even the transfinite Ulm invariants are equal.

REFERENCES

- [ 1 ] S. D. BERMAN and T. ZH. MOLLOV : On group rings of abelian  $p$ -groups of any cardinality, *Mat. Zam* **6** (1969), 381—392 (Translation *Math. Notes* **6** (1969), 686—692).
- [ 2 ] W. MAY : Commutative group algebras, *Trans. Amer. Math. Soc.* **136** (1969), 139—149.
- [ 3 ] S. A. JENNINGS : The structure of the group rings of a  $p$ -group over a modular field, *Trans. Amer. Math. Soc.* **50** (1941), 175—185.
- [ 4 ] S. K. SEHGAL : On the isomorphism of integral group rings II, *Can. J. Math.* **21** (1969), 1182—1188.

UNIVERSITY OF ALBERTA

*(Received February 19, 1972)*