

Mathematical Journal of Okayama University

Volume 4, Issue 2

1954

Article 7

MARCH 1955

A note on matrix rings

Hisao Tominaga*

*Okayama University

Copyright ©1954 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

A NOTE ON MATRIX RINGS

HISAO TOMINAGA

Let R be a ring with an identity. Often, R can be represented as the total matrix ring over an m -irreducible ring (Definition 1). For example, a ring with minimum condition and a homogeneous π -regular ring ([4, Theorem 4.2] or [5, Theorem 5.6])¹⁾ possess this property. For such rings, it seems to be of interest to investigate the problem concerning the uniqueness of the representations as total matrix rings over m -irreducible rings. In this note, we shall examine this problem for a special kind of rings.

Throughout the note, R will be a ring with an identity, and the terms "radical", "primitive ideal" will be in Jacobson's sense ([3]).

We shall begin our course by setting the following definition:

Definition 1. A ring R is said to be *m-irreducible* if R is not representable as the total $n \times n$ matrix ring over any ring with $n > 1$.

A P-ring ([6, Definition 4.1]) is m -irreducible. In general, if a non-zero homomorphic image of R is m -irreducible, then so is R .

Definition 2. Let N be the radical of a ring R . If R/N is a (finite or infinite) complete direct sum of total matrix rings over division rings: $\sum_{\sigma}^c (\bar{D}_{\sigma})_{n_{\sigma}}$, then R is said to be *weakly semi-primary*.

Lemma 1. If R is a weakly semi-primary ring, then

- i) $(R)_n$ is weakly semi-primary,
- ii) eRe is also weakly semi-primary, where e is an idempotent.

Proof. i) As is well-known, the radical of $(R)_n$ is $(N)_n$. Hence $R/N = \sum_{\sigma}^c (\bar{D}_{\sigma})_{n_{\sigma}}$ implies that $(R)_n/(N)_n \cong \sum_{\sigma}^c (\bar{D}_{\sigma})_{nn_{\sigma}}$, that is, $(R)_n$ is weakly semi-primary.

ii) The radical of eRe is $eNe = eRe \cap N$ ([4, Theorem 3.1]). Hence, $eRe/eNe \cong \bar{e}\bar{R}\bar{e} = \sum_{\sigma}^c \bar{e}(\bar{D}_{\sigma})_{n_{\sigma}}\bar{e} = \sum_{\sigma}^c \bar{e}_{\sigma}(\bar{D}_{\sigma})_{n_{\sigma}}\bar{e}_{\sigma}$, where $\bar{}$ will mean the residue class modulo N , and where $\bar{e}_{\sigma}$ will be the σ -component of $\bar{e}$. Clearly $\bar{e}_{\sigma}(\bar{D}_{\sigma})_{n_{\sigma}}\bar{e}_{\sigma}$ is simple and satisfies the minimum condition. These facts show that eRe is weakly semi-primary.

The following is an immediate consequence of Lemma 1:

Corollary. The total $n \times n$ matrix ring $(R)_n$ is weakly semi-primary if and only if R is so.

1) Numbers in brackets refer to the references cited at the end of this note.

Lemma 2. Let $\mathfrak{M} = \mathfrak{m}_1 \oplus \cdots \oplus \mathfrak{m}_n$
 $= \mathfrak{n}_1 \oplus \cdots \oplus \mathfrak{n}_n$

be two direct decompositions of a module $\mathfrak{M}$ with an operator domain $\mathcal{Q}$ into n isomorphic submodules. If the $\mathcal{Q}$ -endomorphism ring of $\mathfrak{m}_1$ is weakly semi-primary, then $\mathfrak{m}_1 \cong \mathfrak{n}_1$.

*Proof.*¹⁾ Let R be the $\mathcal{Q}$ -endomorphism ring of $\mathfrak{M}$, and let e_i and f_j be the projections onto $\mathfrak{m}_i$ and $\mathfrak{n}_j$ respectively. Then $R = Re_1 \oplus \cdots \oplus Re_n (\cong (e_i Re_i)_n)$, whence R is weakly semi-primary by Corollary to Lemma 1. Here we set $R/N = \sum_{\sigma}^c (\bar{D}_{\sigma})_{n_{\sigma}}$, N the radical of R . Next $\bar{e}_i$ and $\bar{f}_j$ will denote the residue classes of e_i and f_j modulo N respectively. Then,

$$\bar{e}_1 + \cdots + \bar{e}_n = \bar{f}_1 + \cdots + \bar{f}_n = \bar{1}.$$

To be easily seen, $\bar{e}_1$ is isomorphic to $\bar{f}_1$ modulo the primitive ideal $\sum_{\sigma \neq 1}^c (\bar{D}_{\sigma})_{n_{\sigma}}$ for each τ . Hence $\bar{e}_1$ is isomorphic to $\bar{f}_1$, whence e_1 is isomorphic to f_1 , that is, $\mathfrak{m}_1$ is isomorphic to $\mathfrak{n}_1$.

Considering R itself and the totality of left-multiplications of elements of R as $\mathfrak{M}$ and $\mathcal{Q}$ in Lemma 2 respectively, we obtain the next:

Corollary. Let $R = (R_1)_n = (R_2)_n$. If R is weakly semi-primary, then $R_1 \cong R_2$.

Definition 3. A weakly semi-primary ring is called a *semi-primary ring* if the radical of the ring is nil.

Now we are going to establish our principal theorem which can be considered as a generalization of the familiar structure theorem of Wedderburn.

Theorem. Let R be a semi-primary ring: $R/N = \sum_{\sigma}^c (\bar{D}_{\sigma})_{n_{\sigma}}$, N the radical, and let n be the greatest common divisor of n_{σ} 's. Then,

- i) R is the total $n \times n$ matrix ring over an m -irreducible ring,
- ii) $R = (R_1)_n = (R_2)_n$ implies that $n_1 = n_2$ and $R_1 \cong R_2$, where R_1 and R_2 are m -irreducible.

1) The present proof is essentially same with that of [7, Theorem 46.3].

2) Let S be a ring and M be its radical. Two idempotents e and f are called *isomorphic* (in S) if there exist two elements a and b such that $ab = e$ and $ba = f$. e and f are isomorphic if and only if the left [right] ideals Se [eS] and Sf [fS] are operator isomorphic ([2, pp. 527 - 528] or [7, Theorem 10.2]). If e and f are isomorphic modulo M , then e and f are virtually isomorphic ([7, Theorem 18.12]). In case S is the endomorphism ring of a module $\mathfrak{M}$, the isomorphism of e with f means the isomorphism of $\mathfrak{M}e$ with $\mathfrak{M}f$ ([7, p. 230]).

Proof. It is almost trivial that $R/N = (T)_n$ with a ring T . Moreover, as N is nil, we obtain that $R = (T_0)_n$ with some T_0 ([1, Theorem 1.21]). Clearly T_0 is m -irreducible. For, if not, n would not be the greatest common divisor of n_i 's. If $R = (R_1)_{n_1}$, R_1 m -irreducible, then n_1 divides n . If $n_1 < n$, then $(R_1)_{n_1} \cong ((T_0)_{n/n_1})_{n_1}$ implies that $(T_0)_{n/n_1} \cong R_1$ by Corollary to Lemma 2. But it contradicts with the m -irreducibility of R_1 . Hence, $n_1 = n$ and $R_1 \cong T_0$.

Remark. Let R be an FI₁-ring of which all the primitive images have the equal degree n . Then R is a total $n \times n$ matrix ring over a P-ring. If $R = (B)_m$, B m -irreducible, then $m = n$.¹⁾

REFERENCES

- [1] K. ASANO, Theory of rings and ideals, Tokyo, Kyōritsu-sha (1949), (in Japanese).
- [2] G. AZUMAYA, On generalized semi-primary rings and Krull-Remak-Schmidt's theorem, Jap. Journ. Math., 19 (1948), 525 - 547.
- [3] N. JACOBSON, The radical and semi-simplicity for arbitrary rings, Amer. Journ. Math., 67 (1945), 300 - 320.
- [4] I. KAPLANSKY, Topological representation of algebras, Trans. Amer. Math. Soc., 68 (1950), 62 - 75.
- [5] J. LEVITZKI, On the structure of algebraic algebras and related rings, Trans. Amer. Math. Soc., 74 (1953), 384 - 409.
- [6] ———, On P-soluble rings, Trans. Amer. Math. Soc., 77 (1954), 216 - 237.
- [7] T. NAKAYAMA and G. AZUMAYA, Algebra II. (Theory of rings), Tokyo, Iwanami (1954), (in Japanese).

DEPARTMENT OF MATHEMATICS,
OKAYAMA UNIVERSITY

(Received February 10, 1955)

1) Cf. [5, Theorem 3.2] and [5, Theorem 5.6].