

Mathematical Journal of Okayama University

Volume 4, Issue 2

1954

Article 1

MARCH 1955

Banach algebras generated by a bounded linear operator

Minoru Tomita*

*Okayama University

Copyright ©1954 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

BANACH ALGEBRAS GENERATED BY A BOUNDED LINEAR OPERATOR

MINORU TOMITA

§1. Banach algebras with generators.

In this paper we shall characterize the spectrum and the semi-spectrums of a bounded linear operator on a Banach space by a ring-theoretical method. And we shall determine all the type of Banach algebras generated by a fixed bounded linear operator in connection with its corresponding semi-spectrums.

Let $\mathbf{M}$ be a Banach algebra which contains the identity I . $\mathbf{M}$ is called *normalized* if $|I| = 1$ and $|AB| \leq |A||B|$. Put

$$\|A\| = \sup_{|x| \leq 1} |AX|,$$

then the norm $\|A\|$ is equivalent to the original norm $|A|$, and by this norm $\mathbf{M}$ is a normalized Banach algebra.

Hereafter we shall assume that $\mathbf{M}$ is normalized and commutative. Given an element A of $\mathbf{M}$ and given a complex rational function $r(z) = (\sum_i a_i z^i) / (\sum_j b_j z^j)$. If $\sum_j b_j A^j$ is invertible, the element $r(A) = (\sum_i a_i A^i) (\sum_j b_j A^j)^{-1}$ is called a *rational form*. The *spectrum* $\mathfrak{S}(\mathbf{M})$ of A is the set of all complex numbers z so that $A - zI$ fail to be invertible. The *resolvent set* of A is the complement of the spectrum in the complex Riemann sphere $\mathbb{R}_0$.

If $\mathbf{M}$ contains a fixed element A whose rational forms are contained everywhere densely in $\mathbf{M}$, then A is called a *generator* of $\mathbf{M}$. The *spectrum* of the algebra $\mathbf{M}$ is the space of all continuous linear functionals f on $\mathbf{M}$ such that $f(I) = 1$ and $f(XY) = f(X)f(Y)$.

A fundamental relationship between spectrums of generators and the spectrum of a Banach algebra will be observed in the sequel.

Theorem 1. *Let $\mathbf{M}$ be a normalized Banach algebra with the identity I and a generator A . Then the spectrum $\mathfrak{S}$ of A is a bounded closed set in the complex number field. In order that a complex number z belong to $\mathfrak{S}$, it is necessary and sufficient that $|r(A)| \geq |r(z)|$ hold for every rational form $r(A)$. Every rational form $r(A)$ satisfies*

$$\lim_{n \rightarrow \infty} |r^n(A)|^{\frac{1}{n}} = \sup_{z \in \mathfrak{S}} |r(z)|.$$

The spectrum $\mathfrak{G}$ of $\mathbf{M}$ is homeomorphic to $\mathfrak{S}$ by the mapping

$$x \in \mathfrak{G} \longrightarrow x(A) \in \mathfrak{S}.$$

Proof. The famous I. Gelfund's theorem states that:

If $\mathbf{M}$ is a commutative Banach algebra, the spectrum $\mathfrak{G}$ of $\mathbf{M}$ is a compact space. For every fixed $f \in \mathbf{M}$, $x \in \mathfrak{G} \rightarrow x(f)$ is a continuous mapping of $\mathfrak{G}$ in the complex field. And every $f \in \mathbf{M}$ satisfies $\lim |f^n|^{1/n} = \sup_{x \in \mathfrak{G}} |x(f)|$.

Now our Banach algebra $\mathbf{M}$ in question contains the generator A . Then especially $x \in \mathfrak{G} \rightarrow x(A)$ is a continuous mapping of $\mathfrak{G}$ in the complex number field. So the range $\mathfrak{G}(A)$ of this mapping is a bounded closed set. We show that this mapping $x \rightarrow x(A)$ is a homeomorphism between $\mathfrak{G}$ and $\mathfrak{G}(A)$. Assume that u and v be two elements in $\mathfrak{G}$ which satisfy $u(A) = v(A) = t$. Then every rational form $r(A)$ satisfies $u(r(A)) = v(r(A)) = r(t)$. Since the set of all rational forms $r(A)$ is everywhere dense in $\mathbf{M}$, u coincides with v . Hence $x \rightarrow x(A)$ is a one-to-one continuous mapping of the compact set $\mathfrak{G}$ on $\mathfrak{G}(A)$, i.e. a homeomorphism.

Next we show that $\mathfrak{G}(A)$ coincides with $\mathfrak{S}$. Let z be an element in $\mathfrak{S}$. Then $A - zI$ fails to be invertible, and the monomial ideal $(A - zI)\mathbf{M}$ of $\mathbf{M}$ is a proper ideal of $\mathbf{M}$. By the Gelfund's theorem there exists at least one element u in $\mathfrak{G}$ which satisfies $u(X) = 0$ for every $X \in (A - zI)\mathbf{M}$. Especially $u(A - zI) = 0$ and $z = u(A) \in \mathfrak{G}(A)$. Therefore $\mathfrak{G}(A)$ contains $\mathfrak{S}$.

Conversely let z be a complex number which does not belong to $\mathfrak{S}$. Then $A - zI$ is invertible, and

$$\sup_{u \in \mathfrak{G}(A)} |(u - z)^{-1}| = \sup_{x \in \mathfrak{G}} |x((A - zI)^{-1})| \leq |(A - zI)^{-1}| = d.$$

That is,

$$\inf_{u \in \mathfrak{G}(A)} |(u - z)| \geq d^{-1} > 0.$$

And $\mathfrak{G}(A)$ does not contain z . Hence $\mathfrak{G}(A)$ coincides with $\mathfrak{S}$.

Now our theorem is clear. In fact, $x \rightarrow x(A)$ is a homeomorphism between $\mathfrak{G}$ and $\mathfrak{S}$. Therefore we can determine $\mathfrak{G}$ as follows: for every $z \in \mathfrak{S}$ there exists a uniquely determined bounded linear functional z on $\mathbf{M}$ defined by $z(r(A)) = r(z)$ for every rational form $r(A)$. $\mathfrak{G}$ is the set of all such bounded linear functionals z . Thus $z \in \mathfrak{G}$ if

and only if $|\tau(A)| \geq |\tau(z)|$ for every rational form $\tau(A)$. And by the Gelfund's theorem,

$$\lim_{n \rightarrow \infty} |\tau^n(A)|^{1/n} = \sup_{x \in \mathfrak{S}} |x(\tau(A))| = \sup_{z \in \mathfrak{S}} |\tau(z)|.$$

q.e.d.

§2. The classification of congenetic subalgebras.

Let $\mathbf{M}$ be a normalized Banach algebra with the identity I and a fixed generator A . We shall now classify all the sub-algebras of $\mathbf{M}$ generated by the same A .

Theorem 2. *Let $\mathbf{M}$ be a normalized Banach algebra with the identity I and a fixed generator A . If $\mathbf{N}$ is a Banach sub-algebra of $\mathbf{M}$ which is also generated by A , then the spectrum $\mathfrak{S}(\mathbf{N})$ of A which corresponds with $\mathbf{N}$ contains the spectrum $\mathfrak{S}$ of A (which corresponds with $\mathbf{M}$). And $\mathfrak{S}(\mathbf{N}) - \mathfrak{S}$ is an open set.*

Conversely let $\mathfrak{Z}$ be a bounded closed set in the complex field containing $\mathfrak{S}$ such that $\mathfrak{Z} - \mathfrak{S}$ is open, then there exists a uniquely determined Banach sub-algebra $\mathbf{N}$ of $\mathbf{M}$ such that its corresponding spectrum $\mathfrak{S}(\mathbf{N})$ of A coincides with $\mathfrak{Z}$.

Proof. Let $\mathbf{N}$ be a Banach sub-algebra of $\mathbf{M}$ which is also generated by A . Then the resolvent set $\mathfrak{R}(\mathbf{N})$ of A which corresponds with $\mathbf{N}$ is contained in the resolvent set $\mathfrak{R}(\mathbf{M})$ of A which corresponds with $\mathbf{M}$. We show that $\mathfrak{R}(\mathbf{M}) - \mathfrak{R}(\mathbf{N})$ is open.

Let $\{z_n\}$ be a sequence in $\mathfrak{R}(\mathbf{N})$ which converges to a number z in $\mathfrak{R}(\mathbf{M})$. Then $(A - z_n I)^{-1}$ belongs to $\mathbf{N}$, and $(A - zI)^{-1}$ belongs to $\mathbf{M}$. In a suitable neighbourhood of z , $(A - wI)^{-1}$ is developed to the power series

$$(A - wI)^{-1} = \sum_{n=0}^{\infty} (w - z)^n (A - zI)^{-(n+1)}.$$

Therefore $(A - z_n I)^{-1}$ converges uniformly to $(A - zI)^{-1}$. This means $(A - zI)^{-1} \in \mathbf{N}$, and $z \in \mathfrak{R}(\mathbf{N})$. Thus $\mathfrak{R}(\mathbf{N})$ is relatively closed in $\mathfrak{R}(\mathbf{M})$, so that $\mathfrak{R}(\mathbf{M}) - \mathfrak{R}(\mathbf{N})$ is an open set. Hence $\mathfrak{S}(\mathbf{N}) = \mathfrak{R}_0 - \mathfrak{R}(\mathbf{N})$ is a bounded closed set which contains $\mathfrak{S} = \mathfrak{R}_0 - \mathfrak{R}(\mathbf{M})$. And $\mathfrak{Z} - \mathfrak{S} = \mathfrak{R}(\mathbf{M}) - \mathfrak{R}(\mathbf{N})$ is open. The former half of Theorem 2 is thus proved.

To prove the latter half of the Theorem, we shall prepare the next Lemma.

Lemma 1. *Let $\mathbf{N}$ be a Banach sub-algebra generated by A . In order that a rational form $\tau(A)$ exist and belong to $\mathbf{N}$, it is necessary*

and sufficient that $r(z)$ have no pole on the spectrum $\mathfrak{S}(\mathbf{N})$ of A corresponding with $\mathbf{N}$.

In fact, if $r(z)$ is a rational function which has no pole on $\mathfrak{X} = \mathfrak{S}(\mathbf{N})$, then $r(z) = p(z) \prod_{i=1}^n (z - a_i)^{-1}$, where $p(z)$ is a polynomial, and every a_i is not contained in $\mathfrak{X}$. Then $r(A) = p(A) \prod_{i=1}^n (A - a_i I)^{-1}$ exists and belongs to $\mathbf{N}$. Conversely if a rational form $r(A)$ belongs to $\mathbf{N}$, then $|r(A)| \geq \sup_{z \in \mathfrak{X}} |r(z)|$. Therefore $r(z)$ is bounded on $\mathfrak{S}(\mathbf{N})$, and has no pole on it. This concludes the Lemma.

We now turn to the proof of our Theorem. Let $\mathfrak{X}$ be a bounded closed set of complex numbers which contains $\mathfrak{S}$ and satisfies that $\mathfrak{X} - \mathfrak{S}$ is open. We denote by $\mathbf{N}$ the smallest Banach sub-algebra of $\mathbf{M}$ which contains all rational forms $r(A)$ for which corresponding rational functions $r(z)$ have no pole on $\mathfrak{X}$. $\mathbf{N}$ is clearly generated by A , and contains the identity. We show that the spectrum $\mathfrak{S}(\mathbf{N})$ of A which corresponds with $\mathbf{N}$ coincides with $\mathfrak{X}$. Let z be any complex number which does not belong to $\mathfrak{X}$. Then $(A - zI)^{-1}$ exists and belongs to $\mathbf{N}$. Therefore z does not belong to $\mathfrak{S}(\mathbf{N})$, so $\mathfrak{X}$ contains $\mathfrak{S}(\mathbf{N})$.

Next we show the converse. Let $r(z)$ be a rational function which has no pole on $\mathfrak{X}$. $r(z)$ is regular and one-valued in the interior of $\mathfrak{X}$. Then by the well-known maximal modulus theorem for analytic functions, the maximal value of $|r(z)|$ in $\mathfrak{X}$ is taken on the boundary $\mathfrak{B}$ of $\mathfrak{X}$. Since $\mathfrak{B} = \mathfrak{X} - \text{int } \mathfrak{X}$ is contained in $\mathfrak{S}$ (because $\mathfrak{X} - \mathfrak{S}$ is open by the assumption), we have

$$\sup_{z \in \mathfrak{X}} |r(z)| = \sup_{z \in \mathfrak{B}} |r(z)| = \sup_{z \in \mathfrak{S}} |r(z)|.$$

Consider a fixed $w \in \mathfrak{X}$, then for every rational form $r(A)$ of which $r(z)$ has no pole on $\mathfrak{X}$, we have

$$|r(A)| \geq \sup_{z \in \mathfrak{S}} |r(z)| = \sup_{z \in \mathfrak{X}} |r(z)| \geq |r(w)|.$$

Those rational forms $r(A)$ are contained everywhere densely in $\mathbf{N}$. By Theorem 1 we conclude that w belongs to $\mathfrak{S}(\mathbf{N})$. Therefore $\mathfrak{S}(\mathbf{N})$ contains $\mathfrak{X}$, and coincides with $\mathfrak{X}$.

Finally we show that the correspondence between Banach algebras $\mathbf{N}$ generated by the same A and the related spectrums $\mathfrak{S}(\mathbf{N})$ is one-to-one. Let $\mathbf{N}$ and $\mathbf{L}$ be two Banach sub-algebras of $\mathbf{M}$ which are generated by A and which have the common related spectrums

$\mathfrak{L} = \mathfrak{S}(\mathbf{N}) = \mathfrak{S}(\mathbf{L})$. We show that $\mathbf{L}$ and $\mathbf{N}$ coincide with each other. By Lemma 1 a rational form $r(A)$ belongs to $\mathbf{N}$ (and $\mathbf{L}$) if and only if the rational function $r(z)$ has no pole on $\mathfrak{L}$. Therefore $\mathbf{L}$ and $\mathbf{N}$ contain all the rational forms $r(A)$ commonly. Hence they must be coincident with each other. q.e.d.

§3. Spectrums and semi-spectrums of bounded linear operators.

We shall now apply our Theorem 1 and 2 to an analysis of the spectrums of general bounded linear operators on Banach spaces.

Let $\mathbf{B}$ be a Banach space and A be a bounded linear operator on it. The *spectrum* of A is the set of all complex numbers z so that $A - zI$ fail to be inversible. There exists the largest Banach algebra $\mathbf{R}(A)$ of bounded linear operators generated by A . $\mathbf{R}(A)$ is the uniform closure of the set of all rational forms $r(A)$. In order to avoid any mis-understand, a spectrum $\mathfrak{S}(\mathbf{N})$ of A corresponding to a Banach algebra $\mathbf{N}$ of bounded linear operators generated by A , is called a *semi-spectrum* of A . Then $\mathfrak{S}(\mathbf{N})$ is the set of all complex numbers z so that $(A - zI)^{-1}$ do not appear in $\mathbf{N}$. Clearly the semi-spectrum of A corresponding to $\mathbf{R}(A)$ is coincident with the usual spectrum $\mathfrak{S}$ of A . Then using Theorem 1 we have

Theorem 3. *Let A be a bounded linear operator on a Banach space B . In order that a complex number z belong to the spectrum of A , it is necessary and sufficient that $|r(A)| \geq |r(z)|$ hold for every rational form $r(A)$. The spectrum $\mathfrak{S}$ of A is a bounded closed set, and every rational form $r(A)$ satisfies*

$$\lim_{n \rightarrow \infty} |r^n(A)|^{1/n} = \sup_{z \in \mathfrak{S}} |r(z)|.$$

Every Banach algebra of bounded operators generated by A is a Banach sub-algebra of $\mathbf{R}(A)$. Then we can determine the type of all such Banach algebras using our Theorem 2.

Theorem 4. *Let $\mathfrak{S}$ be the spectrum of a bounded linear operator A . If $\mathbf{N}$ is a Banach algebra of bounded operators generated by A , then its corresponding semi-spectrum $\mathfrak{S}(\mathbf{N})$ is a bounded closed set containing $\mathfrak{S}$ such that $\mathfrak{S}(\mathbf{N}) - \mathfrak{S}$ is open.*

Conversely if $\mathfrak{L}$ is a bounded closed set which contains $\mathfrak{S}$ such that $\mathfrak{L} - \mathfrak{S}$ is open, then there exists a uniquely determined Banach algebra $\mathbf{N}$ generated by A whose corresponding semi-spectrum coincides with $\mathfrak{L}$.

§4. Some remarks for Theorem 2.

Given a Banach algebra $\mathbf{M}$ generated by an element A . Let $\mathcal{S}$ denote the spectrum of A , and $\mathcal{R}$ denote the resolvent set $\mathcal{R}_0 - \mathcal{S}$. (Conveniently we add the infinity point of the Riemann sphere to the resolvent set $\mathcal{R}$). $\mathcal{R}$ is the sum of mutually disjoint connected components $\mathcal{R} = \mathcal{R}_\infty \cup (\bigcup_{i=1}^{\infty} \mathcal{R}_i)$, where $\mathcal{R}_\infty$ is the connected component of $\mathcal{R}$ which contains the infinity.

Now if $\mathcal{I}$ is a bounded closed set containing $\mathcal{S}$ such that $\mathcal{I} - \mathcal{S}$ is open, then $\mathcal{I} - \mathcal{S}$ is a component of $\mathcal{R}$ which does not contain $\mathcal{R}_\infty$. That is, $\mathcal{I} - \mathcal{S}$ is a sum $\mathcal{I} - \mathcal{S} = \bigcup \mathcal{R}_{n_i}$, where $\mathcal{R}_{n_i}$ are connected components of $\mathcal{R}$ which do not contain the infinity. Since every Banach sub-algebra $\mathbf{N}$ of $\mathbf{M}$ is determined by the corresponding semi-spectrum $\mathcal{S}(\mathbf{N})$, $\mathbf{N}$ is completely determined by the system of connected components $\mathcal{R}_{n_i}$ which are contained in $\mathcal{S}(\mathbf{N})$. Therefore

Theorem 5. *Let $\mathbf{M}$ be a normalized Banach algebra generated by an element A . If the resolvent set $\mathcal{R}$ of A is decomposed into n pieces of connected components, then there exists exactly 2^{n-1} pieces of mutually different Banach sub-algebras of $\mathbf{M}$ generated by A .*

Let $\mathbf{P}$ denote the uniform closure of all rational integral forms $p(A)$. Then $\mathbf{P}$ is the smallest Banach sub-algebra of $\mathbf{M}$ generated by A . The complement of the corresponding semi-spectrum $\mathcal{S}(\mathbf{P})$ coincides with the connected component of the resolvent set $\mathcal{R}$ of A which contains the infinity. Especially when the resolvent set of A is connected, then $\mathbf{M}$ coincides with $\mathbf{P}$, and every rational form $r(A)$ is uniformly approximative by a sequence of rational integral forms $\{p_n(A)\}$.

REFERENCES

- [1] I. GELFUND, Normierte Ringe, Rec. Mat., 9 (1941), 3 - 24.
- [2] HALMOS, Introduction to Hilbert space, (1951).
- [3] F. RIESZ and B. v. SZ. NAGY, Leçons d'analyse Fonctionnelle, (1952).

DEPARTMENT OF MATHEMATICS,
OKAYAMA UNIVERSITY

(Received November 8, 1954)