Mathematical Journal of Okayama University

Volume 37, *Issue* 1 1995 *Article* 7

JANUARY 1995

Prime Ideals in Polynomial Rings over Tame Orders and Hereditary PI-Rings

Hidetoshi Marubayashi* Yang Lee†

Jae Keol Park‡

*Naruto University

†Busan National University

‡Busan National University

Copyright ©1995 by the authors. *Mathematical Journal of Okayama University* is produced by The Berkeley Electronic Press (bepress). http://escholarship.lib.okayama-u.ac.jp/mjou

Math. J. Okayama Univ. 37(1995), 65–74

PRIME IDEALS IN POLYNOMIAL RINGS OVER TAME ORDERS AND HEREDITARY PI-RINGS

HIDETOSHI MARUBAYASHI, YANG LEE and JAE KEOL PARK

For a given hereditary prime PI-ring Λ and a central polynomial $f(x)$ in $\Lambda[x]$ with $f(x)\Lambda[x]$ a prime ideal in $\Lambda[x]$, it was proved in [8] and [13] that the prime factor ring $\Lambda[x]/f(x)\Lambda[x]$ is hereditary if and only if $f(x)$ is not contained in the square of any maximal ideal of $\Lambda[x]$, which is a generalization of a main result in [7].

From this result together with the fact that hereditary prime PI-rings can be tame orders by [17], first in Section 1, we investigate, in Theorem A, a condition of prime factor rings of the polynomial ring $\Lambda[x]$ to be tame orders whenever Λ is a tame order. As applications, when D is a Krull domain we give a criterion for a certain class of prime factor rings of $D[x]$ to be a Krull domain.

Furthermore, in Section 2, when Λ is a hereditary PI-ring, in Theorem B, we give a criterion for prime factor rings of $\Lambda[x]$ to be hereditary, which is a nontrivial extension of the main result in [13, Theorem] and [14, Proposition 1], thereby we can provide an answer to a question of Armendariz [1], i.e., "a characterization of prime ideals P of the polynomial ring $\Lambda[x]$ over a hereditary PI-ring Λ such that $\Lambda[x]/P$ is hereditary", which was raised from a result in [2]: every prime factor ring of a hereditary PI-ring is hereditary.

1. Prime ideals in polynomial rings over tame orders. Let Λ be an order in a simple Artinian ring Q. For a Λ-ideal A in Q, we will use the notation;

$$\begin{aligned} O_\ell(A) &= \{q \in Q \mid qA \subseteq A\}, \\ O_r(A) &= \{q \in Q \mid Aq \subseteq A\}, \\ (\Lambda : A)_\ell &= \{q \in Q \mid qA \subseteq \Lambda\}, \\ (\Lambda : A)_r &= \{q \in Q \mid Aq \subseteq \Lambda\}, \\ A_v = (\Lambda : (\Lambda : A)_\ell)_r \quad &\text{and} \quad {}_vA = (\Lambda : (\Lambda : A)_r)_\ell. \end{aligned}$$

Clearly A_v and ${}_vA$ are again Λ-ideals containing A. We say that A is a *v-ideal* if ${}_vA = A = A_v$. A v-ideal A is *v-invertible* if $(A(\Lambda : A)_r)_v =$

Produced by The Berkeley Electronic Press, 1995

$\Lambda = {}_v((\Lambda : A)_\ell A)$. As it is well known, a Λ-ideal A is projective as left and right Λ-modules if and only if $A(\Lambda : A)_\ell = O_\ell(A)$ and $(\Lambda : A)_r A = O_r(A)$.

The following is a Krull type generalization of projectivities:

(K): ${}_v(A(\Lambda : A)_\ell) = O_\ell(A)$ *for any* Λ*-ideal* A *such that* $A = {}_vA$, *that is,* A *is reflexive as a left* Λ*-module, and* $((\Lambda : A)_r A)_v = O_r(A)$ *for any* Λ*-ideal* A *such that* $A = A_v$.

An order Λ is said to be *v-hereditary* (simply, a VH-*order*) if Λ satisfies the condition (K).

For a right Λ-module X, we denote by $E_\Lambda(X)$ (simply, $E(X)$) the injective hull of X. Let $\mathcal{C} = \mathcal{C}(\Lambda)$ be the right Gabriel topology corresponding to the torsion theory cogenerated by $E(Q/\Lambda)$. Then $\mathcal{C} = \{C$: a right ideal of $\Lambda \mid (\Lambda : r^{-1}C)_\ell = \Lambda$ for any $r \in \Lambda\}$ by Proposition 5.5 of [18, p.147], where $r^{-1}C = \{\lambda \in \Lambda \mid r\lambda \in C\}$. Similarly, we can define the left Gabriel topology $\mathcal{C}'$ on Λ. Let I be any right Λ-ideal (or a right ideal of Λ). We put $\tilde{I} = \{q \in Q \mid qC \subseteq I$ for some $C \in \mathcal{C}\}$. I is said to be $\mathcal{C}$*-closed* if $I = \tilde{I}$. If I is a right ideal of Λ, then we note that $\tilde{I} = \{\lambda \in \Lambda \mid \lambda C \subseteq I$ for some $C \in \mathcal{C}\}$. Also left $\mathcal{C}'$-closed ideals can be defined similarly.

In [3], Chamarie has considered the following condition to get the classical localization Λ_A of Λ at a v-ideal A in case Λ is a maximal order.

(C): Λ *satisfies the maximum condition on right* $\mathcal{C}$*-closed ideals of* Λ *and left* $\mathcal{C}'$*-closed ideals of* Λ.

Following [9], Λ is called a VHC-*order* if it is a VH-order and satisfies the condition (C).

Through this section, *we assume from now on that* D *is a Krull domain which is not a field*; K is the field of fractions of D. Let Λ be a tame D-order in a central simple K-algebra Q with finite dimension over K (see [4]). We denote by $D(\Lambda)$ the set of all v-invertible ideals in Q and by $P(\Lambda)$ the subset of $D(\Lambda)$ consisting of all principal Λ-ideals.

Now we summarize some properties of tame orders which will be necessary.

(i) Λ is a VHC-order with enough v-invertible ideals in the sense of [10] (see [9, Proposition 3.1] and [12, Proposition 1.1]).

(ii) $D(\Lambda)$ is a free abelian group generated by maximal v-invertible ideals (see [9, Theorem 1.13]).

(iii) $\Lambda[x]$ is a tame $D[x]$-order (see [4, Theorem 1.11]).

(iv) $D(\Lambda)/P(\Lambda)$ is naturally group isomorphic to $D(\Lambda[x])/P(\Lambda[x])$ (see [10, Theorem 2.19]).

Marubayashi et al.: Prime Ideals in Polynomial Rings over Tame Orders and Hereditary

(v) The set of all maximal v-invertible ideals of $\Lambda[x]$ is $\{B[x], A \mid B$ is a maximal v-invertible ideal of Λ and $A = A' \cap \Lambda[x]$ for some maximal ideal A' of $Q[x]\}$ (see [9, Theorem 3.9]).

Also throughout this section P is always a non-zero prime ideal of $\Lambda[x]$ with $P \cap \Lambda = 0$. Then in this case we have the following:

(vi) $PQ[x] \cap \Lambda[x] = P$ and $PQ[x]$ is also a prime ideal.

(vii) P is a v-ideal.

(viii) P is a minimal non-zero prime ideal of $\Lambda[x]$.

The proof of (vi) is straightforward. Since $Q[x]$ is a principal ideal ring [16], any ideal of $Q[x]$ is a v-ideal. So (vii) follows from [9, Lemma 2.3] and (vi). (viii) follows from (vi), because any non-zero prime ideal of $Q[x]$ is maximal.

We begin with local case.

Lemma 1. *Assume that D is a discrete rank one valuation domain and that Λ is hereditary. Let P be a non-zero prime ideal of $\Lambda[x]$ with $P \cap \Lambda = 0$. Then there exists $f(x) \in D[x]$ such that $P = f(x)\Lambda[x]$.*

Proof. Let $\wp = P \cap D[x]$, a minimal non-zero prime ideal of the Krull domain $D[x]$. Then, by Nagata's theorem, there exists $f(x) \in D[x]$ such that $\wp = f(x)D[x]$ (see (iv)). Now it is clear that $f(x)K[x]$ is a prime ideal with $f(x)K[x] \cap D[x] = \wp$, and K is embedded in $K[x]/f(x)K[x]$, which is a field. So it follows from [15, Theorem 7.6] that $Q[x]/f(x)Q[x]$ ($\cong Q \otimes_K K[x]/f(x)K[x]$) is a simple Artinian ring. Hence $f(x)Q[x]$ is a maximal ideal contained in $P' = PQ[x]$ and so $P' = f(x)Q[x]$. Since $f(x)\Lambda[x]$ is invertible and $f(x)\Lambda[x] \subseteq P$, we have

$$f(x)\Lambda[x] = P^n \cdot P_1^{n_1} \cdots P_k^{n_k} \cdot J^m[x],$$

where P_i is a prime ideal such that $P_i = P_i' \cap \Lambda[x]$ for some maximal ideal P_i' of $Q[x]$, and $J = J(\Lambda)$, the Jacobson radical of Λ. Note that J is the unique maximal invertible ideal of Λ, because D is a discrete rank one valuation domain. Then it follows that

$$P' = P'^n \cdot P_1'^{n_1} \cdots P_k'^{n_k},$$

and so we have $k = 0$ and $n = 1$. Now to prove that $m = 0$, assume to the contrary that $m \geq 1$, then $f(x) \in J[x] \cap D[x] = J(D)[x]$. In this case $J(D)$ is the unique maximal ideal of D. Thus we have $f(x)D[x] \subseteq J(D)[x]$, a minimal non-zero prime ideal of $D[x]$. So $f(x)D[x] = J(D)[x]$ and this

Produced by The Berkeley Electronic Press, 1995

implies that $f(x)K[x] = K[x]$. This is a contradiction and so $m = 0$, that is, $P = f(x)\Lambda[x]$. This completes the proof.

Let p be a minimal non-zero prime ideal of D. Then Λ_p is hereditary over the discrete rank one valuation domain D_p. So by Lemma 1, there exists $f_p(x) \in D_p[x]$ such that $\wp_p = f_p(x)D_p[x]$ and $P_p = f_p(x)\Lambda_p[x]$.

Lemma 2. *Let M be a prime ideal of $\Lambda[x]$ such that $M \cap (D \setminus p) = \emptyset$ for some minimal non-zero prime ideal p of D. Then* $\mathrm{rank}(M) \leq 2$ *and* $\mathrm{rank}(M \cap \Lambda) \leq 1$.

Proof. $\mathrm{rank}(M) \leq 2$ follows from the fact that the classical Krull dimension of $\Lambda_p[x]$ equals to 2. To prove that $\mathrm{rank}(M \cap \Lambda) \leq 1$, let $\mathcal{M} = M \cap \Lambda$ and suppose that $\mathcal{M}$ is not a minimal non-zero prime ideal. Then $M = \mathcal{M}[x]$, because

$$0 \subsetneq \mathcal{M}_1[x] \subsetneq \mathcal{M}[x] \subseteq M,$$

where $\mathcal{M}_1$ is a non-zero prime ideal strictly contained in $\mathcal{M}$. However, since $\mathcal{M} \not\subseteq p$, we have $M_p = \mathcal{M}_p[x] = \Lambda_p[x]$ and so $M \cap (D \setminus p) \neq \emptyset$, a contradiction. Hence $\mathrm{rank}(\mathcal{M}) \leq 1$.

For a prime ideal M of $\Lambda[x]$ we define $M^{(2)}$ as follows:

If $M \cap \Lambda = 0$, then $M^{(2)} = M^2$. If $M \cap \Lambda \neq 0$, then $m = M \cap D$ is a non-zero prime ideal and

$$M^{(2)} = \{\lambda(x) \in \Lambda[x] \mid \lambda(x)c \in M^2 \text{ for some } c \in D \setminus m\}.$$

Note that $M^{(2)} = (M_m)^2 \cap \Lambda[x]$.

Lemma 3. *The following are equivalent:*

(1) *$f_p(x) \notin (M_p)^2$ for any prime ideal M of $\Lambda[x]$ with $M \cap (D \setminus p) = \emptyset$, where p is a minimal non-zero prime ideal of D.*

(2) *$P \not\subseteq M^{(2)}$ for any prime ideal M of $\Lambda[x]$ with* $\mathrm{rank}(M \cap \Lambda) \leq 1$.

Proof. (1) implies (2). Assume to the contrary that $P \subseteq M^{(2)}$ for some prime ideal M of $\Lambda[x]$ with $\mathrm{rank}(M \cap \Lambda) \leq 1$.

Case I. If $\mathcal{M} = M \cap \Lambda = 0$, then $M = MQ[x] \cap \Lambda[x]$ and so M is a prime v-ideal which is v-invertible by [9, Lemma 3.6]. Furthermore, M is a minimal non-zero prime ideal of $\Lambda[x]$ by [11, Lemma 1.8]. If $M = P$, then, since $M \cap (D \setminus p) = \emptyset$ for any minimal non-zero prime ideal p of D,

we have $P_p = f_p(x)\Lambda_p[x] = M_p \subseteq (M^{(2)})_p = (M_p)^2$, a contradiction. If $M \neq P$, then, of course, we have $P \not\subseteq M^{(2)}$, a contradiction.

Case II. If $\mathcal{M} = M \cap \Lambda$ is non-zero, then $\mathcal{M}$ is a minimal non-zero prime ideal of Λ and $m = M \cap D$ is also a minimal non-zero prime ideal of D with $M \cap (D \setminus m) = \emptyset$. So $f_m(x)\Lambda_m[x] = P_m \subseteq (M^{(2)})_m = (M_m)^2$, a contradiction.

Therefore by Cases I and II, $P \not\subseteq M^{(2)}$ for any prime ideal M of $\Lambda[x]$ with $\mathrm{rank}(M \cap \Lambda) \leq 1$.

(2) implies (1). Suppose that $f_p(x) \in (M_p)^2$ for some prime ideal M of $\Lambda[x]$ with $M \cap (D \setminus p) = \emptyset$, where p is a minimal non-zero prime ideal of D. By Lemma 2, $\mathrm{rank}(M \cap \Lambda) \leq 1$. Let $m = M \cap D$. Then either $m = 0$ or m is a minimal non-zero prime ideal. If $m = 0$, then M is a minimal non-zero prime ideal. First of all, if $P = M$, then $f_p(x)\Lambda_p[x] = P_p \subseteq (P_p)^2$, a contradiction. If $P \neq M$, then $P_p \neq M_p$, because $P = P_p \cap \Lambda[x]$ and $M = M_p \cap \Lambda[x]$. Hence $f_p(x) \notin M_p$ and so $f_p(x) \notin (M_p)^2$, a contradiction. Now if m is a minimal non-zero prime ideal, then $M \cap (D \setminus p) = \emptyset$ implies that $m \subseteq p$ and thus $m = p$. Since $P_p = f_p(x)\Lambda_p[x] \subseteq (M_p)^2$, we have $P = P_p \cap \Lambda[x] \subseteq (M_p)^2 \cap \Lambda[x] = M^{(2)}$, a contradiction.

Therefore $f_p(x) \notin (M_p)^2$ for any prime ideal M of $\Lambda[x]$ with $M \cap (D \setminus p) = \emptyset$, where p is a minimal non-zero prime ideal of D.

Lemma 4. *Let p be a minimal non-zero prime ideal of D. Then $(\Lambda[x]/P)_p = \Lambda_p[x]/P_p$ is hereditary if and only if $f_p(x) \notin (M_p)^2$ for any prime ideal M of $\Lambda[x]$ with $M \cap (D \setminus p) = \emptyset$.*

Proof. There is a one-to-one correspondence between prime ideals M of $\Lambda[x]$ with $M \cap (D \setminus p) = \emptyset$ and prime ideals M' of $\Lambda_p[x]$ corresponding M in $\Lambda[x]$ to $M' = M_p$ and M' in $\Lambda_p[x]$ to $M' \cap \Lambda[x]$. Hence the lemma follows from [13, Theorem].

Lemma 5. *$\Lambda[x]/P = \bigcap_p(\Lambda_p[x]/P_p)$ and $D[x]/\wp = \bigcap_p(D_p[x]/\wp_p)$, where p runs through all minimal non-zero prime ideals of D and $\wp = P \cap D[x]$.*

Proof. Since $\Lambda = \bigcap \Lambda_p$ and $D = \bigcap D_p$, it follows that $\Lambda[x] = \bigcap \Lambda_p[x]$ and $D[x] = \bigcap D_p[x]$. To prove that $P = \bigcap P_p$, let $z \in \bigcap P_p$. Then for any p there exists $c_p \in D \setminus p$ with $zc_p \in P$. Set $\mathcal{A} = \sum c_p\Lambda$. Then $\mathcal{A}_v = \Lambda$, because, on the contrary, assume that $\mathcal{A}_v \subsetneq \Lambda$, then there exists a maximal v-ideal P_0 with $\mathcal{A}_v \subseteq P_0$. Then $p_0 = P_0 \cap D$ is a minimal non-zero prime

ideal of D and for this p_0, we have $c_{p_0} \in P_0 \cap D = p_0$, a contradiction. Now $z\mathcal{A}[x] \subseteq P$ implies that $z \in z\Lambda[x] = z\mathcal{A}_v[x] \subseteq (z\mathcal{A}[x])_v \subseteq P_v = P$. Hence $P = \bigcap P_p$ and similarly $\wp = \bigcap \wp_p$. Therefore $\Lambda[x]/P = \bigcap(\Lambda_p[x]/P_p)$ and $D[x]/\wp = \bigcap(D_p[x]/\wp_p)$.

Lemma 6. *Suppose that one of the conditions in Lemma 3 is satisfied. Then we have the following*:

(1) $D[x]/\wp$ *is a Krull domain and it is the center of* $\Lambda[x]/P$, *where* $\wp = P \cap D[x]$.

(2) $\Lambda[x]/P$ *is a tame* $D[x]/\wp$*-order.*

Proof. (1) Let p be a minimal non-zero prime ideal of D. Then Λ_p is hereditary with its center D_p and $P_p = f_p(x)\Lambda_p[x]$. So it follows from [14, Claim 4, p.1485] and Lemma 4 that $\Lambda_p[x]/P_p$ is hereditary with $Z(\Lambda_p[x]/P_p) = D_p[x]/\wp_p$, where $Z(-)$ denotes the center of a ring. In particular, $D_p[x]/\wp_p$ is a Dedekind domain. Thus $D[x]/\wp$ is a Krull domain by Lemma 5. Furthermore, we have

$$Z(\Lambda[x]/P) = \bigcap_p Z(\Lambda_p[x]/P_p) = \bigcap_p D_p[x]/\wp_p = D[x]/\wp$$

by Lemma 5.

(2) For any minimal non-zero prime ideal p, $\Lambda_p[x]/P_p$ is hereditary with $Z(\Lambda_p[x]/P_p) = D_p[x]/\wp_p$. So it follows that

$$\Lambda_p[x]/P_p = \bigcap(\Lambda_p[x]/P_p)_{q'(p)},$$

where $q'(p)$ ranges over all prime ideals of $D_p[x]/\wp_p$. There is a one-to-one correspondence between prime ideals $q'(p)$ of $D_p[x]/\wp_p$ and minimal non-zero prime ideals $q(p)$ of $D[x]/\wp$ with $q(p) \cap (D \setminus p) = \emptyset$ which is given by;

$$q'(p) \longrightarrow q(p) = q'(p) \cap (D[x]/\wp) \text{ and } q(p) \longrightarrow q(p)(D_p[x]/\wp_p).$$

Furthermore $(\Lambda[x]/P)_{q(p)} = (\Lambda_p[x]/P_p)_{q'(p)} \supseteq (D[x]/\wp)_{q(p)}$. Hence we have that

$$\Lambda[x]/P = \bigcap_p \bigcap_{q(p)} (\Lambda[x]/P)_{q(p)}$$

and $(\Lambda[x]/P)_{q(p)}$ is hereditary. To prove that $\Lambda[x]/P$ has the finite character property, let $\overline{\lambda(x)}$ be any regular element in $\overline{\Lambda[x]} = \Lambda[x]/P$. Since $\overline{\lambda(x)}\cdot\overline{\Lambda[x]}$ is an essential right ideal of $\overline{\Lambda[x]}$, we have $\overline{\lambda(x)}\cdot\overline{\Lambda[x]} \cap Z(\overline{\Lambda[x]}) \neq 0$.

Thus there exists $d(x) \in D[x]$ with $\overline{d(x)} \in \overline{\lambda(x)} \cdot \overline{\Lambda[x]}$. Since $\overline{D[x]} = D[x]/\wp$ is a Krull domain, there are only finite number of minimal non-zero prime ideals of $\overline{D[x]}$, say, $q_1, \ldots, q_n$, (each $q_i = q(p)$ for some $q(p)$) such that $\overline{d(x)} \cdot \overline{D[x]}_{q_i} \subsetneq \overline{D[x]}_{q_i}$. Hence $\Lambda[x]/P$ satisfies the finite character property, because $\overline{\Lambda[x]}_{q(p)} \supseteq \overline{D[x]}_{q(p)}$ for each $q(p)$. Hence $\overline{\Lambda[x]}$ is a tame $\overline{D[x]}$-order by [5, Lemma 1.1] and [9, Proposition 3.1].

The following lemma is implicitly known. However, we could not find the place in which the proof of the lemma is given. So we give a complete proof for our convenience. We denote by *cl.K.dim* Λ the classical Krull dimension of Λ.

Lemma 7. *Let Λ be a Noetherian tame D-order with* cl.K.dim $\Lambda \leq 1$. *Then Λ is hereditary.*

Proof. If cl.K.dim $\Lambda = 0$, then Λ is a simple Artinian ring and so it is hereditary. If cl.K.dim $\Lambda = 1$, then any prime ideal M of Λ is a prime v-ideal by [11, Lemma 1.8]. Furthermore, $\Lambda = \bigcap \Lambda_p$, where p runs through all minimal non-zero prime ideals of D and Λ_p is hereditary. So we have

$$1 \in \Lambda_p \cap O_r(M_p) = \Lambda_p \cap (\Lambda_p : M_p)_r M_p = (\Lambda \cap (\Lambda : M)_r M)\Lambda_p,$$

where $O_r(M_p) = \{q \in Q \mid M_p q \subseteq M_p\}$. Thus

$$(\Lambda \cap (\Lambda : M)_r M)_v = \bigcap_p (\Lambda \cap (\Lambda : M)_r M)\Lambda_p = \Lambda.$$

This implies that $M \subsetneq \Lambda \cap (\Lambda : M)_r M$ and so $\Lambda \cap (\Lambda : M)_r M = \Lambda$, i.e., $1 \in (\Lambda : M)_r M$. Hence it follows that M is left projective and similarly right projective. Then the lemma follows from the same method as in [6, Proposition 1.3].

Now from all lemmas prepared, we have one of our main results of this note as follows;

Theorem A. *Let D be a Krull domain with the field of quotients K and let Λ be a tame D-order in a central simple K-algebra Q with finite dimension over K. Let P be a prime ideal of $\Lambda[x]$ with $P \cap \Lambda = 0$. Then $\Lambda[x]/P$ is a tame order if and only if $P \not\subseteq M^{(2)}$ for any prime ideal M of $\Lambda[x]$ with* rank$(M \cap \Lambda) \leq 1$. *Furthermore, under these conditions, the center $Z(\Lambda[x]/P)$ of $\Lambda[x]/P$ is $D[x]/\wp$, where $\wp = P \cap D[x]$.*

Let M be a prime ideal of $\Lambda[x]$ with $\mathrm{rank}(M \cap \Lambda) \leq 1$ and $P \not\subseteq M$. Then it is easily checked that $P \not\subseteq M^{(2)}$. Thus we have the following:

Remark. $\Lambda[x]/P$ is a tame order if and only if $P \not\subseteq M^{(2)}$ for any prime ideal M of $\Lambda[x]$ satisfying both $\mathrm{rank}(M \cap \Lambda) \leq 1$ and $P \subsetneq M$.

Let $f(x)$ be any polynomial in $D[x]$ with $\deg f(x) \geq 1$. Then $f(x)\Lambda[x] \cap \Lambda = 0$. So we have the following corollary which extends results in [7], [8] and [13].

Corollary 1. *Let $f(x)$ be any polynomial in $D[x]$ such that $P = f(x)\Lambda[x]$ is a prime ideal of $\Lambda[x]$ and deg $f(x) \geq 1$. Then $\Lambda[x]/P$ is a tame order if and only if $f(x) \notin M^{(2)}$ for any prime ideal M of $\Lambda[x]$ with* $\mathrm{rank}(M \cap \Lambda) \leq 1$.

Noting that Krull domains are tame orders, we have following fact immediately, which is Krull domains version of a result in [7].

Corollary 2. *For a Krull domain D, let $f(x)$ be any polynomial in $D[x]$ such that $P = f(x)D[x]$ is a prime ideal and deg $f(x) \geq 1$. Then $D[x]/P$ is a Krull domain if and only if $f(x) \notin M^{(2)}$ for any prime ideal M of $D[x]$ with* $\mathrm{rank}(M \cap D) \leq 1$.

2. Prime ideals in polynomial rings over hereditary PI-rings. As we mentioned, in this section we consider hereditary prime factor rings of the polynomial ring $\Lambda[x]$ over a hereditary PI-ring Λ.

Lemma 8. *Let Λ be a prime hereditary PI-ring and let M be a prime ideal of $\Lambda[x]$ with $\mathcal{M} = M \cap \Lambda \neq 0$. Then* $\mathrm{rank}(\mathcal{M}) = 1$ *and* $M^{(2)} = M^2$.

Proof. It is clear that $\mathrm{rank}(\mathcal{M}) = 1$ and $\mathcal{M}[x] \subseteq M$. First assume that $M = \mathcal{M}[x]$ and let $\lambda(x) \in M^{(2)}$. Then there exists an element $c \in D$, but $c \notin M$ such that $\lambda(x)c \in M^2$ and so $\lambda(x) \in M$. Since $\mathcal{M}$ is a maximal ideal, we have $\mathcal{M} + c\Lambda = \Lambda$ and $\Lambda[x] = \mathcal{M}[x] + c\Lambda[x] = M + c\Lambda[x]$. Hence

$$\lambda(x) \in \lambda(x)\Lambda[x] = \lambda(x)(M + c\Lambda[x]) \subseteq M^2.$$

Next assume that $\mathcal{M}[x] \subsetneq M$, then M is a maximal ideal. So we have $M + c\Lambda[x] = \Lambda[x]$ and hence $\lambda(x) \in M^2$.

By Theorem A and Lemma 8, we answer to a question of Armendariz [1], which is a characterization of hereditary prime factor rings of polynomial rings over a hereditary PI-ring, in the following:

Theorem B. *Let Λ be a hereditary PI-ring and let P be a prime ideal of $\Lambda[x]$. Let $P_0 = P \cap \Lambda$, a prime ideal of Λ. Then we have the following*:

(1) *If $P = P_0[x]$, then $\Lambda[x]/P$ is hereditary if and only if P_0 is a maximal ideal of Λ.*

(2) *If $P_0[x] \subsetneq P$, then $\Lambda[x]/P$ is hereditary if and only if $P \not\subseteq M^2 + P_0[x]$ for any prime ideal M of $\Lambda[x]$ with $P \subsetneq M$.*

Proof. (1) If $P = P_0[x]$, then $\Lambda[x]/P \cong (\Lambda/P_0)[x]$ and Λ/P_0 is a prime hereditary by [2, Theorem]. Hence $(\Lambda/P_0)[x]$ is hereditary if and only if Λ/P_0 is a simple Artinian ring, that is, P_0 is a maximal ideal.

(2) Set $\bar{\Lambda} = \Lambda/P_0$ and consider the natural mapping f from $\Lambda[x]$ to $\bar{\Lambda}[x]$. We just write $f(P)$ by $\bar{P}$. Then $\bar{P}$ is a non-zero prime ideal with $\bar{P} \cap \bar{\Lambda} = \bar{0}$. Hence the result follows from the remark to Theorem A and Lemma 8.

Acknowledgments. This paper was prepared while the first author was visiting Busan National University as a Visiting Professor. He would like to express his appreciation to members of the Mathematics Department of Busan National University for their warm hospitality. The third author was supported in part by KOSEF, RCAA and the Basic Science Research Institute Program, Ministry of Education, Project No. BSRI-95-1402.

The authors are very much indebted to the referee for many useful suggestions.

REFERENCES

[1] E. P. ARMENDARIZ: Private communication, 1990.
[2] E. P. ARMENDARIZ and C. R. HAJARNAVIS: On prime ideals in hereditary PI-rings, J. Algebra **116** (1988), 502–505.
[3] M. CHAMARIE: Anneaux de Krull non commutatif, J. Algebra **72** (1981), 210–222.
[4] R. M. FOSSUM: Maximal orders over Krull domains, J. Algebra **10** (1968), 321–332.
[5] H. FUJITA: Idealizers in generalized Krull orders, Comm. Algebra **11** (1983), 1369–1393.
[6] C. R. HAJARNAVIS and T. H. LENAGAN: Localization in Asano orders, J. Algebra **21** (1972), 441–449.
[7] J. A. HILLMAN: Polynomials determining Dedekind domains, Bull. Australian Math. Soc. **29** (1984), 167–175.

[8] Y. HIRANO, J. K. PARK and K. W. ROGGENKAMP: Global dimension of factor rings, Bull. Australian Math. Soc. **49** (1994), 399–411.
[9] H. MARUBAYASHI: A Krull type generalization of HNP rings with enough invertible ideals, Comm. Algebra **11** (1983), 469–499.
[10] H. MARUBAYASHI: A skew polynomial ring over a v-HC order with enough v-invertible ideals, Comm. Algebra **12** (1984), 1567–1593.
[11] H. MARUBAYASHI: Remarks on VHC-orders in a simple Artinian ring, J. Pure and Applied Algebra **31** (1984), 109–118.
[12] H. MARUBAYASHI: On bounded v-HC orders in a simple Artinian ring, Comm. Algebra **14** (1986), 1229–1251.
[13] H. MARUBAYASHI, Y. LEE and J. K. PARK: Polynomials determining hereditary prime PI-rings, Comm. Algebra **20** (1992), 2503–2511.
[14] J. K. PARK and K. W. ROGGENKAMP: A note on hereditary rings, Comm. Algebra **17** (1989), 1477–1493.
[15] I. REINER: Maximal Orders, Academic Press, New York 1975.
[16] J. C. ROBSON: Pri-rings and ipri-rings, Quart. J. Math. Oxford **18** (1967), 125–145.
[17] J. C. ROBSON and L. W. SMALL: Hereditary prime P.I. rings are classical hereditary orders, J. London Math. Soc. **8** (1974), 499–503.
[18] B. STENSTRÖM: Rings of Quotients, Grundlehren Math. Wiss. 217, Springer-Verlag, Berlin 1975.

H. MARUBAYASHI
DEPARTMENT OF MATHEMATICS
NARUTO UNIVERSITY OF EDUCATION
TAKASHIMA, NARUTO-SHI 772, JAPAN

Y. LEE AND J. K. PARK
DEPARTMENT OF MATHEMATICS
BUSAN NATIONAL UNIVERSITY
BUSAN 609-735, SOUTH KOREA

(*Received October 16, 1995*)