

Mathematical Journal of Okayama University

Volume 28, Issue 1

1986

Article 6

JANUARY 1986

Cozzens domains are hereditary

Carl Faith*

*State University

Copyright ©1986 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

COZZENS DOMAINS ARE HEREDITARY

CARL FAITH

A ring R is said to be a *right V-ring* (after Villamayor) provided that it satisfies the equivalent conditions :

- (V.1) $\text{rad } M = 0$ for all M in $\text{mod-}R$.
- (V.2) Each right ideal is the intersection of maximal right ideals.
- (V.3) Each simple right R -module is injective.

For proof consult [3], p. 356 or [6].

Condition (V.3) was introduced by Kaplansky who showed that (V.3) in commutative R is equivalent to R von Neumann regular.

A *generalized right Cozzens domain* is a right Noetherian right V -domain R for which every indecomposable injective right R -module is either simple or else embeds in the right quotient field of R .

A module M of $\text{mod-}R$ is a *B-object* (after Bass ; see 18.3(b) of [5]) provided that every submodule $S \neq M$ is contained in a maximal submodule. A ring R is a *right B-ring* if every M in $\text{mod-}R$ is a *B-object* (see [5], pp. 330–331 and 155 for further background and related results).

In [1] Cozzens constructed V -domains not fields in providing a negative answer to Bass' question of whether or not a right B -ring with no infinite sets of orthogonal idempotents was necessarily right perfect, equivalently satisfied the d.c.c. on principal left ideals. (Bass proved that the converse held.)

The domain R constructed by Cozzens were right and left hereditary, and right and left Noetherian, in fact, they were principal right (left) ideal rings. Furthermore, R has a unique simple right R -module and a unique indecomposable injective right R -module, namely the right quotient field K of R .

We prove that any right Ore domain with these two properties is necessarily right hereditary. More generally, we have :

Theorem. *Every generalized right Cozzens domain is hereditary.*

Proof. The proof requires a result of [4] which states that a right Noetherian right V -ring is right hereditary provided that each indecomposable injective right R -module E is restricted semisimple, that is, E/S is semisimple, whence injective, for each submodule $S \neq 0$. Since K is the only non-simple indecomposable injective, we check K/M , for a submodule $M \neq 0$.

Now, by the Matlis' structure theory of injective modules over right Noetherian rings, the injective hul $E(K/M)$ is a direct sum of nonzero indecomposable injectives

$$E(K/M) = \oplus E_i$$

where $0 \neq E_i = E(U_i/M)$ for some uniform submodule U_i/M of K/M . Now, there are just 2-possibilities for E_i , namely simple, or $E_i \approx K$. But if $E_i \approx K$, then U_i/M embeds in K , hence there is an R -morphism $f: U_i \rightarrow K$ with $\ker f = M$. Since K is injective, then there exists a mapping $f^*: K \rightarrow K$ extending f . However, M is essential in K , so it follows that for any $k \in K$, there exists $r \neq 0$ in R with $kr = u \in U_i$. Then,

$$f^*(u) = f^*(k)r = 0$$

whence $f^*(k) = 0$, proving that $f^* = 0$. But then, this implies that $f = 0$, whence $U_i = M$, contrary to the choice of U_i/M .

Thus, E_i is simple for all i , so $E(K/M)$ is semisimple, and therefore so is K/M . This proves what we wanted.

Another proof can be given using the results of [6], and the calculations above.

We remark that there exist generalized Cozzens domains with more than one isomorphism class of simple injectives. (See [2] and [7].)

REFERENCES

- [1] J. H. COZZENS : Homological properties of the ring of differential polynomials, Bull. Amer. Math. Soc. **76** (1970), 75–79.
- [2] J. H. COZZENS : Twisted group rings and a problem of Faith, Bull. Austral. Math. Soc. **19** (1973), 11–19.
- [3] C. FAITH : Algebra I: Rings, Modules, and Categories, Springer-Verlag, Berlin-Heidelberg-New York, 1973, corrected reprint 1981.
- [4] C. FAITH : On hereditary rings and Boyle's conjecture, Arch. der Math. **27** (1976), 113–119.
- [5] C. FAITH : Algebra II: Ring Theory, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [6] G. O. MICHLER and O. E. VILLAMAYOR : On rings whose simple modules are injective, J. Algebra **25** (1973), 185–201.
- [7] B. L. OSOFSKY : On twisted polynomial rings, J. Algebra **18** (1971), 597–607.

DEPARTMENT OF MATHEMATICS
RUTGERS, THE STATE UNIVERSITY
NEW BRUNSWICK, NEW JERSEY 08540, U.S.A.

(Received March 15, 1985)