Pteridines Vol. 20, Special issue, pp. 36-41

First Synthesis of a Representative, Natural Pterin Glycoside: 2'-O-(α -D-Glucopyranosyl)biopterin

Tadashi Hanaya^{1§}, Hiroshi Yamamoto² and Wolfgang Pfleiderer³

¹ Department of Chemistry, Faculty of Science, Okayama University, Tsushima-naka, Okayama 700-8530, Japan

² School of Pharmacy, Shujitsu University, Nishigawara, Okayama 703-8516, Japan

³ Fachbereich Chemie, Universität Konstanz, Postfach 5560, D-78457 Konstanz, Germany

Abstract

Glycosylation of N^2 -(N,N-dimethylaminomethylene)-1'-O-(4-methoxybenzyl)-3-[2-(4-nitrophenyl)ethyl]biopterin (14) with the novel donor 4,6-di-O-acetyl-2,3-di-O-(4-methoxybenzyl)- α -D-glucopyranosyl bromide (19) in the presence of silver triflate and tetramethylurea predominantly afforded the corresponding α -D-glucopyranoside (20a), from which 2'-O-(α -D-glucopyranosyl)biopterin (1) was obtained by the successive removal of the protecting groups.

Key words: biopterin D-glucoside, pterin glycoside, glycosylation, protecting groups

Introduction

Some pterins having a hydroxyalkyl side-chain at C-6 have been found as glycosidic forms in certain prokaryotes, representative examples being 2'-O-(α -D-glucopyranosyl)biopterin (1) isolated from various kinds of cyanobacteria [1-6] and limipterin [2'-O-(2-acetamido-2deoxy-β-D-glucopyranosyl)biopterin] (2) isolated from a green sulfur photosynthetic bacterium [7]. Glycosides (3-5) of other pterins such as ciliapterin (L-threo-biopterin), neopterin and 6-hydroxymethylpterin have also been isolated from cyanobacteria, anaerobic photosynthetic bacteria and chemoautotrophic archaebacteria [8–10]. We have undertaken a synthetic exploration of various types of glycosides of biopterin and related pterins owing to a marked interest in their physiological functions and biological activities as well as the structural

Pteridines / Vol. 20 / Special issue

proof of those natural products [11-17]. We present here the first synthesis of 2'-*O*-(α -D-glucopyranosyl)biopterin (1), which remained unprepared since its first discovery in 1958.



⁸Correspondence : Dr. Tadashi Hanaya, Department of Chemistry, Faculty of Science, Okayama University, Tsushima-naka, Okayama 700-8530, Japan; F: +81-86-251-7853; Email: hanaya@cc.okayama-u.ac.jp

Results and Discussion

In our initial studies on glycosylation of a sidechain hydroxy group, we had to device suitable protecting and at the same time solubilizing groups for the pyrimidine ring and the side-chain hydroxy groups of the starting material biopterin (6). This was because pterin derivatives are little soluble, owing to the effectively stabilized intramolecular hydrogen bondings in the solid state [18], in such a nonpolar aprotic solvent as dichloromethane in which glycosylation reactions normally proceed smoothly. Thus, 6 was converted in a five-step procedure via intermediate 1',2'-di-O-acetyl- N^2 -(N,Ndimethylaminomethylene)-3-[2-(4-nitrophenyl)ethyl]biopterin into the sufficiently solubilized 1',2'-di-O-trimethylsilyl derivative (7) in 86% overall yield (Scheme 1) [12, 19]. Glycosylation of 7 with 2,3,4,6-tetra-O-benzoyl-α-Dglucopyranosyl bromide resulted in the formation of a mixture of 2'-O-(β -D-glucopyranosyl) 8a (41%), 1'-O-glycosyl isomer 8b (15%) and the 1',2'-di-O-glycosyl 8c (14%).

These results prompted us to pursue preparations of a suitably 1'-O-protected biopterin derivative in order to achieve complete 2'-O-glycosylation as shown in a retrosynthetic analysis for 9a outlined in Scheme 2. This compound was obviously not derived directly from biopterin but it appeared to be best prepared by the condensation of 2,5,6-triamino-4-hydroxypyrimidine with 3-Oprotected pentos-2-ulose 10, which would be derived from 3-O-protected 5-deoxy-L-arabinose 11. As these compounds were both unknown, we explored a synthetic route starting with D-xylose involving C4 inversion and C5 deoxygenation [15]. Moreover, a rational consideration of the available conditions to remove the protecting groups of the glycoside derived from 1'-O-protected biopterin 9a led us to employ *p*-methoxybenzyl (PMB) group for protection of 1'-hydroxy, N,N-dimethylaminomethylene for 2-amino, and 2-(4-nitrophenyl)ethyl (NPE) for N-3 of the ring.

A 9-step synthetic sequence for the 5-deoxy-L-*erythro*-pentos-2-ulose (10) from D-xylose via a known compound 12 [20] and 3-*O*-PMB-5-



Scheme 2

deoxy-L-arabinose (11) is shown in Scheme 3.

Condensation of 10 with 2,5,6-triamino-4hydroxypyrimidine sulfate in an aqueous sodium bicarbonate solution afforded nearly an 8:2 mixture of the L-biopterin derivative 9a and its 7-substituted isomer (L-primapterin) 9b. Successive treatment of the mixture with *N*,*N*dimethylformamide dimethyl acetal in DMF, with acetic anhydride in pyridine, and then with NPE alcohol (Mitsunobu reaction), gave the versatile L-biopterin derivative 13a (53% overall yield from 10) and L-primapterin derivative 13b (15%). Methanolysis of 13a in the presence of sodium methoxide quantitatively provided the 1'- *O*-PMB derivative 14, an ideal precursor for 2'-*O*-monoglycosylation [13].

An efficient glycosylation was exemplified by the condensation of 14 with tetra-*O*-benzoyl- α -D-glucopyranosyl bromide in the presence of silver triflate and tetramethylurea (TMU) in dichloromethane, affording 2'-*O*- β -Dglucopyranosyl derivative 15 as a sole product in 75% yield (Scheme 4). Deprotection of 15 efficiently afforded 2'-*O*-(β -D-glucopyranosyl) biopterin (16). Similarly, glycosylation of 14 with 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl bromide provided the 2'-*O*-glucopyranosyl derivative (17) in 82% yield, from which natural product limipterin (2) was



Scheme 3





obtained smoothly after deprotection [15].

The stereoselective formation of the β -glycoside (15) from 14 was mainly caused by participation of the 2-*O*-benzoyl group of the glycosyl donor through the formation of an acyloxonium ion intermediate [21]. Accordingly, in order to avoid such a neighboring group participation in synthesis of biopterin α -D-glucoside (1), we sought to introduce an ether substituent for protection of 2-OH of a glycosyl donor; thus PMB and acetyl groups were respectively chosen

for protection of 2,3-OH and 4,6-OH of the glycosyl moiety. A synthetic route is shown in Scheme 5 for such an appropriately protected α -D-glucopyranosyl bromide derivative (19) starting with penta-*O*-acetyl- β -D-glucopyranose (18) via methyl 2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside [22].

Then, glycosylation of 14 was found to give the best result when it was treated with 4.0 mol equiv. of the glycosyl bromide (19) in dichloromethane in the presence of silver triflate (2.0 mol equiv.)



Scheme 6

Pteridines / Vol. 20 / Special issue

and tetramethylurea (1.0 mol equiv.), affording an inseparable anomeric mixture (85:15) of the 2'-O-(α -D-glucopyranosyl)biopterin derivative (20a) and its β -anomer (20b) in 66% yield, along with the recovery of 15 (24%) (Scheme 6). Separation of these isomers was achieved by removal of PMB groups and the subsequent acetylation, affording the 21a in 51% (total yield from 15) and its β -anomer (21b) in 9%. Removal of the protecting groups of 21a was accomplished as usual, furnishing the desired 2'-O-(α -Dglucopyranosyl)biopterin (1). The α -anomeric structure of 21a was derived from its $J_{1,2}$ value (3.9 Hz) of ¹H-NMR, while the larger $J_{1,2}$ value (8.1 Hz) confirmed the β -form of 21b [23].

We have developed an effective way for selective preparation of both pterin 2'-O- β - and 2'-O- α -glycosides. By use of the key intermediate 1'-O-PMB-biopterin derivative (14) and the novel glycosyl donor (19) the first synthesis of biopterin α -D-glucoside (1) was achieved.

Acknowledgements

We are grateful to the SC-NMR Laboratory of Okayama University for the NMR measurements and to WESCO Scientific Promotion Foundation (to T. H.) which partially supported this work.

References

- Forrest HS, van Baalen C, Myers J. Isolation and identification of a new pteridine from a blue-green alga, Arch. Biochem. Biophys 1958; 78: 95-99.
- Hatfield DL, van Baalen C, Forrest HS. Pteridines in blue green algae, Plant Physiol 1961; 36: 240-243.
- Matsunaga T, Burgess JG, Yamada N, Komatsu K, Yoshida S, Wachi Y. An ultraviolet (UV-A) absorbing biopterin glucoside from the marine planktonic cyanobacterium *Oscillatoria* sp, Appl. Microbiol. Biotechnol 1993; 39: 250-253.
- Noguchi Y, Ishii A, Matsushima A, Haishi D, Yasumuro K, Moriguchi T, Wada T, Kodera Y, Hiroto M, Nishimura H, Sekine M, Inada Y. Isolation of biopterin-α-glucoside from Spirulina (*Arthrospira*) platensis and its physiologic function, Mar. Biotechnol 1999; 1: 207-210.
- 5. Chung HJ, Kim Y-A, Kim YJ, Choi

Pteridines / Vol. 20 / Special issue

YK, Hwang YK, Park YS. Purification and characterization of UDP-glucose: tetrahydrobiopterin glucosyltransferase from *Synechococcus* sp. PCC 7942. Biochim. Biophys. Acta 2000; 1524: 183-188.

- Choi YK, Hwang YK, Kang YH, Park YS. Chemical structure of 1-O-(L-erythrobiopterin-2'-yl)-α-glucose isolated from a cyanobacterium Synechococcus sp. PCC 7942, Pteridines 2001; 12: 121-125.
- Cha KW, Pfleiderer W, Yim JJ. Isolation and characterization of limipterin (1-O-(L-erythro-biopterin-2'-yl)-β-Nacetylglucosamine) and its 5,6,7,8-tetrahydro derivative from green sulfur bacterium *Chlorobium limicola f. thiosulfatophilum* NCIB 8327, Helv. Chim. Acta 1995; 78: 600-614.
- Cho SH, Na JU, Youn H, Hwang CS, Lee CH, Kang SO. Tepidopterin, 1-O-(L-threobiopterin-2'-yl)-β-N-acetylglucosamine from *Chlorobium tepidum*, Biochim. Biophys. Acta 1998; 1379: 53-60.
- Lin XL, White RH. Structure of solfapterin (erythro-neopterin-3'-D-2-deoxy-2aminoglucopyranoside) isolated from the thermophilic archaebacterium Sulfolobus solfataricus, J. Bacteriol 1988; 170: 1396-1398.
- Lee HW, Oh CH, Geyer A, Pfleiderer W, Park YS. Characterization of a novel unconjugated pteridine glycoside, cyanopterin, in *Synechocystis* sp. PCC 6803, Biochim. Biophys. Acta 1999; 1410: 61-70.
- Yamamoto H, Hanaya T, Harada K, Kawamoto H, Pfleiderer W. An efficient synthesis of limipterin [2'-O-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-L-biopterin] via N²-(N,N-dimethylamino)methylene-3-[2-(4-nitrophenyl)ethyl]-L-biopterin, Pteridines 1996; 7: 110-112.
- Hanaya T, Soranaka K, Harada K, Yamaguchi H, Suzuki R, Endo Y, Yamamoto H, Pfleiderer W. An efficient synthesis of 2'-O-(β-D-glucopyranosyl)- and 2'-O-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-L-biopterins, Heterocycles 2006; 67: 299-310.
- Hanaya T, Toyota H, Yamamoto H. Novel preparation of a 2'-O-acetyl-1'-O-(4methoxybenzyl)-L-biopterin derivative, a versatile precursor for a selective synthesis of L-biopterin glycosides, Synlett 2006; 2075-2078.

- 14. Hanaya T, Baba H, Yamamoto H. First synthesis of tepidopterin [2'-O-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-L-*threo*-biopterin], Carbohydr. Res 2007; 342: 2159-2162.
- Hanaya T, Baba H, Toyota H, Yamamoto H. Efficient total syntheses of natural pterin glycosides: limipterin and tepidopterin, Tetrahedron 2008; 64: 2090-2100.
- Hanaya T, Baba H, Kanemoto M, Yamamoto H. An efficient synthetic route for a versatile ciliapterin derivative and the first ciliapterin D-mannoside synthesis, Heterocycles 2008; 76: 635-644.
- Hanaya T, Torigoe K, Soranaka K, Fujita H, Yamamoto H, Pfleiderer W. An efficient synthesis of 2'-O-(β-D-ribofuranosyl)biopterin, Pteridines 2008; 19: 72-78.
- Pfleiderer W. In: Katritzky AR, ed. Physical methods in heterocyclic chemistry. New York: Academic Press 1963; 1: 177-188.
- Hanaya T, Torigoe K, Soranaka K, Yamamoto H, Yao Q, Pfleiderer W. Selective N(3)- and O⁴-alkylation of L-biopterin: a convenient synthesis of 3- and O⁴methyl-L-biopterin and the versatile N²-(N, N-dimethylaminomethylene)-N(3)p-nitrophenethyl-protected L-biopterin, Pteridines 1995; 6: 1-7.

- 20. Kiss J, D'Souza R, Taschner P. Präparative Herstellung von 5-Desoxy-L-arabinose, Xylit und D-Ribose aus Diacetonglucose, Helv. Chim. Acta 1975; 58: 311-317.
- Wulff G, Röhle G. Results and problems of O-glycoside synthesis, Angew. Chem., Int. Ed. Engl 1974; 13: 157-170, and references cited therein.
- 22. Ibatullin FM, Shabalin KA, Jänis JV, Shavva AG. Reaction of 1,2-*trans*-glycosyl acetates with thiourea: a new entry to 1-thiosugars, Tetrahedron Lett 2003; 44: 7961-7964.
- 23. A part of the results have been reported as a preliminary communication: Hanaya T, Baba H, Yamamoto H. First synthesis of biopterin α -D-glucoside, Heterocycles 2009; 77: 747-753.