# JOURNAL OF GENERAL VIROLOGY

#### **ICTV VIRUS TAXONOMY SUMMARY**

Rubino et al., Journal of General Virology 2025;106:002114 DOI 10.1099/jgv.0.002114





# Summary of taxonomy changes ratified by the International Committee on Taxonomy of Viruses from the Plant Viruses Subcommittee, 2025

Luisa Rubino<sup>1,\*</sup>, Peter Abrahamian<sup>2</sup>, Wenxia An<sup>3</sup>, Miquel A. Aranda<sup>4</sup>, José T. Ascencio-Ibañez<sup>5</sup>, Nicolas Bejerman<sup>6</sup>, Arnaud G. Blouin<sup>7</sup>, Thierry Candresse<sup>8</sup>, Tomas Canto<sup>9</sup>, Mengji Cao<sup>10</sup>, John P. Carr<sup>11</sup>, Won Kyong Cho<sup>12</sup>, Fiona Constable<sup>13</sup>, Indranil Dasgupta<sup>14</sup>, Humberto Debat<sup>6</sup>, Ralf G. Dietzgen<sup>15</sup>, Michele Digiaro<sup>16</sup>, Livia Donaire<sup>4</sup>, Toufic Elbeaino<sup>16</sup>, Denis Fargette<sup>17</sup>, Fiona Filardo<sup>18</sup>, Matthias G. Fischer<sup>19</sup>, Nuria Fontdevila<sup>7</sup>, Adrian Fox<sup>20</sup>, Juliana Freitas-Astua<sup>21</sup>, Marc Fuchs<sup>22</sup>, Andrew D.W. Geering<sup>15</sup>, Mahan Ghafari<sup>23</sup>, Anders Hafrén<sup>24</sup>, John Hammond<sup>25</sup>, Rosemarie Hammond<sup>26</sup>, Beata Hasiów-Jaroszewska<sup>27</sup>, Eugenie Hebrard<sup>28</sup>, Carmen Hernández<sup>29</sup>, Jean-Michel Hily<sup>30</sup>, Ahmed Hosseini<sup>31</sup>, Roger Hull<sup>32</sup>, Alice K. Inoue-Nagata<sup>33</sup>, Ramon Jordan<sup>25</sup>, Hideki Kondo<sup>34</sup>, Jan F. Kreuze<sup>35</sup>, Mart Krupovic<sup>36</sup>, Kenji Kubota<sup>37</sup>, Jens H. Kuhn<sup>38</sup>, Scott Leisner<sup>39</sup>, Jean-Michel Lett<sup>40</sup>, Chengyu Li<sup>3</sup>, Fan Li<sup>41</sup>, Jun Min Li<sup>42</sup>, Paola M. López-Lambertini<sup>43</sup>, Juan J. Lopez-Moya<sup>44</sup>, Francois Maclot<sup>8</sup>, Kristiina Mäkinen<sup>45</sup>, Darren Martin<sup>46</sup>, Sebastien Massart<sup>47</sup>, W. Allen Miller<sup>48</sup>, Musa Mohammadi<sup>49</sup>, Dimitre Mollov<sup>50</sup>, Emmanuelle Muller<sup>51</sup>, Tatsuya Nagata<sup>52</sup>, Jesús Navas-Castillo<sup>53</sup>, Yutaro Neriya<sup>54</sup>, Francisco M. Ochoa-Corona<sup>55</sup>, Kazusato Ohshima<sup>56</sup>, Vicente Pallás<sup>29</sup>, Hanu Pappu<sup>57</sup>, Karel Petrzik<sup>58</sup>, Mikhail Pooggin<sup>28</sup>, Maria Isabella Prigigallo<sup>1</sup>, Pedro L. Ramos-González<sup>59</sup>, Simone Ribeiro<sup>60</sup>, Katja R. Richert-Pöggeler<sup>61</sup>, Philippe Roumagnac<sup>62</sup>, Avijit Roy<sup>26</sup>, Sead Sabanadzovic<sup>63</sup>, Dana Šafářová<sup>64</sup>, Pasquale Saldarelli<sup>1</sup>, Hélène Sanfaçon<sup>65</sup>, Cecilia Sarmiento<sup>66</sup>, Takahide Sasaya<sup>67</sup>, Kay Scheets<sup>68</sup>, Willem E.W. Schravesande<sup>69</sup>, Susan Seal<sup>70</sup>, Yoshifumi Shimomoto<sup>71</sup>, Merike Sõmera<sup>66</sup>, Livia Stavolone<sup>1</sup>, Lucy R. Stewart<sup>72</sup>, Pierre-Yves Teycheney<sup>73</sup>, John E. Thomas<sup>15</sup>, Jeremy R. Thompson<sup>74</sup>, Antonio Tiberini<sup>75</sup>, Yasuhiro Tomitaka<sup>37</sup>, Ioannis Tzanetakis<sup>76</sup>, Marie Umber<sup>77</sup>, Cica Urbino<sup>28</sup>, Harrold A. van den Burg<sup>69</sup>, René A.A. Van der Vlugt<sup>78</sup>, Arvind Varsani<sup>79</sup>, Adriaan Verhage<sup>80</sup>, Dan Villamor<sup>76</sup>, Susanne von Bargen<sup>81</sup>, Peter J. Walker<sup>82</sup>, Thierry Wetzel<sup>83</sup>, Anna E. Whitfield<sup>84</sup>, Stephen J. Wylie<sup>85</sup>, Caixia Yang<sup>3</sup>, F. Murilo Zerbini<sup>86</sup>, Song Zhang<sup>10</sup> and ICTV Taxonomy Summary Consortium

# Abstract

In March 2025, following the annual International Committee on Taxonomy of Viruses (ICTV) ratification vote, newly proposed taxa were added to those under the mandate of the Plant Viruses Subcommittee. In brief, 1 new order, 3 new families, 6 new genera, 2 new subgenera and 206 new species were created. Some taxa were reorganized. Genus *Cytorhabdovirus* in the family *Rhabdoviridae* was abolished and its taxa were redistributed into three new genera *Alphacytorhabdovirus*, *Betacytorhabdovirus* and *Gammacytorhabdovirus*. Genus *Waikavirus* in the family *Secoviridae* was reorganized into two subgenera (*Actinidivirus* and *Ritunrivirus*). One family and four previously unaffiliated genera were moved to the newly established order *Tombendovirales*. Twelve species not assigned to a genus were abolished. To comply with the ICTV mandate of a binomial format for virus species, eight species were renamed. Demarcation criteria in the absence of biological information were defined in the genus *Ilarvirus* (family *Bromoviridae*). This article presents the updated taxonomy put forth by the Plant Viruses Subcommittee and ratified by the ICTV.

Received 15 April 2025; Accepted 13 May 2025; Published 25 July 2025

For numbered affiliations see end of the article.

\*Correspondence: Luisa Rubino, luisa.rubino@cnr.it

**Abbreviations:** CP, coat protein; ICTV, International Committee on Taxonomy of Viruses; Pol, polymerase; Pro, protease. Supplementary files are available with the online version of this article.

002114



#### INTRODUCTION

The Plant Viruses Subcommittee of the International Committee on Taxonomy of Viruses (ICTV) deals with the taxonomic classification of most viruses infecting or isolated from plants [1]. It consists of 22 Study Groups, composed of more than 180 members, covering 27 virus families.

Viruses, viroids, satellite viruses and satellite nucleic acids originally isolated from plants have been classified into 2,392 species belonging to 247 genera in 55 families, 25 orders, 16 classes, 7 phyla, 3 kingdoms and 2 realms [2]. Following the 2025 ICTV ratification vote, these taxa were expanded by creating 1 new order, 3 new families, 6 new genera, 2 new subgenera and 206 new species.

To comply with the ICTV mandate of a binomial format for virus species [3, 4], species names will consist of the genus name followed by a Latinized or not Latinized (freeform) epithet [5, 6]. The adoption of the binomial format for virus species names was completed by renaming eight species. Similarly, 12 species in the families *Rhabdoviridae* and *Tombusviridae* were abolished because they could not be assigned to a genus due to the lack of a genome sequence. Viruses in many of the new species were discovered by metagenomic analysis and were classified according to Simmonds *et al.* [7].

Some established taxa have been extensively reorganized. The family *Rhabdoviridae* has been expanded by the creation of one new genus, whereas the genus *Cytorhabdovirus* was abolished and replaced with three new genera *Alphacytorhabdovirus*, *Betacytorhabdovirus* and *Gammacytorhabdovirus*, representing distinct evolutionary lineages. In the family *Secoviridae*, one new genus was created and the established genus *Waikavirus* was reorganized into two subgenera. The establishment of the new order *Tombendovirales* allowed the placement of four previously unaffiliated genera of plant satellite viruses into four families based on high-resolution capsid structure studies rather than sequence identity.

The classification of viruses discovered by metagenomics prompted some refinement of demarcation criteria in the absence of biological information. For instance, species demarcation criteria based on 'serology, host range and sequence similarity' in the genus *Ilarvirus*, family *Bromoviridae*, have been refined to require less than 85% amino acid identity for the complete RNA2-encoded 2a protein.

All of these changes contribute to the advancement of virus taxonomy. To disseminate taxonomy decisions, the annual publication of summaries of all taxonomy proposals from each ICTV Subcommittee will provide a compendium of the taxonomy changes [8]. Therefore, the updated taxonomy from the Plant Viruses Subcommittee as now accepted by the ICTV is summarized in this article. It should be noted that the article does not necessarily cover all changes in the taxonomy of plant viruses, and consultation of reports from the Fungal and Protist Viruses Subcommittee [9] and the Animal dsRNA and ssRNA(–) Viruses Subcommittee [10] may be needed. A file including all the Tables of taxonomic changes below is available as a supplementary file to this article.

#### **MAIN TEXT**

#### CONTENTS

2024.001P.Fimoviridae 1nsp 2024.002P.Alphaflexiviridae\_7nsp 2024.003P.Tospoviridae\_2nsp 2024.004P.Konkoviridae 1nsp 2024.005P.Caulimoviridae\_3nsp 2024.006P.Kitaviridae\_3nsp 2024.007P.Geminiviridae Capulavirus 1nsp 2024.008P.Geminiviridae\_Citlodavirus\_2nsp 2024.009P.Geminiviridae\_Mastrevirus\_5nsp 2024.010P.Geminiviridae Begomovirus 18nsp 2024.011P.Bromoviridae\_4nsp 2024.012P.Potyviridae 1ng 10nsp 2024.013P.Secoviridae\_1ng\_2nsg\_34nsp 2024.014P.Rhabdoviridae\_2nsp 2024.015P.Rhabdoviridae Cytorhabdovirus splitgen 2024.016P.Rhabdoviridae\_1ngen\_5nsp 2024.017P.Tombusviridae abolishsp 2024.018P.Tombusviridae 1nsp 2024.019P.Closteroviridae\_1nsp 2024.021P.Riboviria\_1nord 2024.022P.Betaflexiviridae\_abolishsp

2024.023P.Solemoviridae\_rename\_sp 2024.024P.Tymoviridae\_abolish\_sp

# 2024.001P.Fimoviridae\_1nsp

Title: Create Emaravirus clematis as a new species in the genus Emaravirus, family Fimoviridae

Authors: Yang C., An W., Li C., Zhang S., Cao M., Digiaro M. (digiaro@iamb.it), Elbeaino T., Kubota K., Ochoa Corona F.M., von Bargen S.

#### **Summary**

#### Taxonomic rank(s) affected

Species

# Description of current taxonomy

The family Fimoviridae currently includes 32 virus species in the genus Emaravirus.

#### Proposed taxonomic change(s)

Add one (1) new virus species to genus *Emaravirus* within the family *Fimoviridae*.

#### **Justification**

The creation of the new species *Emaravirus clematis* in the genus *Emaravirus*, family *Fimoviridae*, is proposed to accommodate Clematis yellow mottle-associated virus, identified in China on *Clematis brevicaudata* DC. The virus assigned to the new species consists of a fully sequenced five-segmented, linear, single-stranded (ss), negative-sense RNA genome (of which two RNA3s encode the nucleocapsid protein), which shows features common to homologous RNAs of viruses assigned to other *Emaravirus* species but from which they differ significantly in nucleotide and amino acid sequences.

Submitted: 30/04/24; Revised: 16/09/24

**Table 1.** Fimoviridae, 1 new taxon\*

| Operation | Rank    | New taxon name      | Virus name                                  | Exemplar   |
|-----------|---------|---------------------|---|--|
| New taxon | Species | Emaravirus clematis | Clematis yellow mottle-<br>associated virus | RNA1: OP807964; RNA2: OP807965; RNA3a: OP807966; RNA3b: OP807967; RNA4: OP807968 |

 $<sup>*</sup>Source/full\ text:\ https://ictv.global/ictv/proposals/2024.001P.Fimoviridae\_1nsp.zip.$ 

# 2024.002P.Alphaflexiviridae\_7nsp

Title: Create seven new species in the family Alphaflexiviridae

**Authors:** Abrahamian P., Donaire L., Candresse T., Fox A., Hammond J., Hasiów-Jaroszewska B., Kreuze J., Rubino L., Aranda M.A. (m.aranda@cebas.csic.es)

#### **Summary**

# Taxonomic rank(s) affected

Species

#### **Description of current taxonomy**

The family Alphaflexiviridae currently includes 65 virus species in genera Allexivirus (13), Botrexvirus (1), Lolavirus (1), Platypuvirus (1), Potexvirus (48) and Sclerodarnavirus (1).

#### Proposed taxonomic change(s)

Add seven (7) new virus species to genera *Allexivirus* (1), *Botrexvirus* (2) and *Potexvirus* (4) within the family *Alphaflexiviridae*. **Justification** 

Throughout the family, isolates of viruses assigned to different species have less than 72% nucleotide identity (or 80% amino acid identity) among their respective coat protein or polymerase genes (or proteins). Viruses from different genera usually have <45% nucleotide identity in these genes. The nucleotide or amino acid sequences of viruses belonging to the seven newly proposed species fit well within these demarcation criteria.

**Submitted**: 14/06/24

Table 2. Alphaflexiviridae, 7 new taxa\*

| Operation | Rank    | New taxon name              | Virus name                                 | Exemplar |
|-----------|---------|-----------------------------|--|----------|
| New taxon | Species | Allexivirus rehmanniae      | Rehmannia allexivirus                      | PP097219 |
| New taxon | Species | Botrexvirus unosclerotiniae | Sclerotinia sclerotiorum alphaflexivirus 1 | ON993219 |
| New taxon | Species | Botrexvirus duosclerotiniae | Sclerotinia sclerotiorum alphaflexivirus 2 | OQ865609 |
| New taxon | Species | Potexvirus ecsadenii        | Adenium obesum virus X                     | OR039325 |
| New taxon | Species | Potexvirus chaenostomae     | Chaenostoma potexvirus                     | OL979628 |
| New taxon | Species | Potexvirus ecshibisci       | Hibiscus virus X                           | PP115950 |
| New taxon | Species | Potexvirus ecscaricae       | papaya virus X                             | MN265368 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.002P.Alphaflexiviridae\_7nsp.zip.

# 2024.003P.Tospoviridae\_2nsp

Title: Create two new species in the genus Orthotospovirus (Elliovirales: Tospoviridae)

Authors: Tomitaka Y., Shimomoto Y., Sasaya T. (tsasaya@affrc.go.jp)

# **Summary**

#### Taxonomic rank(s) affected

Species

# Description of current taxonomy

The family *Tospoviridae* currently includes 28 virus species in the genus *Orthotospovirus*.

## Proposed taxonomic change(s)

Add two (2) new virus species to genus *Orthotospovirus* within the family *Tospoviridae*.

#### **Justification**

It is proposed that two (2) newly discovered tospovirids be classified into new species in the genus *Orthotospovirus* on the basis of species demarcation criteria in the amino acid sequence of the RNA-directed RNA polymerase and nucleocapsid protein. **Submitted:** 13/06/24

Table 3. Tospoviridae, 2 new taxa\*

| Operation | Rank    | New taxon name           | Virus name                                | Exemplar                     |
|-----------|---------|--------------------------|---|------------------------------|
| New taxon | Species | Orthotospovirus eustomae | lisianthus necrotic ringspot virus        | MF469045; MF469046; MF469047 |
| New taxon | Species | Orthotospovirus fatsiae  | Fatsia japonica ringspot-associated virus | LC626335; LC626336; LC626337 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.003P.Tospoviridae\_2nsp.zip.

# 2024.004P.Konkoviridae\_1nsp

Title: Create one new species in the genus Olpivirus (Hareavirales: Konkoviridae)

Authors: Neriya Y., Schravesande W.E.W., van den Burg H.A., Verhage A., Tomitaka Y., Sasaya T. (tsasaya@affrc.go.jp)

# **Summary**

#### Taxonomic rank(s) affected

Species

#### Description of current taxonomy

The family Konkoviridae currently includes one virus species in the genus Olpivirus.

#### Proposed taxonomic change(s)

Add one (1) new virus species to genus Olpivirus within the family Konkoviridae.

#### Justification

It is proposed that one (1) newly discovered konkovirid be classified into a new species in the genus *Olpivirus* on the basis of a species demarcation criterion of <95% identity in the amino acid sequence of the RNA-directed RNA polymerase.

Submitted: 13/06/24; Revised: 07/10/24

Table 4. Konkoviridae, 1 new taxon\*

| Operation | Rank    | New taxon name     | Virus name                              | Exemplar   |
|-----------|---------|--------------------|---|--|
| New taxon | Species | Olpivirus lactucae | Lactuca big vein associated phlebovirus | RNA1: OR610326; RNA2: OR610327; RNA3: OR610328; RNA4: OR610329 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.004P.Konkoviridae 1nsp.zip.

# 2024.005P.Caulimoviridae\_3nsp

**Title:** Create three new species in the genus *Badnavirus* (*Ortervirales*: *Caulimoviridae*)

**Authors:** Umber M., Dasgupta I., Geering A.D.W., Hafrén A., Hull R., Kreuze J., Leisner S., Muller E., Pappu H., Pooggin M., Richert-Pöggeler K.R., Seal S., Stavolone L., Teycheney P.Y. (teycheney@cirad.fr)

#### **Summary**

#### Taxonomic rank(s) affected

Genus (Badnavirus)

# Description of current taxonomy

The family *Caulimoviridae* [11] currently comprises 11 genera whose members share a similar genome organization. The molecular species demarcation criterion is <80% identity of nucleotide sequences in the reverse transcriptase/ribonuclease H region of the polymerase. Genus *Badnavirus* [12] currently includes 71 species and is the largest genus within the family *Caulimoviridae*.

#### Proposed taxonomic change(s)

Add three (3) new species to the genus Badnavirus (Badnavirus fatsiae, Badnavirus tetainflatheobromae and Badnavirus ziziphi). **Iustification** 

Complete genomes of the exemplar members of the three proposed new species were sequenced and published recently. Their organizations are similar to those of other members of the genus *Badnavirus*. Phylogenetic analyses place them in this genus as distinct representatives of novel species.

**Submitted:** 14/06/2024

Table 5. Caulimoviridae, 3 new taxa\*

| Operation | Rank    | New taxon name                 | Virus name                        | Exemplar |
|-----------|---------|--------------------------------|-----------------------------------|----------|
| New taxon | Species | Badnavirus fatsiae             | Fatsia badnavirus 1               | OM540428 |
| New taxon | Species | Badnavirus tetainflatheobromae | cacao swollen shoot Ghana T virus | MN179342 |
| New taxon | Species | Badnavirus ziziphi             | jujube badnavirus WS              | OL739567 |

 $<sup>*</sup>Source/full\ text: https://ictv.global/ictv/proposals/2024.005P. Caulimoviridae\_3nsp.zip.$ 

# 2024.006P.Kitaviridae\_3nsp

Title: Create a new species in the genus Cilevirus and two in the genus Higrevirus, family Kitaviridae (Martellivirales)

Authors: Li C., An W., Zhang S., Cao M., Yang C. (xueyang27@126.com), Mohammadi M., Hosseini A., Nasrollanejad S., Roy A., Freitas-Astua J., Tiberini A., Li J.M., Ramos-González P.L.

#### **Summary**

#### Taxonomic rank(s) affected

Species

#### Description of current taxonomy

Family *Kitaviridae*, order *Martellivirales*, includes plant-infecting viruses having linear single-stranded (ss) positive-sense (+) segmented RNA genomes. Viruses in this family are assigned to the genera *Cilevirus*, *Higrevirus* or *Blunervirus* [13, 14].

## Proposed taxonomic change(s)

Add three (3) new species to the family *Kitaviridae*, one (1) to the genus *Cilevirus* and two (2) to the genus *Higrevirus*.

#### **Justification**

The genomes of the three novel viruses show an arrangement that resembles that of kitavirids, and their core conserved proteins share relatively low amino acid sequence identities (<85%) with recognized members of the family *Kitaviridae*. In phylogenetic analyses, the three viruses group with characterized members of the genera *Cilevirus* and *Higrevirus*, but they

are well-separated and supported by bootstrap values >95%. All new species meet the already established or the demarcation criteria defined in this proposal.

**Submitted:** 11/06/2024

Table 6. Kitaviridae, 3 new taxa\*

| Operation | Rank    | New taxon name       | Virus name                                | Exemplar                                       |
|-----------|---------|----------------------|---|--|
| New taxon | Species | Higrevirus amurense  | Phellodendron-associated higre-like virus | RNA1: OP324809; RNA2: OP324810; RNA3: OP324811 |
| New taxon | Species | Higrevirus pistaciae | pistachio virus X                         | RNA1: MT334620; RNA2: MT334619; RNA3: MT334618 |
| New taxon | Species | Cilevirus pistaciae  | pistachio virus Y                         | RNA1: MT362606; RNA2: MT362605                 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.006P.Kitaviridae\_3nsp.zip.

# 2024.007P.Geminiviridae Capulavirus 1nsp

Title: Create one new species in the genus Capulavirus (Geplafuvirales: Geminiviridae)

Authors: Roumagnac P. (philippe.roumagnac@cirad.fr), Ascencio-Ibanez J., Lett J.-M., López-Lambertini P.M., Martin D.P., Navas-Castillo J., Ribeiro S., Urbino C., Varsani A., Zerbini F.M.

#### **Summary**

#### Taxonomic rank(s) affected

Genus (Capulavirus)

#### **Description of current taxonomy**

Monodnaviria: Shotokuvirae: Cressdnaviricota: Repensiviricetes: Geplafuvirales: Geminiviridae: Capulavirus.

#### Proposed taxonomic change(s)

Add one (1) new species to the genus Capulavirus (Capulavirus trifolii).

#### **Justification**

Similar to members of the *Capulavirus* genus, members of the proposed new species *Capulavirus trifolii* have the virion-strand origin of replication nonanucleotide motif 'TAATATTAC' and show a typical capulavirus genome organization, with putative multiple overlapping short ORFs (V3 and V4) upstream of the coat protein gene that encode putative movement proteins. In addition, genome-wide pairwise analysis of the representative genomes of capulaviruses showed that viruses in the species *Capulavirus trifolii* share <78% identity with all representative genomes of capulaviruses. Because 78% nucleotide identity is the genome-wide species demarcation threshold for capulaviruses, it was concluded that *Capulavirus trifolii* represents a new species in the genus *Capulavirus*.

**Submitted:** 10/06/24

Table 7. Geminiviridae, 1 new taxon\*

| Operation | Rank    | New taxon name       | Virus name        | Exemplar |
|-----------|---------|----------------------|-------------------|----------|
| New taxon | Species | Capulavirus trifolii | Trifolium virus 1 | MW698813 |

 $<sup>*</sup>Source/full\ text:\ https://ictv.global/ictv/proposals/2024.007P. Geminiviridae\_Capulavirus\_1 nsp.zip.$ 

# 2024.008P.Geminiviridae\_Citlodavirus\_2nsp

Title: Create two new species in the genus Citlodavirus (Geplafuvirales: Geminiviridae)

**Authors:** Roumagnac P. (philippe.roumagnac@cirad.fr), Ascencio-Ibanez J., Lett J.-M., López-Lambertini P.M., Martin D.P., Navas-Castillo J., Ribeiro S., Urbino C., Varsani A., Zerbini F.M.

#### Summary

# Taxonomic rank(s) affected

Genus (Citlodavirus)

#### Description of current taxonomy

Monodnaviria: Shotokuvirae: Cressdnaviricota: Repensiviricetes: Geplafuvirales: Geminiviridae: Citlodavirus.

# Proposed taxonomic change(s)

Add two (2) new species to genus Citlodavirus (Citlodavirus apijamaicaense and Citlodavirus myricae).

#### Iustification

Similar to members of the *Citlodavirus* genus, members of the proposed new species *Citlodavirus apijamaicaense* and *Citlodavirus myricae* have the virion-strand origin of replication nonanucleotide motif 'TAA TAT TAC', a relatively large genome (3,918 and 3,775 nt, respectively) and unique genome arrangements that, in both cases, include the putative *mp* gene (888 and 912 nt, respectively) that is similar in size to the *mp* gene in the DNA-B of bipartite begomoviruses. Genome-wide pairwise analysis of the representative genomes of citlodaviruses showed that viruses in the species *Citlodavirus apijamaicaense* and *Citlodavirus myricae* share <78% identity with all representative genomes of citlodaviruses and with each other. Since 78% nucleotide identity is the genome-wide species demarcation threshold for citlodaviruses, it has been concluded that *Citlodavirus apijamaicaense* and *Citlodavirus myricae* represent two new species in the genus *Citlodavirus*.

Submitted: 10/06/2024

Table 8. Geminiviridae, 2 new taxa\*

| Operation | Rank    | New taxon name              | Virus name                  | Exemplar |
|-----------|---------|-----------------------------|-----------------------------|----------|
| New taxon | Species | Citlodavirus apijamaicaense | apiscitlodal virus          | PP467584 |
| New taxon | Species | Citlodavirus myricae        | Myrica rubra citlodavirus 1 | OP374189 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.008P.Geminiviridae\_Citlodavirus\_2nsp.zip.

# 2024.009P.Geminiviridae\_Mastrevirus\_5nsp

Title: Establish five new species in the genus Mastrevirus

**Authors:** Varsani A. (Arvind.varsani@asu.edu), Martin D.P., Roumagnac P., Ascencio-Ibanez J., Lett J.-M., López-Lambertini P.M., Navas-Castillo J., Ribeiro S., Urbino C., Zerbini F.M.

#### **Summary**

#### Taxonomic rank(s) affected

Species

#### **Description of current taxonomy**

Monodnaviria: Shotokuvirae: Cressdnaviricota: Repensiviricetes: Geplafuvirales: Geminiviridae: Mastrevirus. Within the genus Mastrevirus, viruses are classified into species based on a 78% genome-wide pairwise identity threshold [15].

#### Proposed taxonomic change(s)

Add five (5) new species to classify a suite of new mastreviruses that have been identified over the last year or so.

#### **Justification**

The members of the five new proposed species in the genus *Mastrevirus* share <78% genome-wide pairwise identity with sequences of members of currently established mastrevirus species.

**Submitted:** 10/06/24

Table 9. Geminiviridae, 5 new taxa\*

| Operation | Rank    | New taxon name                  | Virus name  | Exemplar |
|-----------|---------|---------------------------------|---|----------|
| New taxon | Species | Mastrevirus urochloareunionense | Urochloa decumbens associated virus                         | OQ451139 |
| New taxon | Species | Mastrevirus nomiae              | nomiamastrel virus  | PP467585 |
| New taxon | Species | Mastrevirus brachypodiumprimi   | Brachypodium phoenicoides associated virus $\boldsymbol{1}$ | OR596402 |
| New taxon | Species | Mastrevirus bothriochloae       | Bothriochloa barbinodis associated virus                    | OR596403 |
| New taxon | Species | Mastrevirus brachypodiumsecundi | Brachypodium phoenicoides associated virus 2                | OR596405 |

<sup>\*</sup>Source/full text:https://ictv.global/ictv/proposals/2024.009P.Geminiviridae\_Mastrevirus\_5nsp.zip.

# 2024.010P.Geminiviridae\_Begomovirus\_18nsp

**Title:** Create 18 new species in the genus *Begomovirus* (*Geplafuvirales*: *Geminiviridae*)

**Authors:** Zerbini F.M. (zerbini@ufv.br), Ascencio-Ibanez J., Lett J.M., Navas-Castillo J., Urbino C., López-Lambertini P.M., Martin D.P., Ribeiro S.G., Roumagnac P., Varsani A.

#### **Summary**

# Taxonomic rank(s) affected

Species in the genus *Begomovirus* 

# Description of current taxonomy

Monodnaviria: Shotokuvirae: Cressdnaviricota: Repensiviricetes: Geplafuvirales: Geminiviridae: Begomovirus. Within the genus Begomovirus, viruses are classified into species based on a 91% genome-wide (or DNA-A in the case of bipartite viruses) pairwise identity threshold [16].

# Proposed taxonomic change(s)

Creation of 18 new species to classify new begomoviruses that have been identified and described in the literature over the last 3 years.

# **Justification**

All 18 proposed new species have <91% genome-wide (or DNA-A in the case of bipartite viruses) pairwise identity with sequences of members of currently established begomovirus species.

**Submitted:** 10/06/24

Table 10. Begomovirus, 18 new taxa\*

| Operation | Rank    | New taxon name                     | Virus name                                | Exemplar           |
|-----------|---------|------------------------------------|---|--------------------|
| New taxon | Species | Begomovirus chuxiongense           | tomato leaf curl Chuxiong virus           | OR543988           |
| New taxon | Species | Begomovirus<br>solanumaureusreti   | tomato golden net virus                   | MT214095           |
| New taxon | Species | Begomovirus<br>solanumflavusreti   | tomato yellow net virus                   | MT214096           |
| New taxon | Species | Begomovirus whitaniae              | Withania leaf curl virus                  | OP617239           |
| New taxon | Species | Begomovirus cajani                 | Cajanus scarabaeoides yellow mosaic virus | OM397101; OM397102 |
| New taxon | Species | Begomovirus hortuscrotoni          | garden croton enation leaf<br>curl virus  | MW816855; MW816857 |
| New taxon | Species | Begomovirus<br>jatrophagunturense  | Jatropha leaf curl Guntur<br>virus        | MZ217773           |
| New taxon | Species | Begomovirus hyptidis               | Hyptis golden mosaic virus                | ON073795; ON073796 |
| New taxon | Species | Begomovirus galii                  | Galium leaf distortion virus              | OL689630           |
| New taxon | Species | Begomovirus myanmarense            | tobacco curly shoot<br>Myanmar virus      | MK920410           |
| New taxon | Species | Begomovirus caboniensis            | Cnidoscolus mild mosaic virus             | MZ465527; MZ465585 |
| New taxon | Species | Begomovirus pyrenacanthae          | Pyrenacantha yellow mosaic virus          | MZ390982; MZ390984 |
| New taxon | Species | Begomovirus puerense               | tobacco leaf curl Puer virus              | MZ465370           |
| New taxon | Species | Begomovirus<br>solanumdistorsionis | tomato mottle leaf distortion virus       | MW561191; MW650837 |
| New taxon | Species | Begomovirus alceacrispi            | hollyhock vein yellowing virus            | LK028571           |
| New taxon | Species | Begomovirus muntiflavi             | Muntingia yellow spot virus               | MW032664; MW032665 |
| New taxon | Species | Begomovirus flavintervenae         | tomato interveinal yellowing virus        | MW057360           |
| New taxon | Species | Begomovirus sidaflavitessellati    | Sida yellow mosaic Gujarat<br>virus       | KX513859           |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.010P.Geminiviridae\_Begomovirus\_18nsp.zip.

# 2024.011P.Bromoviridae\_4nsp

**Title:** Create four (4) new species in the genus *Ilarvirus* (*Martellivirales*: *Bromoviridae*)

Authors: Thompson J.R. (jeremy.thompson@mpi.govt.nz), Canto T., Carr J.P., Pallás V., Šafářová D.

# **Summary**

# Taxonomic rank(s) affected

**Species** 

#### Description of current taxonomy

The family *Bromoviridae* currently includes 49 virus species in genera *Alfamovirus* (1), *Anulavirus* (4), *Bromovirus* (7), *Cucumovirus* (4), *Ilarvirus* (32) and *Oleavirus* (1).

#### Proposed taxonomic change(s)

Add four (4) new virus species to genus *Ilarvirus* within the family *Bromoviridae*.

#### **Justification**

This taxonomic proposal considers the recognition of four new virus species based on species demarcation criteria in the family *Bromoviridae*, genus *Ilarvirus* of 'serology, host range and sequence similarity'. In the absence of biological information, it is proposed that a refinement of the 'sequence similarity' criterion be made to require that members of different species display <85% amino acid identity for the complete RNA2-encoded 2a protein.

Submitted: 10/06/24; Revised: 07/10/24

Table 11. Bromoviridae. 4 new taxa\*

| Operation | Rank    | New taxon name   | Virus name                  | Exemplar                     |
|-----------|---------|------------------|-----------------------------|------------------------------|
| New taxon | Species | Ilarvirus ApNMV  | apple necrotic mosaic virus | LC108993; LC108994; LC108995 |
| New taxon | Species | Ilarvirus BabIV1 | babaco ilarvirus 1          | OQ256238; OQ256239; OQ256240 |
| New taxon | Species | Ilarvirus TIV1   | tomato ilarvirus 1          | OL472057; OL472058; OL472059 |
| New taxon | Species | Ilarvirus ToNSV  | tomato necrotic spot virus  | MH780154; MH780155; MH780156 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.011P.Bromoviridae\_4nsp.zip.

# 2024.012P.Potyviridae\_1ng\_10nsp

Title: Create 1 new genus (Phragmivirus) with 2 species and 8 new species in the genus Potyvirus (Patatavirales: Potyviridae)

**Authors:** Inoue-Nagata A.K. (alice.nagata@embrapa.br), Jordan R., Kreuze J.F., Li F., Lopez-Moya J.J., Mäkinen K., Ohshima K., Wylie S.J.

# **Summary**

#### Taxonomic rank(s) affected

Genus within the family *Potyviridae* and species within the genus *Potyvirus* and the newly proposed genus *Phragmivirus* **Description of current taxonomy** 

According to the ICTV Report chapter on *Potyviridae*, 12 genera are differentiated by biological criteria, mainly transmission by specific vectors, and by molecular data, in which members of different genera are <46% identical in nucleotide sequence. Members of different species have complete ORF sequences that are generally <76% identical in nucleotide sequence and <82% identical in amino acid sequence. In considering the evidence for new species or genera in the family *Potyviridae*, the Study Group will evaluate each new case based on complete- or near-complete genome sequence(s) together with host and biological characteristics.

#### Proposed taxonomic change(s)

Create one (1) new genus (*Phragmivirus*), two (2) new species in the genus *Phragmivirus* (*Phragmivirus* phragmii and *Phragmivirus* spartinae) and eight (8) new species in the genus *Potyvirus* (*Potyvirus* aconiti, *Potyvirus* puerariae, *Potyvirus* alilii, *Potyvirus* parisflavitessellati, *Potyvirus* catharanthiflavitessellati, *Potyvirus* polygonatimaculae, *Potyvirus* crocitessellati and *Potyvirus* galanthi).

# Justification

The genomes of the proposed members in the new genus *Phragmivirus* share sequence identity below the threshold for genus differentiation in the family *Potyviridae*; members of the proposed species have a genome strategy typical of members of genus *Phragmivirus* (two species) and *Potyvirus* (eight species), and their nucleotide and amino acid sequences are below the threshold for species demarcation criteria for the genera.

Submitted: 11/06/24; Revised: 21/09/24

Table 12. Potyviridae, 11 new taxa\*

| Operation | Rank    | New taxon name                          | Virus name                               | Exemplar |
|-----------|---------|---|--|----------|
| New taxon | Species | Potyvirus aconiti                       | Aconitum virus 2                         | MZ389235 |
| New taxon | Species | Potyvirus puerariae                     | kudzu chlorotic ring blotch virus        | OQ148665 |
| New taxon | Species | Potyvirus alilii                        | lily virus A                             | OR879085 |
| New taxon | Species | Potyvirus parisflavitessellati          | Paris yunnanensis mosaic chlorotic virus | ON871824 |
| New taxon | Species | Potyvirus<br>catharanthiflavitessellati | periwinkle mild yellow<br>mosaic virus   | PP382205 |
| New taxon | Species | Potyvirus polygonatimaculae             | Polygonatum kingianum<br>mottle virus    | ON428226 |
| New taxon | Species | Potyvirus crocitessellati               | saffron yellow mosaic virus              | OK632024 |
| New taxon | Species | Potyvirus galanthi                      | snowdrop virus Y                         | OP871788 |
| New taxon | Genus   | Phragmivirus                            |  |          |
| New taxon | Species | Phragmivirus phragmii                   | common reed chlorotic stripe virus       | KY612317 |
| New taxon | Species | Phragmivirus spartinae                  | Spartina mottle virus                    | MN788417 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.012P.Potyviridae\_1ng\_10nsp.zip.

# 2024.013P.Secoviridae\_1ng\_2nsg\_34nsp

Title: Create a new genus, two new subgenera and 34 new species in the family Secoviridae (Picornavirales)

Authors: Fuchs M. (mf13@cornell.edu), Hily J.-M., Petrzik K., Sanfaçon H., Stewart L., Thompson J.R., Van der Vlugt R.A.A., Wetzel T.

#### Summary

#### Taxonomic rank(s) affected

Genus, subgenus, species

#### Description of current taxonomy

The recognition of new virus species is based on demarcation criteria in the family Secoviridae of <75% amino acid sequence identity in the coat protein (CP)(s) and/or <80% amino acid sequence identity in the conserved protease (Pro)-polymerase (Pol) region (from the protease CG motif to the polymerase GDD motif) and/or distinct plant hosts and biological properties.

#### Proposed taxonomic change(s)

Create one (1) new genus (Mersevirus), two (2) new subgenera in the genus Waikavirus (Ritunrivirus and Actinidivirus), create two (2) new species in the genus Fabavirus (Fabavirus betavitis and Fabavirus cirsii), create four (4) new species in the new genus Mersevirus (Mersevirus mercurialis, Mersevirus paris, Mersevirus boehmeriae and Mersevirus jujubae), create two (2) new species in the genus Nepovirus (Nepovirus betaparis and Nepovirus mirae), create three (3) new species in the genus Sadwavirus (Sadwavirus cattleyae, Sadwavirus gymnemae and Sadwavirus chrysanthemi), create three (3) new species in the genus Torradovirus (Torradovirus physalis, Torradovirus nanorugosum and Torradovirus arctii) and create 20 new species in the genus Waikavirus (Waikavirus ajugae, Waikavirus anacycli, Waikavirus betacamelliae, Waikavirus eleocharis, Waikavirus hirtae, Waikavirus juglandis, Waikavirus ligustici, Waikavirus mertensiae, Waikavirus populi, Waikavirus pedicularis, Waikavirus primulae, Waikavirus querci, Waikavirus ranunculi, Waikavirus thymi, Waikavirus trifoccidentale, Waikavirus thapsiae, Waikavirus violae, Waikavirus carotae, Waikavirus celtis and Waikavirus pittospori).

#### **Justification**

The proposed new genus *Mersevirus* is based on the distinct genome organization of *Mersevirus mercurialis*, *Mersevirus paris*, *Mersevirus boehmeriae* and *Mersevirus jujubae* with a Ham1 domain with predicted inosine triphosphate pyrophosphatase activity at the C-terminus of the RNA-directed RNA polymerase – a feature unique among members of the family *Secoviridae* – and a grouping on a monophyletic clade of the amino acid sequence of the CPs and conserved Pro-Pol region. The proposed new subgenus *Ritunrivirus* is based on a statistically supported single lineage of 22 distinct species in the genus *Waikavirus* defined by the amino acid sequence of the combined three CPs and the conserved Pro-Pol region. The proposed new subgenus *Actinidivirus* is based on a statistically supported single lineage of 16 distinct species in the genus *Waikavirus* defined by the

amino acid sequence of the combined 3 CPs and conserved Pro-Pol region. The creation of the proposed new 34 species is based on <75% amino acid sequence identity in the CP(s) and/or <80% amino acid sequence identity in the conserved Pro-Pol region (from the protease CG motif to the polymerase GDD motif) compared with members of classified species of the family *Secoviridae*.

**Submitted:** 10/06/24; **Revised**: 11/10/24

Table 13. Secoviridae, 37 new taxa\*

| Operation | Rank     | New taxon name             | Virus name                            | Exemplar           |
|-----------|----------|----------------------------|---------------------------------------|--------------------|
| New taxon | Genus    | Mersevirus                 |                                       |                    |
| New taxon | Species  | Mersevirus boehmeriae      | Boehmeria nivea secovirus             | BK061322; BK061323 |
| New taxon | Species  | Mersevirus jujubae         | jujube-associated virus 1             | MT375548; MT375547 |
| New taxon | Species  | Mersevirus mercurialis     | Mercurialis secovirus 1               | OR544055; OR544056 |
| New taxon | Species  | Mersevirus paris           | Paris polyphylla secovirus 2          | BK061330; BK061331 |
| New taxon | Subgenus | Actinidivirus              |                                       |                    |
| New taxon | Species  | Waikavirus betacamelliae   | Camellia virus B                      | BK062984           |
| New taxon | Species  | Waikavirus carotae         | carrot psyllid-borne associated virus | OM801008           |
| New taxon | Species  | Waikavirus celtis          | hackberry virus A                     | OP533794           |
| New taxon | Species  | Waikavirus hirtae          | Ficus hirta waikavirus                | BK062987           |
| New taxon | Species  | Waikavirus juglandis       | Juglans nigra waikavirus              | BK062989           |
| New taxon | Species  | Waikavirus pittospori      | Pittosporum tobira virus              | OR659471           |
| New taxon | Species  | Waikavirus populi          | Populus alba waikavirus               | BK062992           |
| New taxon | Species  | Waikavirus querci          | Quercus robur waikavirus              | BK062996           |
| New taxon | Species  | Waikavirus trifoccidentale | Trifolium occidentale waikavirus      | BK063000           |
| New taxon | Subgenus | Ritunrivirus               |                                       |                    |
| New taxon | Species  | Waikavirus ajugae          | Ajuga reptans waikavirus              | BK062980           |
| New taxon | Species  | Waikavirus anacycli        | Anacyclus depressus waikavirus        | BK062979           |
| New taxon | Species  | Waikavirus eleocharis      | Eleocharis dulcis waikavirus          | BK062986           |
| New taxon | Species  | Waikavirus ligustici       | Ligusticum chuanxiong waikavirus      | BK062990           |
| New taxon | Species  | Waikavirus mertensiae      | Mertensia paniculata waikavirus       | BK062991           |
| New taxon | Species  | Waikavirus pedicularis     | Pedicularis rex waikavirus            | BK062993           |
| New taxon | Species  | Waikavirus primulae        | Primula vulgaris waikavirus           | BK062995           |
| New taxon | Species  | Waikavirus ranunculi       | Ranunculus cantoniensis waikavirus    | BK062997           |
| New taxon | Species  | Waikavirus thapsiae        | Thapsia villosa waikavirus            | BK063001           |
| New taxon | Species  | Waikavirus thymi           | Thymus vulgaris waikavirus            | BK062999           |
| New taxon | Species  | Waikavirus violae          | Viola inconspicua waikavirus          | BK063002           |
| New taxon | Species  | Fabavirus betavitis        | grapevine secovirus                   | OR947508; OR947509 |
| New taxon | Species  | Fabavirus cirsii           | Cirsium virus A                       | OP794357; OP794358 |
| New taxon | Species  | Nepovirus betaparis        | Paris polyphylla secovirus 1          | BK061328; BK061329 |
| New taxon | Species  | Nepovirus mirae            | Prunus mira virus A                   | BK064709; BK064710 |
| New taxon | Species  | Sadwavirus cattleyae       | Cattleya purple ringspot virus        | OR439368; OR439369 |

Continued

Table 13. Continued

| Operation | Rank    | New taxon name           | Virus name                   | Exemplar           |
|-----------|---------|--------------------------|------------------------------|--------------------|
| New taxon | Species | Sadwavirus gymnemae      | Gymnema sylvestre virus 1    | BK062888; BK062889 |
| New taxon | Species | Sadwavirus chysanthemi   | chrysanthemum sadwavirus     | OR413567; OR413568 |
| New taxon | Species | Torradovirus arctii      | burdock mosaic virus         | OQ087134; OQ087135 |
| New taxon | Species | Torradovirus nanorugosum | potato rugose stunting virus | ON871623; ON871624 |
| New taxon | Species | Torradovirus physalis    | Physalis torrado virus       | MZ357183; MZ357184 |

Table 14. Secoviridae, 13 move taxa\*

| Operation  | Rank    | Taxon name             | Old parent taxon | New parent taxon       |
|------------|---------|------------------------|------------------|------------------------|
| Move taxon | Species | Waikavirus actinidiae  | Waikavirus       | Subgenus Actinidivirus |
| Move taxon | Species | Waikavirus camelliae   | Waikavirus       | Subgenus Actinidivirus |
| Move taxon | Species | Waikavirus diospyri    | Waikavirus       | Subgenus Actinidivirus |
| Move taxon | Species | Waikavirus liegense    | Waikavirus       | Subgenus Actinidivirus |
| Move taxon | Species | Waikavirus rhododendri | Waikavirus       | Subgenus Actinidivirus |
| Move taxon | Species | Waikavirus brassicae   | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus campanulae  | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus lactucae    | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus oryzae      | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus ribesnigri  | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus rosae       | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus trifolii    | Waikavirus       | Subgenus Ritunrivirus  |
| Move taxon | Species | Waikavirus zeae        | Waikavirus       | Subgenus Ritunrivirus  |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.013P.Secoviridae\_1ng\_2nsg\_34nsp.zip.

# 2024.014P.Rhabdoviridae\_2nsp

**Title:** Create one new species in the genus *Alphanucleorhabdovirus* and one species in the genus *Betanucleorhabdovirus*, subfamily *Betarhabdovirinae* (*Mononegavirales: Rhabdoviridae*)

Authors: Bejerman N. (bejerman.nicolas@inta.gob.ar), Debat H., Dietzgen R., Freitas-Astua J., Kondo H., Ramos-Gonzalez P., Whitfield A., Walker P.

# **Summary**

# Taxonomic rank(s) affected

Species

#### Description of current taxonomy

Viruses classified in the genera *Alphanucleorhabdovirus* and *Betanucleorhabdovirus* infect a wide range of plants, and the assignment of viruses to these genera is based on the placement of the viruses on maximum likelihood trees inferred from complete L protein sequences.

#### Proposed taxonomic change(s)

Add one (1) new species to the genus *Alphanucleorhabdovirus* (*Alphanucleorhabdovirus babaci*) and one (1) new species in the genus *Betanucleorhabdovirus* (*Betanucleorhabdovirus paridis*).

#### Justification

Two novel rhabdoviruses were identified in babaco [17] and  $Paris\ polyphylla\ [18]$ . The characterization of both viruses showed that the babaco-associated virus should be classified as a novel species within the genus Alphanucleorhabdovirus, while the Pappen Pap

**Submitted:** 10/06/24

Table 15. Rhabdoviridae. 2 new taxa\*

| Operation | Rank    | New taxon name                   | Virus name                         | Exemplar |
|-----------|---------|----------------------------------|------------------------------------|----------|
| New taxon | Species | Alphanucleorhabdovirus<br>babaci | babaco nucleorhabdovirus 1         | OQ256237 |
| New taxon | Species | Betanucleorhabdovirus<br>paridis | Paris yunnanensis<br>rhabdovirus 1 | OL439478 |

<sup>\*</sup>Source/full text; https://ictv.global/ictv/proposals/2024.014P.Rhabdoviridae\_2nsp.zip

# 2024.015P.Rhabdoviridae\_Cytorhabdovirus\_splitgen

**Title:** Abolish one genus and create three new genera to include 98 new species in the subfamily *Betarhabdovirinae* (*Mononegavirales*: *Rhabdoviridae*)

Authors: Bejerman N. (bejerman.nicolas@inta.gob.ar), Debat H., Dietzgen R., Freitas-Astua J., Kondo H., Ramos-Gonzalez P., Whitfield A., Walker P.

#### **Summary**

# Taxonomic rank(s) affected

Genus and species

# Description of current taxonomy

Viruses classified in the genus *Cytorhabdovirus* infect a wide range of plants, and the assignment of viruses to this genus is based on the placement of the viruses on maximum likelihood trees inferred from complete L protein sequences.

#### Proposed taxonomic change(s)

Abolish the genus *Cytorhabdovirus* and add three (3) new genera (*Alphacytorhabdovirus*, *Betacytorhabdovirus* and *Gammacytorhabdovirus*), including 98 new species in the subfamily *Betarhabdovirinae* (*Mononegavirales*: *Rhabdoviridae*). Abolish four (4) *Cytorhabdovirus* species and reassign the remaining species to the new genera.

#### **Justification**

Recently, 98 new putative cytorhabdoviruses were discovered. The phylogenetic relationships of the now significantly expanded number of known cytorhabdoviruses provide support for splitting the genus *Cytorhabdovirus* into three genera that represent distinct evolutionary lineages to be named *Alphacytorhabdovirus*, *Betacytorhabdovirus* and *Gammacytorhabdovirus*. Also, it is proposed that four *Cytorhabdovirus* species be abolished based on the lack of sequence data for them.

Submitted: 10/06/24; Revised: 03/10/24

Table 16. Rhabdoviridae, 101 new taxa\*. Table too large, see supplementary information sheet supp\_info\_tab\_16

Table 17. Rhabdoviridae, 51 move; rename taxa\*. Table too large, see supplementary information sheet supp\_info\_tab\_17

Table 18. Rhabdoviridae, 4 abolish taxa\*

| Operation     | Rank    | Abolished taxon name      |
|---------------|---------|---------------------------|
| Abolish taxon | Species | Cytorhabdovirus brassicae |
| Abolish taxon | Species | Cytorhabdovirus festucae  |
| Abolish taxon | Species | Cytorhabdovirus sonchi    |
| Abolish taxon | Species | Cytorhabdovirus tritici   |

 $<sup>*</sup>Source/full\ text:\ https://ictv.global/ictv/proposals/2024.015P.Rhabdoviridae\_Cytorhabdovirus\_splitgen.zip.$ 

# 2024.016P.Rhabdoviridae\_1ngen\_5nsp

**Title:** Create one new genus to include five new species in the subfamily *Betarhabdovirinae* (*Mononegavirales: Rhabdoviridae*)

**Authors:** Bejerman N. (bejerman.nicolas@inta.gob.ar), Debat H., Dietzgen R.G., Freitas-Astua J., Kondo H., Ramos-Gonzalez P., Whitfield A., Walker P.

#### **Summary**

### Taxonomic rank(s) affected

Genus and species

#### **Description of current taxonomy**

Almost all viruses classified in the subfamily *Betarhabdovirinae*, family *Rhabdoviridae*, are unsegmented, but plant-associated rhabdoviruses with bi-segmented genomes have also been identified and included in the genera *Varicosavirus* and *Dichorhavirus* within the subfamily *Betarhabdovirinae*. The assignment of viruses to these genera is based on the placement of the viruses on maximum likelihood trees inferred from complete L protein sequences.

#### Proposed taxonomic change(s)

Add one (1) new genus to include five (5) new species in the subfamily *Betarhabdovirinae*, family *Rhabdoviridae*. These new species (*Trirhavirus alni*, *Trirhavirus chrysanthemi*, *Trirhavirus erysimi*, *Trirhavirus medicagonis* and *Trirhavirus picridis*) are placed in a new genus named *Trirhavirus*.

#### **Justification**

Five novel rhabdoviruses were identified in *Alnus rubra*, *Chrysanthemum morifolium*, *Erysimum nevadense*, *Medicago sativa* and *Picris echioides* [19]. Unexpectedly, these five viruses have tri-segmented genomes, which represent the first tri-segmented genomes among rhabdoviruses. The characterization of these five viruses showed they should be classified as members of novel species within a novel genus within the subfamily *Betarhabdovirinae*, family *Rhabdoviridae*, for which the name *Trirhavirus* is proposed [19].

Submitted: 10/06/24

Table 19. Rhabdoviridae. 6 new taxa\*

| Operation | Rank    | New taxon name           | Virus name                  | Exemplar                     |
|-----------|---------|--------------------------|-----------------------------|------------------------------|
| New taxon | Genus   | Trirhavirus              |                             |                              |
| New taxon | Species | Trirhavirus alni         | Alnus trirhavirus 1         | BK064247; BK064248; BK064249 |
| New taxon | Species | Trirhavirus chrysanthemi | chrysanthemum trirhavirus 1 | BK064250; BK064251; BK064252 |
| New taxon | Species | Trirhavirus erysimi      | Erysimum trirhavirus 1      | BK064253; BK064254; BK064255 |
| New taxon | Species | Trirhavirus medicagonis  | Medicago trirhavirus        | BK064256; BK064257; BK064258 |
| New taxon | Species | Trirhavirus picridis     | Picris trirhavirus 1        | BK064259; BK064260; BK064261 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.016P.Rhabdoviridae\_1ngen\_5nsp.zip.

# 2024.017P.Tombusviridae abolishsp

Title: Abolish five unassigned species in the family Tombusviridae

Authors: Scheets K. (kay.scheets@okstate.edu), Hernández C., Jordan R., Miller W.A., Prigigallo M.I., Rubino L.

# **Summary**

#### Taxonomic rank(s) affected

Species in the family Tombusviridae

# **Description of current taxonomy**

Ahlum waterborne virus, Bean mild mosaic virus, Chenopodium necrosis virus, Cucumber soil-borne virus and Weddel waterborne virus are currently classified as unassigned species in the family *Tombusviridae*.

#### Proposed taxonomic change(s)

Abolish five (5) species.

# Justification

The proposed abolishment of these species is based on the lack of sequence data for them.

**Submitted:** 21/06/24

Table 20. Tombusviridae, 5 abolish taxa\*

| Operation     | Rank    | Abolished taxon name   |
|---------------|---------|------------------------|
| Abolish taxon | Species | Ahlum waterborne virus |
| Abolish taxon | Species | Bean mild mosaic virus |

Continued

Table 20. Continued

| Operation     | Rank    | Abolished taxon name       |
|---------------|---------|----------------------------|
| Abolish taxon | Species | Chenopodium necrosis virus |
| Abolish taxon | Species | Cucumber soil-borne virus  |
| Abolish taxon | Species | Weddel waterborne virus    |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.017P.Tombusviridae\_abolishsp.zip.

# 2024.018P.Tombusviridae\_1nsp

**Title:** Create one new species in the genus *Machlomovirus* (*Tolivirales*: *Tombusviridae*)

Authors: Maclot F., Massart S. (sebastien.massart@uliege.be)

### **Summary**

#### Taxonomic rank(s) affected

Genus Machlomovirus (Tolivirales: Tombusviridae)

#### Description of current taxonomy

One virus species, Machlomovirus zeae, is currently described within the genus Machlomovirus.

#### Proposed taxonomic change(s)

Add a second species (*Machlomovirus liegense*) to the genus *Machlomovirus* to accommodate a recently identified virus in the wild grass common bent (*Agrostis capillaris*), tentatively named Poaceae Liege machlomovirus.

# Justification

Analysis of Poaceae Liege machlomovirus genomic structure and phylogenetic analyses of its complete sequence and specific genes (polymerase and coat protein) placed this virus as a member of a novel species in the genus *Machlomovirus*.

Submitted: 21/06/24

Table 21. Tombusviridae, 1 new taxon\*

| Operation | Rank    | New taxon name         | Virus name                  | Exemplar |
|-----------|---------|------------------------|-----------------------------|----------|
| New taxon | Species | Machlomovirus liegense | Poaceae Liege machlomovirus | ON137711 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.018P.Tombusviridae\_1nsp.zip

# 2024.019P.Closteroviridae\_1nsp

Title: Create one new species in the genus Velarivirus (order Martellivirales, family Closteroviridae)

Authors: Fontdevila N., Massart S. (sebastien.massart@uliege.be)

#### Summary

# Taxonomic rank(s) affected

Genus Velarivirus (order Martellivirales, family Closteroviridae)

#### Description of current taxonomy

The family *Closteroviridae* comprises plant viruses with long, filamentous particles (650–2,200 nm in length) and large positive-sense RNA genomes (mono-, bi- or tripartite). There are 57 recognized species in the family, classified in 1 of the 7 existing genera (*Ampelovirus*, *Bluvavirus*, *Closterovirus*, *Crinivirus*, *Menthavirus*, *Olivavirus* and *Velarivirus*). Within the genus *Velarivirus*, there are currently eight recognized species.

#### Proposed taxonomic change(s)

The authors propose adding a ninth species in the genus *Velarivirus*, named *Velarivirus gembloutense*, to classify the recently identified virus Pyrus virus A in pear trees (*Pyrus communis* L.).

#### Justification

Analysis of the genomic structure of Pyrus virus A and subsequent phylogenetic analyses of the specific HSP70h gene placed this virus as a member of a novel species of the genus *Velarivirus* within the family *Closteroviridae*.

Submitted: 21/06/24; Revised: 07/10/24

Table 22. Closteroviridae, 1 new taxon\*

| Operation | Rank    | New taxon name           | Virus name    | Exemplar |
|-----------|---------|--------------------------|---------------|----------|
| New taxon | Species | Velarivirus gembloutense | Pyrus virus A | OR887735 |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.019P.Closteroviridae\_1nsp.zip.

# 2024.021P.Riboviria\_1nord

**Title:** Create one new unassigned order in realm *Riboviria*, and three new families for four currently unassigned genera of plant satellite viruses

Authors: Krupovic M. (mart.krupovic@pasteur.fr), Fischer M.G., Kuhn J.H.

#### **Summary**

#### Taxonomic rank(s) affected

Species, genus, family

# Description of current taxonomy

Riboviria: unassigned family Sarthroviridae and genera Albetovirus, Aumaivirus, Papanivirus and Virtovirus.

# Proposed taxonomic change(s)

Riboviria: Tombendovirales to include family Sarthroviridae and two (2) new families, Pamosaviridae (Papanivirus) and Tomosaviridae (Virtovirus); Riboviria: Tonesaviridae (Albetovirus, Aumaivirus); renaming of all species in the four genera to fulfil the ICTV's binomial naming mandate.

#### **Justification**

Structural comparison of the satellite virus capsid proteins indicates that these viruses are not monophyletic and form two distinct assemblages.

**Submitted:** 21/06/24

Table 23. Riboviria, 6 rename taxa\*

| Operation    | Rank    | New taxon name          | Previous taxon name   |
|--------------|---------|-------------------------|-----------------------|
| Rename taxon | Species | Papanivirus panici      | Panicum papanivirus 1 |
| Rename taxon | Species | Virtovirus tabaci       | Tobacco virtovirus 1  |
| Rename taxon | Species | Albetovirus alphatabaci | Tobacco albetovirus 1 |
| Rename taxon | Species | Albetovirus betatabaci  | Tobacco albetovirus 2 |
| Rename taxon | Species | Albetovirus gammatabaci | Tobacco albetovirus 3 |
| Rename taxon | Species | Aumaivirus maidis       | Maize aumaivirus 1    |

Table 24. Riboviria, 5 move taxa\*

| Operation  | Rank   | Taxon name     | New parent taxon | Old parent taxon |
|------------|--------|----------------|------------------|------------------|
| Move taxon | Family | Sarthroviridae | Tombendovirales  | Riboviria        |
| Move taxon | Genus  | Papanivirus    | Pamosaviridae    | Riboviria        |
| Move taxon | Genus  | Virtovirus     | Tomosaviridae    | Riboviria        |
| Move taxon | Genus  | Albetovirus    | Tonesaviridae    | Riboviria        |
| Move taxon | Genus  | Aumaivirus     | Tonesaviridae    | Riboviria        |

Table 25. Riboviria, 4 new taxa\*

| Operation | Rank   | New taxon name  |
|-----------|--------|-----------------|
| New taxon | Order  | Tombendovirales |
| New taxon | Family | Pamosaviridae   |

Continued

Table 25. Continued

| Operation | Rank   | New taxon name |
|-----------|--------|----------------|
| New taxon | Family | Tomosaviridae  |
| New taxon | Family | Tonesaviridae  |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.021P.Riboviria\_1nord.zip.

# 2024.022P.Betaflexiviridae\_abolishsp

**Title:** Abolish one unassigned species in the family *Betaflexiviridae* 

Authors: Nagata T. (tatsuya@unb.br), Blouin A., Candresse T., Cao M., Cho W.K., Constable F., Sabanadzovic S., Saldarelli P., Tzanetakis I., Villamor D.

### **Summary**

#### Taxonomic rank(s) affected

Species in the family Betaflexiviridae

# Description of current taxonomy

Banana virus X is currently classified as an unassigned species in the family Betaflexiviridae.

#### Proposed taxonomic change(s)

Abolish one (1) species.

#### **Justification**

The proposed abolishment of this betaflexivirid species is based on the lack of Rep sequence data.

**Submitted:** 30/06/24

Table 26. Betaflexiviridae. 1 abolish taxon\*

| Operation     | Rank    | Abolished taxon name |
|---------------|---------|----------------------|
| Abolish taxon | Species | Banana virus X       |

<sup>\*</sup>Source/full text: https://ictv.global/ictv/proposals/2024.022P.Betaflexiviridae\_abolishsp.zip.

# 2024.023P.Solemoviridae\_rename\_sp

Title: Rename two species in the genus Sobemovirus (family Solemoviridae)

Authors: Somera M. (merike.somera@taltech.ee), Fargette D., Filardo F., Ghafari M., Hebrard E., Sarmiento C., Thomas J.E.

#### **Summary**

# Taxonomic rank(s) affected

**Species** 

#### Description of current taxonomy

Riboviria: Orthornavirae: Pisuviricota: Pisoniviricetes: Sobelivirales: Solemoviridae: Sobemovirus: Cocksfoot mottle virus. Riboviria: Orthornavirae: Pisuviricota: Pisoniviricetes: Sobelivirales: Solemoviridae: Sobemovirus: Sobemovirus smamv.

# Proposed taxonomic change(s)

Renaming of Cocksfoot mottle virus and Sobemovirus smamv.

#### **Justification**

Renaming of *Cocksfoot mottle virus* to fulfil the ICTV's binomial naming mandate. Renaming of *Sobemovirus smamv* for consistency with other species in the family *Solemoviridae*.

Submitted: 28/06/24; Revised: 07/10/24

**Table 27.** Solemoviridae, 2 rename taxa\*

| Operation    | Rank    | New taxon name    | Previous taxon name    |
|--------------|---------|-------------------|------------------------|
| Rename taxon | Species | Sobemovirus CFMV  | Cocksfoot mottle virus |
| Rename taxon | Species | Sobemovirus SMAMV | Sobemovirus smamv      |

 $<sup>*</sup>Source/full\ text: https://ictv.global/ictv/proposals/2024.023P. Solemoviridae\_rename\_sp.zip. A solution of the control of$ 

# 2024.024P.Tymoviridae\_abolish\_sp

Title: Abolish two unassigned species in the family Tymoviridae

Authors: Hammond R. (rose.hammond@usda.gov), Abrahamian P., Bejerman N., Mollov D., Nagata T., Sabanadzovic S.

#### **Summary**

#### Taxonomic rank(s) affected

Species in the family *Tymoviridae* 

#### Description of current taxonomy

Bombyx mori latent virus and Poinsettia mosaic virus are currently classified as unassigned species in the family Tymoviridae.

# Proposed taxonomic change(s)

Abolish two (2) species.

#### **Justification**

Bombyx mori latent virus and Poinsettia mosaic virus are unassigned species in the family *Tymoviridae* which cannot be assigned to a genus because of the lack of genome sequence. Therefore, it is not possible to comply with the ICTV mandate of a binomial format for virus species and it is proposed that these tymovirid species be abolished.

**Submitted:** 14/07/24

Table 28. Tymoviridae, 2 abolish taxa\*

| Operation     | Rank    | Abolished taxon name     |
|---------------|---------|--------------------------|
| Abolish taxon | Species | Bombyx mori latent virus |
| Abolish taxon | Species | Poinsettia mosaic virus  |

\*Source/full text: https://ictv.global/ictv/proposals/2024.024P.Tymoviridae\_abolish\_sp.zip.

Author affiliations: 1 Istituto per la Protezione Sostenibile delle Piante, CNR, Bari, Italy; 2 USDA-ARS, BARC, National Germplasm Resources Laboratory, Beltsville, MD, USA; <sup>3</sup>Liaoning Key Laboratory of Urban Integrated Pest Management and Ecological Security, Shenyang University, Dadong, Shenyang, Liaoning, PR China; "Centro de Edafología y Biología Aplicada del Segura-CSIC, Murcia, Spain; Department of Molecular and Structural Biochemistry, North Carolina State University, Raleigh, NC, USA; éUnidad de Fitopatología y Modelización Agrícola (UFYMA) INTA-CONICET, Buenos Aires, Argentina; Plant Protection Department, Agroscope, Nyon, Switzerland; 8UMR 1332 Biologie du Fruit et Pathologie, University of Bordeaux, INRAE, Domaine de la Grande Ferrade, 71 Avenue Edouard Bourlaux, CS 20032, 33882 Villenave d'Ornon Cedex, France; Margarita Salas Center for Biological Research (CIB-CSIC) Spanish Council for Scientific Research (CSIC), Madrid, Spain; 10 National Citrus Engineering and Technology Research Center, Integrative Science Center of Germplasm Creation in Western China (CHONGQING) Science City, Citrus Research Institute, Southwest University, Beibei, Chongging, PR China; 11 Department of Plant Sciences, University of Cambridge, Cambridge, UK; 12 Agriculture and Life Sciences Research Institute, Kangwon National University, Chuncheon, Republic of Korea; 13 Agriculture Victoria Research, Department of Energy, Environment and Climate Action and School of Applied Systems Biology, La Trobe University, Bundoora, Australia; 14University of Delhi South Campus, Benito Juarez Road, New Delhi, India; 15Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, St Lucia QLD, Australia; 16CIHEAM, Istituto Agronomico Mediterraneo of Bari, Via Ceglie 9, Valenzano (BA), Italy; 17Virus South Data, CDV 230980, La Mure, France; 18Queensland Department of Primary Industries, Brisbane, Australia; 19 Max Planck Institute for Marine Microbiology, Bremen, Germany; 20 Fera Science Ltd (Fera), York Biotech Campus, Sand Hutton, York, UK; <sup>21</sup>Embrapa Cassava and Fruits, Brazilian Agricultural Research Corporation, Cruz das Almas, Brazil; <sup>22</sup>Plant Pathology, Cornell University, Geneva, NY, USA; <sup>23</sup>Department of Biology, University of Oxford, Oxford, UK; <sup>24</sup>Swedish University of Agriculture, Almas Allé 5, Uppsala, Sweden; <sup>25</sup>USDA-ARS, USNA, Floral and Nursery Plants Research Unit, Beltsville, MD, USA; <sup>26</sup>USDA-ARS, BARC, Molecular Plant Pathology Laboratory, Beltsville, MD, USA; <sup>27</sup>Institute of Plant Protection-NRI, ul. Wł. Węgorka 20, Poznań, Poland; <sup>28</sup>PHIM Plant Health Institute, University of Montpellier, INRAE, CIRAD, IRD, Institute Agro, Montpellier, France; 29 Instituto de Biología Molecular y Celular de Plantas (IBMCP), Universitat Politècnica de Valencia-CSIC, Valencia, Spain; 30Institut Français de la Vigne et du Vin, Le Grau du Roi, France; 31Vali-e-Asr University of Rafsanjan, Department of Plant Protection, Rafsanjan, Iran; 32Retired from John Innes Centre, Norwich, UK; 33Embrapa Hortaliças, Brasília, DF, Brazil; 34Institute of Plant Science and Resources, Okayama University, Kurashiki, Japan; 35International Potato Center (CIP), Lima, Peru; 36Institut Pasteur, Université Paris Cité, CNRS UMR6047, Archaeal Virology Unit, Paris, France; 37Institute for Plant Protection, NARO, 2-1-18, Kannondai, Tsukuba, Japan; 38Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, MD, USA; 39 Department of Biological Sciences, University of Toledo, Toledo, OH, USA; 40CIRAD, UMR PVBMT, Saint Pierre, La Réunion, France; 41State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan Agricultural University, Kunming, PR China; 42Institute of Plant Virology, Ningbo University, Ningbo, PR China; 43 Instituto de Patología Vegetal (IPAVE), INTA, Unidad de Fitopatología y Modelización Agrícola (UFYMA) INTA-CONICET, Córdoba, Argentina; 44Centre for Research in Agricultural Genomics, CRAG (CSIC-IRTA-UAB-UB), Barcelona, Spain; 45Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland; "

Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa; 47 Plant Pathology Laboratory, TERRA Gembloux Agro-Bio Tech, University of Liege, Liege, Belgium; 48 Department of Plant Pathology, Entomology and Microbiology, Iowa State University, Ames, IA, USA; 49Department of Plant Protection, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran; 50 USDA-APHIS, Plant Protection and Quarantine, 4700 River Road, Riverdale, MD, USA; 51 CIRAD, AGAP Institut; AGAP Institut, University of Montpellier; CIRAD, INRAE, Institut Agro, 34398 Montpellier, France; 52 Instituto de Ciências Biológicas, Universidade de Brasília, Brasília, Brazil; 53 nstituto de Hortofruticultura Subtropical y Mediterránea "La Mayora" (IHSM-UMA-CSIC), Consejo Superior de Investigaciones Científicas, Algarrobo-Costa, Málaga, Spain; 54Utsunomiya University, Utsunomiya, Japan; 55Oklahoma State University, Institute for Biosecurity & Microbial Forensics, 127 NRC Stillwater, OK, USA; 56Saga University, Saga, Japan; 57Department of Plant Pathology, Washington State University, Pullman, WA, USA; 58 Institute of Plant Molecular Biology, České Budějovice, Czech Republic; 59 Applied Molecular Biology Laboratory, Instituto Biológico de São Paulo, São Paulo, Brazil; 🗠 Embrapa Recursos Genéticos e Biotecnologia, Brasília, Brazil; 🕯 Julius Kühn Institute, Federal Research Centre for Cultivated Plants,

Institute for Epidemiology and Pathogen Diagnostics, Braunschweig, Germany; 62CIRAD, UMR PHIM, Montpellier, France; 63Department of Agricultural Science and Plant Protection, Mississippi State University, Mississippi State, MS, USA; 64Department of Cell Biology and Genetics, Faculty of Science, Palacký University Olomouc, Olomouc, Czech Republic; 65Summerland Research and Development Centre, Agriculture and Agri-Food Canada, Summerland, Canada; <sup>66</sup>Department of Chemistry and Biotechnology, Tallinn University of Technology, Tallinn, Estonia; <sup>67</sup>Strategic Planning Headquarters, NARO, 3-1-1 Kannondai, Tsukuba, Japan; 68 Department of Plant Pathology, Ecology and Evolution, Oklahoma State University, Stillwater, OK, USA; 69 Molecular Plant Pathology, University of Amsterdam, Amsterdam, Netherlands; 70 Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent, UK; 7¹Kochi Agricultural Research Center, Nankoku, Japan; 7²Currently unaffiliated; 7³CIRAD, UMR PVBMT & UMR PVBMT, Université de la Réunion, F-97410 Saint Pierre de La Réunion, France; 74Plant Health and Environment Laboratory, 231 Morrin Rd, St. Johns, Auckland, New Zealand; 75 Council for Agricultural Research and Economics, Research Centre for Plant Protection and Certification, Rome, Italy; <sup>76</sup>Department of Entomology and Plant Pathology, Division of Agriculture, University of Arkansas System, Fayetteville, AR, USA; <sup>77</sup>INRAE, UR ASTRO, Domaine de Duclos, Petit Bourg, France; 78 Wageningen University and Research, Wageningen, Netherlands; 79 The Biodesign Center for Fundamental and Applied Microbiomics, Center for Evolution and Medicine, School of Life Sciences, Arizona State University, Tempe, AZ, USA; 80 Rijk Zwaan Breeding B.V., De Lier, Netherlands; 81Humboldt-Universität zu Berlin, Thaer-Institute of Agricultural and Horticultural Sciences, Unter den Linden 6, Berlin, Germany; 82The University of Queensland, St Lucia, Australia; 83Dienstleistungszentrum Ländlicher Raum Rheinpfalz, Breitenweg 71, Neustadt an der Weinstrasse, Germany; 84North Carolina State University, Raleigh, NC, USA; 85Food Futures Institute, Murdoch University, 90 South Street, Perth, Australia; 86 Dep. de Fitopatologia/BIOAGRO, Universidade Federal de Viçosa, Viçosa, Brazil.

Keywords: Actinidivirus; Ahlum waterborne virus; Albetovirus alphatabaci; Albetovirus betatabaci; Albetovirus gammatabaci; Allexivirus rehmanniae; Alphacytorhabdovirus; Alphacytorhabdovirus actinidiae; Alphacytorhabdovirus allii; Alphacytorhabdovirus alphacytorhabdovirus alphachysanthemi; Alphacytorhabdovirus alphafragariae; Alphacytorhabdovirus alphamedicagonis; Alphacytorhabdovirus alphapogostemi; Alphacytorhabdovirus alphaprimulae; Alphacytorhabdovirus alpharubi; Alphacytorhabdovirus alpha Alphacytorhabdovirus arctii; Alphacytorhabdovirus asclepiadis; Alphacytorhabdovirus baccharis; Alphacytorhabdovirus baccharis betaartemisiae; Alphacytorhabdovirus betachrysanthemi; Alphacytorhabdovirus betafragariae; Alphacytorhabdovirus betamedicagonis; Alphacytorhabdovirus betapogostemi; Alphacytorhabdovirus betaprimulae; Alphacytorhabdovirus betarubi; Alphacytorhabdovirus betarifolii; Alphacytorhabdovirus betawuhaninsectum; Alphacytorhabdovirus brassicicolae; Alphacytorhabdovirus cardaminis; Alphacytorhabdovirus chelidonii; Alphacytorhabdovirus cnidii; Alphacytorhabdovirus conopholis; Alphacytorhabdovirus coriandri; Alphacytorhabdovirus cynarae; Alphacytorhabdovirus daphnis; Alphacytorhabdovirus deltapogostemi; Alphacytorhabdovirus euphorbiae; Alphacytorhabdovirus fici; Alphacytorhabdovirus fragariarugosus; Alphacytorhabdovirus qammaartemisiae; Alphacytorhabdovirus qammapoqostemi; Alphacytorhabdovirus qammawuhaninsectum; Alphacytorhabdovirus gei; Alphacytorhabdovirus glehniae; Alphacytorhabdovirus hederae; Alphacytorhabdovirus hyptisis; Alphacytorhabdovirus ilicis; Alphacytorhabdovirus kenyatuberosum; Alphacytorhabdovirus lactucanecante; Alphacytorhabdovirus lactucamaculante; Alphacytorhabdovirus lycopersici; Alphacytorhabdovirus menthae; Alphacytorhabdovirus morindae; Alphacytorhabdovirus nymphaeae; Alphacytorhabdovirus ocimi; Alphacytorhabdovirus paludis; Alphacytorhabdovirus pastinacae; Alphacytorhabdovirus pelargonii; Alphacytorhabdovirus persimmon; Alphacytorhabdovirus phyllostachysis; Alphacytorhabdovirus pinelliae; Alphacytorhabdovirus plumagonis; Alphacytorhabdovirus querci; Alphacytorhabdovirus ribes; Alphacytorhabdovirus rosae; Alphacytorhabdovirus sambuci; Alphacytorhabdovirus scutellariae; Alphacytorhabdovirus taraxaci; Alphacytorhabdovirus tolmieae; Alphacytorhabdovirus trichosantheir, Alphacytorhabdovirus tritici; Alphacytorhabdovirus utriculariae; Alphacytorhabdovirus wurfbainiae; Alphacytorhabdovirus zeae; Alphanucleorhabdovirus babaci; Aumaivirus maidis; Badnavirus fatsiae; Badnavirus tetainflatheobromae; Badnavirus ziziphi; Banana virus X; Bean mild mosaic virus; Begomovirus alceacrispi; Begomovirus caboniensis; Begomovirus cajani; Begomovirus chuxiongense; Begomovirus flavintervenae; Begomovirus galii; Begomovirus hortuscrotoni; Begomovirus hyptidis; Begomovirus jatrophagunturense; Begomovirus muntiflavi; Begomovirus myanmarense; Begomovirus puerense; Begomovirus pyrenacanthae; Begomovirus sidaflavitessellati; Begomovirus solanumaureusreti; Begomovirus solanumdistorsionis; Begomovirus solanumflavusreti; Begomovirus whitaniae; Betacytorhabdovirus; Betacytorhabdovirus alphabetulae; Betacytorhabdovirus alphacucurbitae; Betacytorhabdovirus alpharosae; Betacytorhabdovirus alphaturbaesoli; Betacytorhabdovirus alphazanthoxyli; Betacytorhabdovirus anthurii; Betacytorhabdovirus aristolochiae; Betacytorhabdovirus artemisiae; Betacytorhabdovirus begoniae; Betacytorhabdovirus bemisiae; Betacytorhabdovirus betabetulae; Betacytorhabdovirus betacucurbitae; Betacytorhabdovirus betarosae; Betacytorhabdovirus betaturbaesoli; Betacytorhabdovirus betazanthoxyli; Betacytorhabdovirus bouteloae; Betacytorhabdovirus broussonetiae; Betacytorhabdovirus caricae; Betacytorhabdovirus chrysanthemi; Betacytorhabdovirus colocasiae; Betacytorhabdovirus coryli; Betacytorhabdovirus cypripedii; Betacytorhabdovirus dryobalanopis; Betacytorhabdovirus durionis; Betacytorhabdovirus flaviyerbamate; Betacytorhabdovirus gammazanthoxyli; Betacytorhabdovirus gleditsiae; Betacytorhabdovirus glycinis; Betacytorhabdovirus glycyrrhizae; Betacytorhabdovirus gramineae: Betacytorhabdovirus hepaticae: Betacytorhabdovirus hordei: Betacytorhabdovirus howeae: Betacytorhabdovirus ipomoeae; Betacytorhabdovirus ixeris; Betacytorhabdovirus justiciae; Betacytorhabdovirus kobresiae; Betacytorhabdovirus leucadendri; . Betacytorhabdovirus lycii; Betacytorhabdovirus mangonis; Betacytorhabdovirus maydis; Betacytorhabdovirus maysflavostriatis; Betacytorhabdovirus mori; Betacytorhabdovirus nitrariae; Betacytorhabdovirus oryzae; Betacytorhabdovirus panici; Betacytorhabdovirus passiflorae; Betacytorhabdovirus pentaphragmae; Betacytorhabdovirus phellodendri; Betacytorhabdovirus populi; Betacytorhabdovirus puerariae; Betacytorhabdovirus rudbeckiae; Betacytorhabdovirus schiedeae; Betacytorhabdovirus sesami, Betacytorhabdovirus sophorae; Betacytorhabdovirus tagetis; Betacytorhabdovirus tiliae; Betacytorhabdovirus trifolii; Betacytorhabdovirus viciae; Betacytorhabdovirus yerbamate; Betanucleorhabdovirus paridis; Bombyx mori latent virus; Botrexvirus duosclerotiniae; Botrexvirus unosclerotiniae; Capulavirus trifolii; Chenopodium necrosis virus; Cilevirus pistaciae; Citlodavirus apijamaicaense; Citlodavirus myricae; Cocksfoot mottle virus; Cucumber soil-borne virus; Cytorhabdovirus; Cytorhabdovirus brassicae; Cytorhabdovirus festucae; Cytorhabdovirus sonchi; Cytorhabdovirus tritici; Emaravirus clematis; Fabavirus betavitis; Fabavirus cirsii; Gammacytorhabdovirus; Gammacytorhabdovirus alphacuscutae; Gammacytorhabdovirus alphafraxini; Gammacytorhabdovirus apii; Gammacytorhabdovirus argyranthemi; Gammacytorhabdovirus betacuscutae; Gammacytorhabdovirus betafraxini; Gammacytorhabdovirus coptis; Gammacytorhabdovirus cypripedii; Gammacytorhabdovirus dauci; Gammacytorhabdovirus epipactis; Gammacytorhabdovirus gymnadeniae; Gammacytorhabdovirus heliospermae; Gammacytorhabdovirus hibisci; Gammacytorhabdovirus lonatis; Gammacytorhabdovirus lupinis; Gammacytorhabdovirus rhopalocnemis; Gammacytorhabdovirus silenis; Gammacytorhabdovirus trachyspermi; Higrevirus amurense; Higrevirus pistaciae; Ilarvirus ApNMV; Ilarvirus BabIV1; Ilarvirus TIV1; Ilarvirus ToNSV; Machlomovirus liegense; Maize aumaivirus 1; Mastrevirus bothriochloae; Mastrevirus brachypodiumprimi; Mastrevirus brachypodiumsecundi; Mastrevirus nomiae; Mastrevirus urochloareunionense; Mersevirus; Mersevirus boehmeriae; Mersevirus jujubae; Mersevirus mercurialis; Mersevirus paris; Nepovirus betaparis; Nepovirus mirae; Olpivirus lactucae; Orthotospovirus eustomae; Orthotospovirus fatsiae; Pamosaviridae; Panicum papanivirus 1; Papanivirus panici; Phragmivirus; Phragmivirus phragmii; Phragmivirus spartinae; Poinsettia mosaic virus; Potexvirus chaenostomae; Potexvirus ecsadenii; Potexvirus ecscaricae; Potexvirus ecshibisci; Potyvirus aconiti; Potyvirus alilii; Potyvirus catharanthiflavitessellati; Potyvirus crocitessellati; Potyvirus galanthi; Potyvirus parisflavitessellati; Potyvirus polygonatimaculae; Potyvirus puerariae; Ritunrivirus; Sadwavirus cattleyae; Sadwavirus chysanthemi; Sadwavirus gymnemae; Sobemovirus CFMV; Sobemovirus SMAMV; Sobemovirus smamv; Tobacco albetovirus 1; Tobacco albetovirus 2; Tobacco albetovirus 3; Tobacco virtovirus 1; Tombendovirales; Tomosaviridae; Tonesaviridae; Torradovirus arctii, Torradovirus nanorugosum; Torradovirus physalis; Trirhavirus; Trirhavirus alni; Trirhavirus chrysanthemi; Trirhavirus erysimi; Trirhavirus medicagonis; Trirhavirus picridis; Velarivirus gembloutense; Virtovirus tabaci; Waikavirus; Waikavirus ajugae; Waikavirus anacycli; Waikavirus betacamelliae; Waikavirus carotae; Waikavirus celtis; Waikavirus eleocharis; Waikavirus hirtae; Waikavirus juglandis; Waikavirus ligustici; Waikavirus

mertensiae; Waikavirus pedicularis; Waikavirus pittospori; Waikavirus populi; Waikavirus primulae; Waikavirus querci; Waikavirus ranunculi; Waikavirus thapsiae; Waikavirus thymi; Waikavirus trifoccidentale; Waikavirus violae; Weddel waterborne virus.

ICTV Taxonomy Summary Consortium: Adriaenssens E.M.; Alfenas-Zerbini P.; Aylward F.O.; Hendrickson R.C.; Hughes H.R.; Lefkowitz E.J.; Łobocka M.; Mayne R.; Mushegian A.R.; Oksanen H.M.; Penzes J.; Reyes Muñoz A.; Robertson D.L.; Roux S.; Simmonds P.; Smith D.B.; Suzuki N.; Turner D.; Van Doorslaer K..

#### Funding information

L.R. and M.I.P. acknowledge that part of this work was granted by the European Commission – NextGenerationEU, Project SUS-MIRRI.IT "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", code n. IR0000005. H.M.O. was supported by the University of Helsinki and the Research Council of Finland by funding for FINStruct and Instruct Centre FI, part of Biocenter Finland and Instruct-ERIC and Horizon MSCA 101120407. This publication was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award No. U24Al162625. This project was supported in part through a Laulima Government Solutions, LLC, prime contract with the U.S. National Institute of Allergy and Infectious Diseases (NIAID) under Contract No. HHSN272201800013C. J.H.K. performed this work as an employee of Tunnell Government Services (TGS), a subcontractor of Laulima Government Solutions, LLC, under Contract No. HHSN272201800013C. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Department of Health and Human Services or of the institutions and companies affiliated with the authors, nor does mention of trade names, commercial products or organisations imply endorsement by the U.S. Government. P.Y.T. was funded by the European Regional Development Fund (ERDF, contract REU005756) and Région Réunion through the project "Dispositif de partenariat en santé et biodiversité". S.S. acknowledges support from the Mississippi Agricultural and Forestry Experiment Station (MAFES), USDA-ARS project 58-6066-3-044, USDA-NIFA SCRI Project 1029242 and NIFA-USDA Hatch Project 7006130.

#### Acknowledgements

The authors thank Anya Crane (Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Fort Detrick, Frederick, MD, USA) for critically editing the manuscript. This is a publication of the International Committee on Taxonomy of Viruses (ICTV), a non-profit, volunteer committee of the Virology Division of the International Union of Microbiology Societies (IUMS). The ICTV does not represent or endorse the views and actions of governments or the institutions or organisations for which its co-authors work.

#### Conflicts of interest

The authors declare that there are no conflicts of interest.

#### References

- Siddell SG, Smith DB, Adriaenssens E, Alfenas-Zerbini P, Dutilh BE, et al. Virus taxonomy and the role of the International Committee on Taxonomy of Viruses (ICTV). J Gen Virol 2023;104:e001840.
- Simmonds P, Adriaenssens EM, Lefkowitz EJ, Oksanen HM, Siddell SG, et al. Changes to virus taxonomy and the ICTV Statutes ratified by the International Committee on Taxonomy of Viruses (2024). Arch Virol 2024;169:236.
- Siddell SG, Walker PJ, Lefkowitz EJ, Mushegian AR, Dutilh BE, et al. Binomial nomenclature for virus species: a consultation. Arch Virol 2020:165:519–525.
- Walker PJ, Siddell SG, Lefkowitz EJ, Mushegian AR, Adriaenssens EM, et al. Changes to virus taxonomy and to the International Code of Virus Classification and Nomenclature ratified by the International Committee on Taxonomy of Viruses (2021). Arch Virol 2021;166:2633–2648.
- Postler TS, Rubino L, Adriaenssens EM, Dutilh BE, Harrach B, et al. Guidance for creating individual and batch latinized binomial virus species names. J Gen Virol 2022;103:001800.
- Zerbini FM, Siddell SG, Mushegian AR, Walker PJ, Lefkowitz EJ, et al. Differentiating between viruses and virus species by writing their names correctly. Arch Virol 2022;167:1231–1234.
- Simmonds P, Adams MJ, Benkő M, Breitbart M, Brister JR, et al. Consensus statement: virus taxonomy in the age of metagenomics. Nat Rev Microbiol 2017;15:161–168.
- Mayne R, Simmonds P, Smith D, Adriaenssense E, Lefkowitz EJ, et al. Virus taxonomy proposal summaries: a searchable and citeable resource to disseminate virus taxonomy advances. J Gen Virol 2025;106.
- Sabanadzovic S, Abergel C, Ayllón M, Botella L, Canuti M, et al. Summary of taxonomy changes ratified by the International Committee on Taxonomy of Viruses (ICTV) from the Fungal and Protist Viruses Subcommittee. J Gen Virol 2025;106.

- Hughes HR, Ballinger MJ, Bao Y, Bejerman N, Blasdell KR, et al. Summary of taxonomy changes ratified by the International Committee on Taxonomy of Viruses (ICTV) from the Animal dsRNA and ssRNA(-) Viruses Subcommittee, 2025. J Gen Virol 2025;106.
- Teycheney P-Y, Geering ADW, Dasgupta I, Hull R, Kreuze JF, et al. ICTV Virus Taxonomy Profile: Caulimoviridae. J Gen Virol 2020:101:1025–1026.
- Geering A. Badnaviruses (Caulimoviridae). In: Bamford DH and Zuckerman M (eds). Encyclopedia of Virology, 4th Edition. 3. Oxford: Academic Press; 2021. pp. 158–168.
- 13. **Quito-Avila DF, Freitas-Astúa J, Melzer MJ**. Bluner-, cile-, and higreviruses (*Kitaviridae*). *Encyclopedia of virology* 2021:247–251.
- Ramos-González PL, Dias Arena G, Tassi AD, Chabi-Jesus C, Watanabe Kitajima E, et al. Kitaviruses: a window to atypical plant viruses causing nonsystemic diseases. Annu Rev Phytopathol 2023;61:97–118.
- 15. Muhire B, Martin DP, Brown JK, Navas-Castillo J, Moriones E, et al. A genome-wide pairwise-identity-based proposal for the classification of viruses in the genus *Mastrevirus* (family *Geminiviridae*). *Arch Virol* 2013;158:1411–1424.
- Brown JK, Zerbini FM, Navas-Castillo J, Moriones E, Ramos-Sobrinho R, et al. Revision of Begomovirus taxonomy based on pairwise sequence comparisons. Arch Virol 2015;160:1593–1619.
- 17. Reyes-Proaño EG, Cañada-Bautista MG, Cornejo-Franco JF, Alvarez-Quinto RA, Mollov D, et al. The virome of babaco (Vasconcellea × heilbornii) expands to include new members of the Rhabdoviridae and Bromoviridae. Viruses 2023;15:1380.
- Hu J, Miao T, Que K, Rahman MdS, Zhang L, et al. Identification, molecular characterization and phylogenetic analysis of a novel nucleorhabdovirus infecting Paris polyphylla var. yunnanensis. Sci Rep 2023;13:10040.
- Bejerman N, Dietzgen R, Debat H. Novel tri-segmented rhabdoviruses: a data mining expedition unveils the cryptic diversity of cytorhabdoviruses. Viruses 2023;15:2402.